0003

File ACT/007/006 #2

State of Utah
DEPARTMENT OF NATURAL RESOURCES
DIVISION OF WILDLIFE RESOURCES

Norman H. Bangerter
Governor
Dee C. Hansen
Executive Director
Timothy H. Provan
Division Director

Southeastern Region
455 West Railroad Avenue
Price, Utah 84501-2829
801-637-3310

RECEIVED
DEC 08 1989
DIVISION OF
OIL, GAS & MINING

December 5, 1989

Mr. Ira Hatch
Price District Ranger
Manti LaSal National Forest
599 West Price River Drive
Price, UT 84501

Attn: Walt Nowak

Dear Ira:

In regard to readjustment of Cypress-Western Coal Company's federal lease (4-7949) located 2/3 of a mile west of their Wattis mine, we are concerned about the effects of subsidence on wildlife. As you know, the company will access the tract from their existing Starpoint Mines. Generally speaking, impacts of subsidence on wildlife lie in two general areas: hydrologic ecosystems and terrestrial ecosystems. The following is offered for your information.

Subsidence can result in drying up of impounded water bodies or modification to flows at seeps, springs, perennial or even intermittent channels. This can result from the capture in subsidence cracks of water and its resultant migration into other geological strata. Some strata may not allow water to discharge to the surface. Such an impact can have serious consequence to a local area's wildlife in that drinking water may become reduced in value or unavailable to terrestrial animals.

Seep or springs providing flow during periods when wildlife are present represent a critical valued resource to all of the local area's wildlife. Most wildlife have small and limited home ranges. As a result, when one of these critical valued aquatic resources is lost, the animal does not have the physical capability of "packing his bag" and moving to another area of acceptable habitat. Those few species that have such a physical capability usually find the home ranges in adjoining areas already filled to capacity. It is for that reason that the Division holds firm to the philosophy that each and every seep and spring is critical resource for wildlife.

In the event that coal mining results in subsidence that impacts the flows at seeps and springs, mitigation is anticipated. An impact would be deemed substantial if the daily flow from a seep or spring was reduced by 50% or more. Mitigation that would be expected is simply the replacement of the water. Unquestionably, there would be many techniques that could achieve this goal: guzzlers are considered to be the most effective technique. They should be fenced with a 3-rail/pole fence having a maximum height of 42 inches. Clearance between the ground and the bottom pole, as well as space between the top two poles, should be at least 14 inches. This will allow passage of wildlife while disallowing domestic livestock.

Aquatic life, particularly hydrophytes, invertebrates, mollusks and fishes could suffer due to reduced or lost flows. It is even possible that subsidence fractures could expose undesirable mineral deposits to aquatic systems. If such situations were to occur, aquatic and terrestrial animals would either perish or be forced to reduce the size of their use areas. Ultimately, the carrying capacity of the area for wildlife would be reduced.

Increased flows resulting from subsidence that may appear as a benefit could, in actuality, mean that some other aquatic system has lost flows. But benefits are possible if subsidence cracks access an isolated hydrologic basin.

Reduced or lost flows in surface water systems and ground water systems can negatively impact terrestrial habitats. Mesic habitats (riparian, wetland and aspen types) associated with those systems could be degraded by the reduction or loss of water. In all ecological situations (desert, submontane and montane) riparian or wetland ecosystems due to their high level of biological productivity, limited acreage and intense use by wildlife, represent a critical valued habitat. Similar comments can be made for the aspen habitat in the montane ecological situation.

Beyond problems associated with aquatic systems, subsidence impacts to terrestrial wildlife and habitat are primarily associated with surface movement of the earth. However, methane gas has been known to travel along subsidence fractures to the surface. The escaping gas affects rhizobium in the soil and can kill adjacent rhizobium dependent vegetation.

The surface cracks alone have little impact on wildlife. It is possible that individual animals could perish in some of these holes or cracks. It has even been suggested that the movement in the earth could collapse the burrows of small rodents, thus trapping and killing them. At this point in time, I doubt whether that is a valid concern in that small rodents are extremely abundant. Since subsidence occurs over such small and limited areas, impacts to rodents would not be of consequence.

Most rodents probably have trouble with their burrows caving at times anyway. Thus they are adapted to digging around such cave-ins. It is likely that cracks and surface displacement created by subsidence represent escape cover for small animals, and to some degree, access points for burrowing animals.

Subsidence has caused escarpment failures. When raptor nests exist in the escarpments, such failures would be of negative consequence, since raptors typically return to reuse their nests over the years. Where escarpment failure occurs and there are not raptor nests, such failure could create suitable raptor nesting habitat.

Many surface displacement lines from subsidence in Utah's coal mining areas are utilized extensively by big game as travel corridors. These fracture lines, once they become filled in, represent a flat trail on which the animals can easily walk around the contours of a mountain or across ridge tops.

It is hoped that the aforementioned information will prove useful to you in coal leasing decisions. If the Division can be of any further service, please don't hesitate to call.

Sincerely,

Larry B. Dalton
Wildlife Program Manager
Resource Analysis/Habitat Protection

LBD/dd

cc: Ralph Miles, DWR
Lowell Braxton, DOGM
Clark Johnson, USFWS
Linda Seibert, BLM

UNITED STATES
DEPARTMENT OF THE INTERIOR
GEOLOGICAL SURVEY

STATE OF UTAH
UTAH GEOLOGICAL AND MINERAL SURVEY

WATTIS QUADRANGLE
UTAH
7.5 MINUTE SERIES (TOPOGRAPHIC)



Mapped, edited, and published by the Geological Survey

Control by USGS, NOS/NOAA, and U. S. Forest Service

Topography by photogrammetric methods from aerial photographs taken 1976. Field checked 1976. Map edited 1979

Projection and 10,000-foot grid ticks: Utah coordinate system, central zone (Lambert conformal conic)
1000-meter Universal Transverse Mercator grid ticks, zone 12, shown in blue. 1927 North American datum

Fine red dashed lines indicate selected fence lines

There may be private inholdings within the boundaries of the National or State reservations shown on this map

SCALE 1:24 000

CONTOUR INTERVAL 80 FEET
NATIONAL GEODETIC VERTICAL DATUM OF 1929

THIS MAP COMPLIES WITH NATIONAL MAP ACCURACY STANDARDS
FOR SALE BY U. S. GEOLOGICAL SURVEY, DENVER, COLORADO 80225, OR RESTON, VIRGINIA 22092
A FOLDER DESCRIBING TOPOGRAPHIC MAPS AND SYMBOLS IS AVAILABLE ON REQUEST

ROAD CLASSIFICATION

Primary highway, hard surface — Light-duty road, hard or improved surface
Secondary highway, hard surface — Unimproved road

Interstate Route — U. S. Route — State Route

WATTIS, UTAH

Federal Coal Lease
U-7949

Norman H. Bangerter
Governor
Dee C. Hansen
Executive Director
Timothy H. Provan
Division Director

State of Utah
DEPARTMENT OF NATURAL RESOURCES
DIVISION OF WILDLIFE RESOURCES

Southeastern Region
455 West Railroad Avenue
Price, Utah 84501-2829
801-637-3310

File Deer Creek
ACT/015/018
#2

December 5, 1989

Mr. Ira Hatch
Price District Ranger
Manti LaSal National Forest
599 West Price River Drive
Price, UT 84501

Attn: Walt Nowak

Dear Ira:

In regard to readjustment of Carrol County Coal Company's federal lease (4-7653) located on East Mountain between Meetinghouse Canyon and the Left Fork of Rilda Canyon, we are concerned about the effects of subsidence on wildlife. As you know, the company will access the tract from their existing Starpoint Mines. Generally speaking, impacts of subsidence on wildlife lie in two general areas: hydrologic ecosystems and terrestrial ecosystems. The following is offered for your information.

Subsidence can result in drying up of impounded water bodies or modification to flows at seeps, springs, perennial or even intermittent channels. This can result from the capture in subsidence cracks of water and its resultant migration into other geological strata. Some strata may not allow water to discharge to the surface. Such an impact can have serious consequence to a local area's wildlife in that drinking water may become reduced in value or unavailable to terrestrial animals.

Seep or springs providing flow during periods when wildlife are present represent a critical valued resource to all of the local area's wildlife. Most wildlife have small and limited home ranges. As a result, when one of these critical valued aquatic resources is lost, the animal does not have the physical capability of "packing his bag" and moving to another area of acceptable habitat. Those few species that have such a physical capability usually find the home ranges in adjoining areas already filled to capacity. It is for that reason that the Division holds firm to the philosophy that each and every seep and spring is critical resource for wildlife.

In the event that coal mining results in subsidence that impacts the flows at seeps and springs, mitigation is anticipated. An impact would be deemed substantial if the daily flow from a seep or spring was reduced by 50% or

more. Mitigation that would be expected is simply the replacement of the water. Unquestionably, there would be many techniques that could achieve this goal; guzzlers are considered to be the most effective technique. They should be fenced with a 3-rail/pole fence having a maximum height of 42 inches. Clearance between the ground and the bottom pole, as well as space between the top two poles, should be at least 14 inches. This will allow passage of wildlife while disallowing domestic livestock.

Aquatic life, particularly hydrophytes, invertebrates, mollusks and fishes could suffer due to reduced or lost flows. It is even possible that subsidence fractures could expose undesirable mineral deposits to aquatic systems. If such situations were to occur, aquatic and terrestrial animals would either perish or be forced to reduce the size of their use areas. Ultimately, the carrying capacity of the area for wildlife would be reduced.

Increased flows resulting from subsidence that may appear as a benefit could, in actuality, mean that some other aquatic system has lost flows. But benefits are possible if subsidence cracks access an isolated hydrologic basin.

Reduced or lost flows in surface water systems and ground water systems can negatively impact terrestrial habitats. Mesic habitats (riparian, wetland and aspen types) associated with those systems could be degraded by the reduction or loss of water. In all ecological situations (desert, submontane and montane) riparian or wetland ecosystems due to their high level of biological productivity, limited acreage and intense use by wildlife, represent a critical valued habitat. Similar comments can be made for the aspen habitat in the montane ecological situation.

Beyond problems associated with aquatic systems, subsidence impacts to terrestrial wildlife and habitat are primarily associated with surface movement of the earth. However, methane gas has been known to travel along subsidence fractures to the surface. The escaping gas affects rhizobium in the soil and can kill adjacent rhizobium dependent vegetation.

The surface cracks alone have little impact on wildlife. It is possible that individual animals could perish in some of these holes or cracks. It has even been suggested that the movement in the earth could collapse the burrows of small rodents, thus trapping and killing them. At this point in time, I doubt whether that is a valid concern in that small rodents are extremely abundant. Since subsidence occurs over such small and limited areas, impacts to rodents would not be of consequence.

Most rodents probably have trouble with their burrows caving at times anyway. Thus they are adapted to digging around such cave-ins. It is likely that cracks and surface displacement created by subsidence represent escape cover for small animals, and to some degree, access points for burrowing animals.

Subsidence has caused escarpment failures. When raptor nests exist in the escarpments, such failures would be of negative consequence, since raptors typically return to reuse their nests over the years. Where escarpment failure occurs and there are not raptor nests, such failure could create suitable raptor nesting habitat.

Many surface displacement lines from subsidence in Utah's coal mining areas are utilized extensively by big game as travel corridors. These fracture lines, once they become filled in, represent a flat trail on which the animals can easily walk around the contours of a mountain or across ridge tops.

It is hoped that the aforementioned information will prove useful to you in coal leasing decisions. If the Division can be of any further service, please don't hesitate to call.

Sincerely,

Larry B. Dalton
Wildlife Program Manager
Resource Analysis/Habitat Protection

LBD/dd

cc: Ralph Miles, DWR
Lowell Braxton, DOGM
Clark Johnson, USFWS
Linda Seibert, BLM

UNITED STATES
DEPARTMENT OF THE INTERIOR
GEOLOGICAL SURVEY

STATE OF UTAH
UTAH GEOLOGICAL AND MINERAL SURVEY

RILDA CANYON QUADRANGLE
UTAH
7.5 MINUTE SERIES (TOPOGRAPHIC)



Mapped, edited, and published by the Geological Survey

SCALE 1:24 000

CONTOUR INTERVAL 80 FEET
NATIONAL GEODETIC VERTICAL DATUM OF 1929

THIS MAP COMPLIES WITH NATIONAL MAP ACCURACY STANDARDS
FOR SALE BY U.S. GEOLOGICAL SURVEY, DENVER, COLORADO 80225, OR RESTON, VIRGINIA 22092
A FOLDER DESCRIBING TOPOGRAPHIC MAPS AND SYMBOLS IS AVAILABLE ON REQUEST

ROAD CLASSIFICATION

RILDA CANYON, UTAH

1979

Federal Coal Lease U-7653