

**NOTICE OF INTENT TO CONDUCT
MINOR COAL EXPLORATION
2015**

**SCHOOL AND INSTITUTIONAL TRUST LANDS
ADMINISTRATION MUDDY TRACT
AND UTAH STATE COAL LEASE
ML 49443-OBA**

CANYON FUEL COMPANY, LLC

FEB. 2015



**Canyon Fuel
Company, LLC**

A Subsidiary of Bowie Resource Holdings, LLC

C/041/002 Incoming

#4804

SUFco Mine

Mark Bunnell
Geologist
597 S SR 24
Salina, Utah 84654
(435) 286-4403
Fax (435) 286-4499

Feb. 24, 2015

Daron Haddock
Permit Supervisor
Utah Coal Regulatory Program
Division of Oil, Gas & Mining
1594 West North Temple, Suite 1210
PO Box 145801
Salt Lake City, Utah 84114-5801

RECEIVED

FEB 25 2015

DIV. OF OIL, GAS & MINING

Re: NOI, Minor Coal Exploration, 2015, SITLA Muddy Tract and Utah State Coal Lease ML 49443-OBA, Canyon Fuel Co. LLC, Sufco Mine, C/041/002

Dear Daron:

Attached are 5 copies of a Canyon Fuel Company Minor Coal Exploration NOI for proposed exploration drilling north of Sufco Mine on the SITLA Muddy Tract, including our leased portion of that tract.

The proposed drilling includes 5 drill holes, all of which are located on Manti-La Sal National Forest land and SITLA coal ownership.

If possible we would like to start work by mid-June.

Attached with the NOI is the C1/C2 form. Copies are also inserted into each of the NOI binders. If you or your staff have questions please don't hesitate to contact me as follows: Sufco Mine 435-286-4403; cell 435-650-8940; or email at mbunnell@bowieresources.com.

Sincerely,

Mark Bunnell
Senior Geologist
Canyon Fuel Company, LLC

Encl.
MDB:mdb

APPLICATION FOR COAL PERMIT PROCESSING

Permit Change [] New Permit [] Renewal [] Exploration [x] Bond Release [] Transfer []

Applicant: Canyon Fuel Company, LLC

Applicant: SUFCO

Permit Number: C/041/002

Title: NOTICE OF INTENT TO CONDUCT MINOR COAL EXPLORATION -- SITLA MUDDY TRACT AND UTAH STATE COAL LEASE ML49443-OBA, 2015

Description, Include reason for application and timing required to implement:

FIVE COAL EXPLORATION BOREHOLES TO BE DRILLED BEGINNING MID JUNE, 2015

Instructions: If you answer yes to any of the first eight (gray) questions, this application may require Public Notice publication.

- 1. Change in the size of the Permit Area? Acres: _____ Disturbed Area: _____ [] increase [] decrease.
2. Is the application submitted as a result of a Division Order? DO# _____
3. Does the application include operations outside a previously identified Cumulative Hydrologic Impact Area?
4. Does the application include operations in hydrologic basins other than as currently approved?
5. Does the application result from cancellation, reduction or increase of insurance or reclamation bond?
6. Does the application require or include public notice publication?
7. Does the application require or include ownership, control, right-of-entry, or compliance information?
8. Is proposed activity within 100 feet of a public road or cemetery or 300 feet of an occupied dwelling?
9. Is the application submitted as a result of a Violation? NOV # _____
10. Is the application submitted as a result of other laws or regulations or policies?
11. Does the application affect the surface landowner or change the post mining land use?
12. Does the application require or include underground design or mine sequence and timing? (Modification of R2P2)
13. Does the application require or include collection and reporting of any baseline information?
14. Could the application have any effect on wildlife or vegetation outside the current disturbed area?
15. Does the application require or include soil removal, storage or placement?
16. Does the application require or include vegetation monitoring, removal or revegetation activities?
17. Does the application require or include construction, modification, or removal of surface facilities?
18. Does the application require or include water monitoring, sediment or drainage control measures?
19. Does the application require or include certified designs, maps or calculation?
20. Does the application require or include subsidence control or monitoring?
21. Have reclamation costs for bonding been provided?
22. Does the application involve a perennial stream, a stream buffer zone or discharges to a stream?
23. Does the application affect permits issued by other agencies or permits issued to other entities?

Please attach four (4) review copies of the application. If the mine is on or adjacent to Forest Service land please submit five (5) copies, thank you. (These numbers include a copy for the Price Field Office)

I hereby certify that I am a responsible official of the applicant and that the information contained in this application is true and correct to the best of my information and belief in all respects with the laws of Utah in reference to commitments, undertakings, and obligations, herein.

Mark Bunnell
Print Name

Mark Bunnell, Geologist
Sign Name, Position, Date Feb 24 2015

Subscribed and sworn to before me this 24 day of February, 2015

Jacquelyn Nebeker
Notary Public

My commission Expires: _____, 20____
Attest: State of _____ } ss:
County of _____



RECEIVED

FEB 25 2015

DIV. OF OIL, GAS & MINING

For Office Use Only:	Assigned Tracking Number:	Received by Oil, Gas & Mining RECEIVED FEB 25 2015 DIV. OF OIL, GAS & MINING
-----------------------------	----------------------------------	---

Form DOGM- C1 (Revised March 12, 2002)

**NOTICE OF INTENT TO CONDUCT
MINOR COAL EXPLORATION
2015**

**SCHOOL AND INSTITUTIONAL TRUST LANDS
ADMINISTRATION MUDDY TRACT
AND UTAH STATE COAL LEASE
ML 49443-OBA**

CANYON FUEL COMPANY, LLC

FEB. 2015

INTRODUCTION

Canyon Fuel Co. Company is submitting this Notice of Intent to Conduct Minor Coal Exploration on behalf of Canyon Fuel Company, LLC, Sufco Mine to the Utah Division of Oil, Gas, and Mining (UDOGM) in order to obtain approval to conduct coal exploration and reclamation activities in the summer field seasons of 2015 thru 2019. The type of exploration proposed is wireline core drilling. Up to 5 holes will be drilled on the School and Institutional Trust Lands Administration (SITLA) Muddy Coal Tract. The holes are located within and immediately north of Canyon Fuel Company's leased portion that tract. All exploration activities will occur on U.S. Forest Service, Manti-LaSal National Forest surface and will involve both leased and unleased coal resources administered by the School and Institutional Trust Lands Administration.

The planned project will utilize heli-portable drilling equipment and will require no road or drill pad construction. This application is formatted to address the specific requirements of R645-201-200. The U.S. Forest Service/BLM typical exploration stipulations (utilized during 2014 projects) are addressed in Appendix A. Other related information is given in Appendix B through E (submitted for confidential filing). Five copies of this notice are submitted for distribution by the UDOGM to other agencies.

R645-201 Coal Exploration: Requirements for Exploration Approval

The proposed exploration plan qualifies as minor exploration as described in the State of Utah Coal Mining Rules R645 section R645-201-200.

R645-201-221

The name, address and telephone number of the applicant are:

Canyon Fuel Company, LLC
C/o SUFCO Mine
597 S. SR 24
Salina, Utah 84654 (work) 435-286-4403

The applicant is the same as the operator of the proposed exploration plan. Correspondence regarding this exploration plan should be addressed to:

Mark Bunnell
Canyon Fuel Company, LLC
C/o SUFCO Mine
597 S. SR 24
Salina, Utah 84654 (work) 435-286-4403

R645-201-222

The name, address and telephone number of the representative of the applicant who will be present during and be responsible for conducting the exploration is:

Mark Bunnell
Canyon Fuel Company
C/o SUFCO Mine
597 S. SR 24
Salina, Utah 84654
(work) 435-286-4403 (cell) 435-650-8940

At times a consulting geologist may act as representative of the applicant. The UDOGM and USFS will be notified of the consulting geologist's name and address if one is used.

R645-201-223

The exploration area is generally located in central Utah 10 miles northwest of Emery (Map 1). The legal description of the SITLA Muddy Coal Tract is as follows:

T. 20 S., R. 5 E., Salt Lake Base and Meridian
Sec. 32, S1/2, S1/2
Sec. 33, S1/2, S1/2

T.21 S., R. 5 E., Salt Lake Base and Meridian
Sec. 4, all
Sec. 5, all
Sec. 7, all
Sec. 8, all
Sec. 9, all


Containing 2,553.84 acres


The tract is entirely located in Sevier County, Utah. Map 2 shows proposed borehole locations and Forest Development Roads that will be used to access the exploration area. Holes A-15 and C-15 are located on an unleased portion of the SITLA Muddy Tract. Holes B-15, D-15, and M-15 are located on the leased portion of the tract (Utah State Coal Lease ML 49443-OBA).

Surface topography includes a large ridge (known as Big Ridge) with small drainages on both sides. Elevations range from 8,400 to 9,100 ft. The major drainages in the area are Cowboy Creek and Muddy Creek. The holes are located on the north side of the ridge (Map 2).

Rocks exposed in the exploration area belong to the Cretaceous Blackhawk and Price River Formations, as well as the Cretaceous-Tertiary North Horn Formation. Rock types are predominantly sandstones, siltstones, shale and coal. The major geologic feature in the exploration area is the escarpment created by the outcrop of the 200 ft. thick Castlegate Sandstone member of the Price River Formation in Box Canyon.

Vegetation in the exploration area is comprised mostly of conifer timber, aspen, and sage communities with some mountain brush (oak, serviceberry, mahogany). Hole locations are generally in grass and sagebrush areas. The streams in the exploration area are not considered habitat for the four endangered Colorado River drainage fish species and none are known to occur in the drainages. The streams are not capable of supporting game fish. The exploration area is important habitat for raptors, elk, mule deer, cougar, bobcat, black bear, and small mammals. The area is habitat for a limited number of reptiles and amphibians.





R645-201-225

Threatened, endangered, or special interest species in the exploration area include the sage grouse, goshawk, bald eagle and peregrine falcon. Exploration and reclamation activities will not occur within one half mile of known breeding and nesting areas during breeding or nesting periods. A site specific raptor survey will also be conducted prior to commencement of drilling operations. Information concerning threatened and endangered species is included in the Pines Tract Project Final Environmental Impact Statement, Jan. 1999. The U.S. Forest Service also completed wildlife and BEBA analyses of the Muddy Tract area which were included in the U.S.F.S. decision document relative to Canyon Fuel Co. Company's approved 2004 Muddy Drilling project. The 2004 Wildlife Resources Report and BEBA reports are attached as Appendix B and C. Also included in Appendix B is the "Muddy Technical Report: Wildlife" prepared for the Muddy Tract EIS by the USFS. Canyon Fuel Company's annual raptor survey was also completed in the area in June, 2014 (Appendix D, confidential file). The 2015 survey is planned for approximately the same time frame and will be forwarded to the Division upon completion. No Mexican Spotted owls are known to occur in the area. Appendix D also includes the two-year Mexican Spotted Owl Survey, 2002 & 2003 Final Report prepared for the USFS as part of the Muddy Tract EIS.

There are no known districts, sites, buildings, structures, or objects listed on, or eligible for listing on, the National Register of Historic Places in the proposed exploration area. There are known archeological resources located in the proposed exploration area. A cultural resource evaluation of previous exploratory drilling locations on the Muddy Tract and nearby Pines Lease was submitted to the Division in November, 1999 (Utah State Project No. UT-99-AF-0684f). The Pines Tract Project Final Environmental Impact Statement (Jan., 1999) also includes cultural resource analysis on the area. A site-specific cultural resource evaluation has been completed for this project (U-15-EZ-0056f) and is included in Appendix E (confidential file).

R645-201-224

A timetable for each year's exploration-related activities is given below. It is anticipated that exploration activities will start in early June of each year and be completed by late October. This timetable may vary somewhat depending on U.S. Forest Service stipulations and other factors such as weather. It should be noted that holes may be completed as part of larger drilling projects on adjacent Federal coal and lands.

EVENT	WEEK 1	WEEK 2	WEEK 3	WEEK 4	
Set pumps, frac tanks and run water lines to sites					
Move drill equipment to site and drilling					
Reclaim any disturbance					

R645-201-225

The general method to be followed during drill hole exploration, reclamation, and abandonment is: 1) place temporary water system including lines, portable pumps, and portable water tanks 2) prep the drill site which will include minor leveling and brush removal with hand tools, 3) drill and log holes, and 4) reclaim/seed drill site remove all waterlines, tanks, and pumps. No blasting will be done for road building or repair. Repair of Forest System Roads may include hauling of gravel to fill rough areas on bedrock ledges and grading rutted areas. U.S. Forest Service road use permits will be obtained prior to initiation of the project. Equipment operators will use pick-up trucks for transportation.

Drilling will be accomplished utilizing plug and core drilling techniques. Drilling will involve up to two heli-portable wireline core rigs capable of drilling 2200 ft. with necessary heli-portable support equipment such as rod trays, supply tubs, supply trailers, portable mud tanks, etc. Drill water will be hauled by water truck on Forest system roads to one or two 18,000 gal. frac tanks. From there, water will be pumped to each drill site (Map 2). The drilling procedure will be to rotary/plug drill to core depth and continuously core through the coal zones to total depth utilizing water, polymer, and/or mud as drilling medium.

Water for drilling and road watering will be obtained from the North and/or South Forks of Quitcupah Creek (locations shown on Map 2) and hauled to 18,000 gal. frac tanks located near the intersection of Forest roads 52101 and 50044 (Map 2). A Triplex pump or equivalent will be located at the roadside tank site for pumping water to the drill sites. Water will be pumped via 1, 2, and 3 inch HDPE waterline as well as 2 inch high pressure hose. If necessary, due to inadequate water flow in those drainages, additional water may be hauled from the Sufco minesite. Water will generally be hauled in 4000 gal. water trucks. All necessary arrangements will be made for water usage, including a Temporary Water Change approval from the Division of Water Rights. Copies of approved Temporary Water Changes will be forwarded to the Division prior to project startup. Where not located adjacent to a road, waterlines and pumps will be placed and removed via horseback, on foot, or helicopter.

Supply trailers will carry drill steels, coring equipment, drilling additives, cutting and welding equipment, and other supplies to the staging area/helicopter LZ. Also located at the staging area

will be a double-lined diesel fuel tank for the drill rig and support equipment, an 18,000 gal. frac tank and pump, construction equipment, and other supplies such as waterline, etc. Support vehicles such as pickup trucks and a geophysical logging truck will be parked at the staging area. Access to the staging area will be via U.S. Forest Service roads 007 and 044 (Map 2).

The only coal to be removed during exploration activities will be cores. Cores will nominally be 1.5 inches (BQ) in diameter. Given an approximate projected thickness of 7 ft. for the Upper Hiawatha seam and 12 ft. for the Lower Hiawatha seam, approximately 20 to 50 lbs. of coal will be removed.

No temporary road construction is planned for this project.

To eliminate the need for road and drillpad construction, the planned drilling method is helicopter-supported wireline plug and core drilling.

Core drilling will involve up to two skid-mounted 2200 ft rated core drill, one or two 1000 gal. poly water tanks at the drill rig, two water trough-type mudtanks, and 4 to 6 drill rod trays. Other support equipment will include 4000 gallon water trucks to supply the 18,000 gal. frac tanks at staging areas, one 1500 gallon water/fire truck, two to four 18,000 gallon frac tanks at staging areas depending on water usage, two to three supply trailers parked at staging areas, up to 8 pick-up trucks, and a geophysical logging truck. One hundred to 200 ft. of surface casing will be set in each hole depending on hole conditions. Fifth-wheel supply trailers or transport trailers will carry the heli-portable equipment, including drills, drill steels, coring equipment, drilling additives, cutting and welding equipment, and other supplies to the staging area. One pick-up truck will be used for each drill rig by the drillers to carry personnel, fuel, and supplies and two to three pickup trucks will be used by the dirt contractor. The logging contractor will use a single axle 1 ton rated truck. The company representative and geological consultant will also use pick-up trucks for transportation.

Backup and auxiliary equipment to be located at the staging areas will include but not be limited to one or two 18,000 gal. frac (water) tank(s), two to three supply trailers, drilling fluid containers, two 4000 gal. water trucks, a fuel truck or or double lined fuel tank, four to 8 pickup trucks, a covered tool supply trailer, and a geophysical logging truck.

Regulations cited in R645-202-232 relative to roads will be followed. Disturbance to wildlife will be minimized by utilizing helicopter-supported drilling methods. A site specific raptor survey has will be completed and submitted to the Division and the USFS prior to project startup. No wetlands or riparian will be disturbed by drilling activities. Existing roads will be maintained during the project by grading as needed under an approved USFS road use permit. Spot placement of gravel may be necessary depending on weather conditions and USFS stipulations. No utility or support facilities are present in the area.

Drill site reclamation will include filling in and returning any disturbed topsoil to original contour with hand tools. The surface will be reseeded with the USFS-approved seed mix. No damage to public or private property will occur.

No drillpad construction will occur. Some minor topsoil leveling and removing of brush will occur to allow placement of the low-impact heli-portable drill stand and rod trays. All work will be done with hand tools. The drill engine and hydraulics will be underlain with brattice cloth to eliminate the chance of any petroleum products such as diesel fuel or hydraulic oil dripping or spilling onto topsoil. Drill sites will be approx.. 12 ft. X 30 ft. (figure 1). No drilling fluids, oil and grease, or diesel fuel will be allowed to contact the topsoil. Mudpits will be pumped out and/or allowed to dry before being reclaimed. Pumped fluids will be transported to an approved disposal site off USFS lands. Mudpits will be fenced when unattended to prevent wildlife from possible entry.

Reclamation will occur as soon as possible upon completion of drilling operations. Reclamation will include filling in any hand excavations and reseeded the disturbed surface with the approved seed mix. No damage to public or private property will occur.

The drill location will be setup approximately as shown on Figure 1. Earth excavation for the drill site will be minimal using hand tools only. Some minor leveling for placement of wood crib blocking for leveling of drill may be required. Minor amounts of topsoil that may be removed will be stored and replaced upon completion of drilling. No mud pits will be excavated. Portable mudtanks will be utilized. Cuttings will be stored and hauled away to the Skyline Mine waste rock site by helicopter or truck upon completion of drilling.

Reclamation is an integral part of the exploration activities and will progress as contemporaneously as practical with the other exploration activities. Upon completion of the hole, all hand excavations will be filled in to original contour, topsoil replaced, all equipment will be removed, and all trash will be hauled away. An approved seed mix will then be applied to the drill area.

There will be no diversion of overland flows.

It is not anticipated that acid- or toxic- forming materials will be encountered during exploration because none have been encountered previously. Samples of drill core will be analyzed for acid- and toxic-forming materials. These samples will be taken from the 10 ft. interval above and below each seam of minable thickness.




Figure 1. Typical drillpad construction and setup.

The method of revegetation is intended to encourage prompt revegetation and recovery of a diverse, effective, and permanent vegetative cover. The following seed mix has been proscribed by the U.S. Forest Service for reclamation of nearby 2014 drill holes (the seed mix proscribed by the USFS in 2015 will be utilized):

Seed Mix

		<u>Pounds PLS/acre</u>
Western Wheatgrass	<i>Elymus smithii</i>	2
Basin Wild Ryegrass	<i>Elymus cinereus</i>	1
Intermediate Wheatgrass	<i>Elymus hispidus</i>	2
Blue Leaf Aster	<i>Aster glaucodes</i>	0.25
Lewis Flax	<i>Linum lewisii</i>	0.50
Small Burnet	<i>Sanguisorbia minor</i>	1
Silvery Lupine	<i>Lupinus argenteus</i>	1
True Mahogany	<i>Cercocarpus montanus</i>	1
Bitterbrush	<i>Purshia tridentata</i>	1
	TOTAL	9.75

The pure live seed (PLS) rating will be 99% and only seed meeting the State Seed Act will be used. Certification tags will be retained by the permittee. The vegetative cover resulting from this seed mix is considered capable of stabilizing the soil surface from erosion.

Map 2 shows the location of the proposed drill sites and the Forest Service roads used for access. Equipment access to the exploration area will be via FDR 007 which traverses federal coal leases SLI-062583, U-47080, and U-63214 on both the Fishlake and Manti-LaSal National Forests then FDR 044 to the proposed tank/pump location and staging area/landing zone location.

The main drill hole diameter will be nominally 2.36 inch diameter. Approximately 200 to 300 ft. of surface casing (3 inch) will be set. Estimated depth and other drill hole information is given in the following table. Disturbed area will include one drill site. Total disturbed area acreage is estimated at 0.030 acres.

Drill Site	Location	Total Depth (ft)	Disturbance (ft)	Disturbed Area (acres)
A-15	NW, NW, 7, T21S, R5E	2100	8 X 30	0.006
B-15	NW,NW, 8, T21S, R5E	1900	8 X 30	0.006
C-15	SE,SE, 32, T20S, R5E	1200	8 X 30	0.006
M-15	NE,NW, 4, T21S, R5E	1600	8 X 30	0.006
D-15	NE,NE, 4, T21S, R5E	1100	8 X 30	0.006
			TOTAL	0.030

There are no occupied dwellings or pipelines located in the exploration area. A shallow trench and hole will be dug with hand tools near the drill hole to allow drill fluid to accumulate and be recirculated into the mud tank. The trench and hole will be lined with brattice or pit liner to eliminate contamination to the topsoil. No structures will be constructed nor debris disposed of in the exploration area. The permittee or his representative will have a copy of this Notice of Intention To Conduct Minor Coal Exploration while in the exploration area available for review by an authorized representative of the Division upon request.

Upon completion of drilling, the holes will be plugged with a cement, bentonite, or cement/bentonite slurry to full depth. A brass tag will be placed at the top of the drill hole stating the operator's name, drill hole number, and legal description. The tag will be placed in cement at ground level. One or more of the proposed drill holes may be completed as water monitor wells. The completion method includes: 1) plug the hole with cement, bentonite, or cement/bentonite slurry from full depth to the planned monitor completion depth, 2) clean the hole of drill cuttings by circulating with air or water, 3) insert an approx. 1 inch diameter steel or PVC casing with a 10 to 30 ft section of 0.010" slot screen section with an end cap, 4) fill the annulus in the screened section with washed sand or pea gravel, 5) pack off the screened section or seal it off with bentonite, and 5) fill the remainder of the hole annulus to the surface

with a cement slurry. A steel protective casing with locking cap will be placed 1 to 2 ft. above ground level. Figure 2 gives the typical design to be used in completing the piezometer.

Holes left as water monitor wells will be plugged as above when no longer in use, and any surface standpipe removed.




Figure 2. Typical shallow water monitor well installation.

R645-203-200

Canyon Fuel Company requests the Division not make any drilling information available for public inspection relative to coal seam thickness or quality. This information is considered crucial to Canyon Fuel Company's competitive rights.

R645-202.230

No adverse impacts to stream channels will occur during water pumping or drilling activities. The BLM, USFS and the Division will be notified as to points of diversion. Stream flows will not be pumped dry during pumping activities. In the past, the BLM and USFS have authorized the placement of a water tank at the pump location to allow more gradual pumping and water storage. If, due to drought conditions, stream flow drops too low to pump, water will be hauled from the Sufco minesite. No water will be pumped from Quitchapah Creek without an approved "Temporary Change of Water" from the Division of Water Rights. Approved Temporary Change documents will be forwarded to the Division and USFS prior to startup of drilling operations. It is projected that approx. 0.2 acre-ft. of water will be utilized during the project during any given year. Total water usage is estimated at 0.6 acre-ft.

R645-202-231

A cultural resource survey was conducted in 2014 (U-15-EZ-0056f) and is included in Appendix E (confidential file). If additional survey work is required by the U.S. Forest Service, it will be conducted in late spring 2015 and forwarded to the Division prior to startup of drilling activities. Copies of additional recent cultural resource surveys in the area are included in Appendix E (confidential). Threatened, endangered, and sensitive plant and animal survey information has been developed by the U.S.F.S. during their work relative to Canyon Fuel's 2004 Muddy Tract drilling license application (Appendix B and C). The USFS also conducted baseline studies for the Muddy Tract EIS. Appendix D (confidential) shows the raptor information developed during the SUFCO's 2014 raptor survey. SUFCO's 2015 raptor survey will be conducted in early summer 2015. That survey together with site specific survey information will be forwarded to the Division prior to startup of drilling activities.

TES protection measures include the use of non-surface disturbing water pumping equipment which will minimize surface disturbance. Pumping of most project water through waterlines will minimize water truck traffic on permanent and temporary access roads.

R645-202-232

No access road or trail construction is planned for this project.

R645-202-235 (R645-301-624.210, R645-301-731.121, R645-301-731.215))

Geologic logs of drilling will be kept. Any appreciable water encountered during drilling will be logged, noting depth, geology, and estimated flow. Due to the use of low-impact heli-portable drilling equipment, only shallow water monitor well installation is possible. One or more of the holes may be completed as shallow monitor wells as shown in section R645-201-225 (pg. 13).

Figure 1 shows a drawing of the approximate drill site setup. If the drill hole begins to make excess water, such water will be pumped to a larger 18,000 gal. frac tank at the water tank staging area. From there it will be hauled to an approved waste water disposal site off Forest lands. At no time will excess drill water generated in the drill hole be allowed to run on topsoil on the surface.

It should be noted that fresh incoming water from the water tank/pump supply system has to be circulated through the engine cooling system at all times during drill operation. At times, excess fresh cooling water will be allowed to drain onto the surface. Care will be taken to ensure this fresh water is controlled such that it is dispersed and not allowed to create undo erosion of the topsoil.

R645-301-525-200

No major utilities pass over, under, or through the exploration area. Use of roads and development of the exploration site will not disrupt or damage any utility service.

R645-301-527.230

U.S. Forest system roads utilized as part of this minor coal exploration plan will be maintained as per U.S.F.S. authorized road use permit, including proper control of fugitive dust to minimize effects to fish, wildlife, and related environmental values. Approved road use permits will be obtained prior to startup of the drilling project.

R645-301-731.100

Approved Temporary Changes of Water for water to be used in the drilling process will be obtained prior to startup of drilling activities.

R645-301-742.410 thru 742.420

Minimal surface disturbance with hand tools will be required for the drilling project. Disturbance will be limited to the drill site. Changes to drainage patterns will be minimal and temporary. As shown on Figure 1, the drilling equipment at the drill site will be set on top of existing topsoil with only minor leveling with hand tools beneath leveling stands, drill rod stands and mud tanks. Brattice or pit liner will be anchored beneath the drill to catch any oil leaks or drill fluid spills that may occur. Absorbent pads will be utilized to soak up any leaks on the brattice or pit liner material. Brattice and/or pit liner material will also be laid in any minor hand-excavated ditches and the small hand-excavated mud-pit to prevent drilling fluid from contacting topsoil. No perennial or intermittent stream drainages will be crossed. Excess water will be removed and placed in the drill water tank for use in the drilling process or pumped to a tank at the staging area for collection and transport to an approved waste water disposal site. Contributions of suspended solids during any pumping activity from the stream will be minimized by placing native stream gravel or rocks beneath the screens and suction hoses wherever they can potentially stir up sediment at pump-pickup points.

The potential for water pollution will be minimized by keeping pollutants away from the drill hole and in their containers. Materials used during drilling operations will be selected to be as non-polluting as possible. All spills of polluting materials will be removed from the area and properly disposed of.

No mixing of surface and ground waters is possible because all drill sites will be above perennial and ephemeral stream drainages.

Drill fluids and/or cuttings will be contained within mudtanks. If necessary, excess fluids will be pumped out and excess drill cuttings and core will be hauled off Forest Service land and disposed

of properly.

SUFCO will retain all drill and geophysical logs. Copies of the drilling and geophysical logs will be provided to the School and Institutional Trust Lands Administration.

APPENDIX A

The surface management agency is the U.S. Forest Service, Manti-La Sal and Fishlake National Forests. This Appendix gives responses to applicable Special Coal Lease Stipulations and all the Standardized Stipulations for Coal Drilling Operations from the Manti-La Sal National Forest Land and Resource Management Plan (1986). Other Forest Service requirements are addressed at the end.

SPECIAL COAL LEASE STIPULATIONS

1. Before undertaking activities that may disturb the surface of previously undisturbed leased lands, the Lessee may be required to conduct a cultural resource inventory and a paleontological appraisal of the areas to be disturbed. These studies shall be conducted by qualified professional cultural resource specialists or qualified paleontologists, as appropriate, and a report prepared itemizing the findings. A plan will then be submitted making recommendations for the protection of, or measures to be taken to mitigate impacts, for identified cultural or paleontological resources.

If cultural resources or paleontological remains (fossils) of significant scientific interest are discovered during operations under this lease, the Lessee, prior to disturbance shall immediately bring them to the attention of the appropriate authority. Paleontological remains of significant scientific interest do not include leaves, ferns, or dinosaur tracks commonly encountered during underground mining operations.

The cost of conducting the inventory, preparing reports, and carrying out mitigating measures shall be borne by the Lessor.

Response/Action – If required, Canyon Fuel Co. will have a cultural resource inventory and paleontological appraisal done by a third party contractor in the spring/early summer, 2015. The work will be done by a qualified contractor approved by the USFS. A plan will be submitted with recommendations for the protection of, or measures to be taken to mitigate impacts, should any cultural or paleontological resources be identified. A cultural resource inventory was conducted for the drill sites in 2014 (U-15-EZ-0056f).

2. If there is reason to believe that Threatened or Endangered (T&E) species of plants or animals, or migratory bird species of high Federal interest occur in the area, the Lessee shall be required to conduct an intensive field inventory of the area to be disturbed and/or impacted. The inventory shall be conducted by a qualified specialist and a report of findings will be prepared. A plan will be prepared making recommendations for the protection of these species or action necessary to mitigate the disturbance. The cost of conducting the inventory, preparing reports, and carrying out mitigating measures shall be borne by the Lessor.

Response/Action – If required, Canyon Fuel Co. will have field inventories done of threatened or endangered fauna and flora by a third party contractor in the spring/early summer of 2015. The work will be done by qualified contractors approved by the USFS. A plan will be submitted with

recommendations for the protection of, or measures to be taken to mitigate impacts, should any threatened or endangered fauna or flora be found. Canyon Fuel Co. will bear the cost of conducting the inventories, preparing reports, and carrying out any mitigating measures.

4. If removal of timber is required for clearing of construction sites, etc., such timber shall be removed in accordance with the regulations of the surface management agency.

Response/Action – Canyon Fuel Co. does not plan to remove timber.

5. Existing Forest Service owned or permitted surface improvements will need to be protected, restored, or replaced to provide for the continuance of current land uses.

Response/Action – Canyon Fuel Co. will improve the surface of Forest Development Roads as needed following the recommendations of the USFS and according to the USFS Road Use Permit. A final grading of any ruts or other damage will be made after all activities are completed. Stock watering ponds will not be used and otherwise will not be affected by the exploration and reclamation activities.

6. In order to protect big-game wintering areas, elk calving and deer-fawning areas, sage grouse strutting areas, and other key wildlife habitat and/or activities, specific surface uses outside the mine development area may be curtailed during specified periods of the year.

Response/Action – Canyon Fuel Co. will not conduct exploration and reclamation activities during periods specified by the USFS to protect wildlife habitat and/or activities.

STANDARDIZED STIPULATIONS FOR COAL DRILLING OPERATIONS

Stipulations To Be Included in the Coal Drilling Permit

1. A pre-work meeting including the responsible company representative(s), contractors, and the Forest Service will be conducted at the project location prior to commencement of operations. Site-specific Forest Service requirements will be discussed at this time.

Response/Action – Canyon Fuel Co. will hold a pre-work meeting with the responsible company representative(s), contractors, and the Forest Service at the project location prior to commencement of operations. Site-specific Forest Service requirements will be discussed at this time.

2. A Road-Use Permit must be obtained from the Forest Service before equipment is transported onto National Forest System lands.

Response/Action – Canyon Fuel Co. will obtain a Road Use Permit from the Forest Service before equipment is transported onto National Forest System lands.

3. The Forest Service will specify times and locations, if any, that exploration activities may not occur.

Response/Action – Canyon Fuel Co. will obey any times and locations specified by the Forest

Service in which exploration activities may not occur.

4. All surface disturbing activities including reclamation will be supervised by a responsible representative of the permittee/licensee who is aware of the terms and conditions of the project permits and licenses. A copy of the appropriate permits and licenses must be available for review at the project site.

Response/Action – Canyon Fuel Co. will have an employee or representative who is aware of the terms and conditions of the project permits and licenses present during surface disturbing activities including reclamation.

5. The Forest Service must be notified 48 hours in advance that heavy equipment will be moved onto National Forest System lands and that surface disturbing activities will commence.

Response/Action – Canyon Fuel Co. will notify the Forest Service at least 48 hours in advance of the movement of heavy equipment onto National Forest System lands and of when surface disturbing activities will commence.

6. Establishment of campsites and staging areas on National Forest System lands in support of this project is subject to Forest Service approval.

Response/Action – Canyon Fuel Co. will request Forest service approval before establishing campsites or staging areas for this exploration project.

7. The Forest Service will be notified of any proposed alterations to the plan of operations. Any changes to the existing plan are subject to Forest Service review and concurrence.

Response/Action – Canyon Fuel Co. will notify the Forest Service of any proposed alterations to the plan of operations and will not make alterations to the plan until the Forest Service has reviewed and concurred with the change.

8. Fire suppression equipment will be available to all personnel working at the project site. Equipment will include at least one hand tool per crew member consisting of shovels and pulaskis and one properly rated fire extinguisher per vehicle and/or internal combustion engine.

Response/Action – Canyon Fuel Co. or its contractors will provide fire suppression to all personnel working at the project site, including at least one hand tool per crew member consisting of shovels and pulaskis and one properly rated fire extinguisher per vehicle and/or internal combustion engine.

9. All gasoline, diesel and steam-powered equipment will be equipped with effective spark arresters and mufflers. Spark arresters will meet Forest Service specifications discussed in the USDA Forest Service Spark Arrester Guide, June, 1981. In addition, all electrical equipment will be properly insulated to prevent sparks.

Response/Action - Canyon Fuel Co. will ensure that all gasoline, diesel and steam-powered equipment is equipped with effective spark arresters and mufflers meeting USDA Forest Service Spark Arrester Guide, June, 1981. Also, all electrical equipment will be properly insulated to prevent sparks.

10. The permittee/licensee will be held responsible for damage and suppression costs for fires started as a result of operations. Fires will be reported to the Forest Service as soon as possible.

Response/Action - Canyon Fuel Co. understands that it will be held responsible for damage and suppression costs for fires started as a result of operations. Fires will be reported to the Forest Service as soon as possible.

11. The Forest Service reserves the right to suspend operations during periods of high fire potential.

Response/Action - Canyon Fuel Co. will, at Forest Service request, suspend operations during periods of high fire potential.

12. Water needed in support of operations will be properly and legally obtained according to State Water Laws. The locations of diversion, if on National Forest System lands, are subject to Forest Service review and approval.

Response/Action - Canyon Fuel Co. will properly and legally obtain the water needed in support of operations according to State Water Laws. Any diversions, if on National Forest System lands, will be reviewed and approved by the Forest Service. Water for exploration and reclamation activities will be taken from either the South Fork and/or the North Fork of Quitcupah Creek where they flow under Forest Development Road FDR 007; or from the Sufco minesite. It is estimated that 0.2 acre-feet will be used for drilling and dust suppression per year and approx. 0.5 acre-feet for the entire project.

13. There will be no unauthorized off-road vehicular travel.

Response/Action - Canyon Fuel Co. will ensure that no unauthorized off-road vehicular travel occurs.

14. Section corners or other survey markers, including claim corners, in the project area will be located and flagged for preservation prior to commencement of surface disturbing activities. The removal, displacement, or disturbance of markers will be approved by the proper authority.

Response/Action - Canyon Fuel Co. will locate and flag any section corners, survey markers, and claim corners that are in areas of planned surface disturbance for preservation prior to commencement of surface disturbing activities. Any removal, displacement, or disturbance of markers will be approved by the proper authority.

15. If cultural or paleontological resources are discovered during operations, all operations which may result in disturbance to the resource will cease and the Forest Service will be notified of the discovery.

Response/Action - Canyon Fuel Co. will cease all operations should cultural or paleontological resources be discovered and the Forest Service will be notified of the discovery.

Forest Service Stipulations To Be Discussed at the Pre-Work Meeting

1. Gates will be kept closed unless otherwise notified.

Response/Action - Canyon Fuel Co. will keep all gates closed unless otherwise notified.

2. The permittee/licensee will be held responsible for all damages to fences, cattle guards, resource improvements, roads, and other structures on National Forest system lands which result from operations. The Forest Service will be notified of damages as soon as possible.

Response/Action - Canyon Fuel Co. acknowledges responsibility for all damages to fences, cattle guards, resource improvements, roads, and other structures on National Forest system lands which result from operations. The Forest Service will be notified of damages as soon as possible.

3. All drilling fluids, muds, and cuttings will be contained on the project site in mud pits or portable containers. The pits will not be used for disposal of garbage, trash or other refuse.

Response/Action - Canyon Fuel Co. will contain all drilling fluids, muds, and cuttings in heli-portable containers to be transported to an approved site. An MSDS will be provided the Forest Service for all drilling fluids and muds used. Garbage, trash or other refuse will be properly disposed of off Forest Service lands.

4. All trees and brush must be cleared as the first step for new access and site construction. Topsoil must be stripped and stockpiled at a location where loss and contamination is minimized.

Response/Action – Minimal trees or brush will be cleared with hand tools only. No topsoil removal will be necessary due to the use of heli-portable equipment. Topsoil will be protected by brattice or pit liner material to prevent contamination by oil or drilling fluids.

5. Disturbed areas must be reclaimed by the end of the field season. Exceptions require Forest Service approval.

Response/Action - The disturbed area will be reclaimed as soon as possible upon completion of drilling. The drill site will be reclaimed by filling in any holes or leveled areas to approx. original contour with hand tools. The soil will be raked and grubbed material will be distributed over the area and the area will be seeded. The estimated acreage disturbed by drilling is given in the table in the section discussing R645-201-225.

6. Contaminated soil and gravel must be stripped and placed in the mud pit prior to site reclamation.

Response/Action - Contaminated soil will be removed and transported to an approved disposal site.

7. Mud pits must be allowed to dry before they are backfilled and reclaimed.

Response/Action – The small mud pit that may be utilized for recirculation of fluids will be allowed to dry before backfilling or reclamation with hand tools.

8. When dry, mud pits must be reclaimed by selectively back filling excavated materials, topsoil last, such that the disturbed area is replaced to approximate original contour. The disturbed area must be seeded with the specified seed mix when topsoil is replaced.

Response/Action – Mud pit will be reclaimed as specified and the site will be seeded with specified seed mix.

9. Roads to be obliterated must be reclaimed by ripping the surface, replacing the disturbed area to approximate original contour, replacing stockpiled topsoil, and seeding with the specified seed mix. Seeding must take place when topsoil is replaced. Water diversion structures, if needed, must be constructed as specified by the Forest Service.

Response/Action – No roads will be constructed.

10. All disturbed drainages must be replaced to their approximate original configuration when the project area is reclaimed.

Response/Action – It is expected that no drainages will be disturbed. If disturbance does occur, drainages will be replaced to their approximate original configuration.

11. All significant water encountered during drilling must be reported to the Forest Service, including the depth and formation at which it was encountered, and an estimate of flow.

Response/Action - All significant groundwater encountered during drilling will be reported to the Forest Service, including the depth and formation at which it was encountered, and an estimate of flow.

12. The operator must clean-up and remove all drilling equipment, trash, garbage, flagging, vehicles and other such materials from National Forest System lands.

Response/Action - The permittee will clean-up all drilling equipment, trash, garbage, flagging, vehicles, and other such materials from National Forest System lands upon completion of exploration and reclamation operations.

13. All trash, garbage, and other refuse must be properly contained on the project site prior to disposal.

Response/Action - The permittee will properly contain all trash, garbage, and other refuse on the project site prior to disposal.

14. All drill holes must be plugged in accordance with Federal and State regulations.

Response/Action - Canyon Fuel Co. Company will plug all drill holes in accordance with Federal and State regulations.

15. Operations must be coordinated with grazing permittees to prevent conflicts.

Response/Action - Exploration and reclamation operations will be coordinated with grazing permittees to prevent conflicts.

16. Harassment of wildlife and livestock is prohibited.

Response/Action - Canyon Fuel Co. will ensure that harassment of wildlife and livestock does not occur.

Stipulations To Be Included in Road-Use Permits

1. Roads must not be used when they are wet and susceptible to damage.

Response/Action - Roads will not be used when they are wet and susceptible to damage.

2. The permittee is responsible for repair of damages to roads caused by its operations.

Response/Action - Canyon Fuel Co. will repair damage to roads caused its operations.

3. All traffic must maintain speeds commensurate with existing conditions.

Response/Action - Canyon Fuel Co. will ensure that all traffic related to exploration and reclamation operations maintain speeds commensurate with existing conditions.

4. Roads must be watered if dust becomes a problem or if excessive loss of material occurs.

Response/Action - Roads will be watered if dust becomes a problem or if excessive loss of material occurs.

5. Heavy equipment will not be transported along FDR 007 and FDR 044 during holiday weekends and the opening weekend of the regular big game hunting seasons.

Response/Action - Canyon Fuel Co. will not transport heavy equipment on Forest

Development Roads at times specified by the Forest Service.

Other Forest Service Requirements

1. Outside berms will not be constructed on any roads.
2. The permittee will notify the USFS District Ranger of any artesian groundwater flows encountered prior to plugging the drill hole to determine whether or not the Forest Service would elect to establish a permanent water development at the site.
3. The permittee will be responsible for control of noxious weed infestations found to be a result of this exploration. All equipment coming into the exploration area from outside Utah will be steam cleaned to remove foreign seeds.
4. No timber will be removed during exploration and reclamation.

Stipulation for Lands of the National Forest System Under Jurisdiction of the Department of Agriculture

1. The licensee/permittee/lessee must comply with all the rules and regulations of the Secretary of Agriculture set forth at Title 36, Chapter II, of the Code of Federal Regulations governing the use and management of the National Forest System (NFS) when not inconsistent with the rights granted by the Secretary of the Interior in the license/permit/lease. The Secretary of Agriculture's rules and regulations must be complied with for (1) all use and occupancy of the NFS prior to approval of a permit/operation by the Secretary of the Interior, (2) uses of all existing improvements, such as Forest Development roads, within and outside the area licensed, permitted or leased by the Secretary of the Interior, and (3) use and occupancy of the NFS not authorized by a permit/operating plan approved by the Secretary of the Interior.

APPENDIX B

**WILDLIFE RESOURCES REPORT
SUFCO 2004 COAL
EXPLORATION DRILLING
PROJECT
&
MUDDY CREEK TECHNICAL REPORT
WILDLIFE**

WILDLIFE RESOURCES REPORT
for the
**SUFCO 2004 HELICOPTER-ASSISTED
COAL EXPLORATION DRILLING PROJECT**

Ferron/Price Ranger District
Manti-La Sal National Forest
San Pete and Sevier Counties, Utah

Prepared by: /s/ Terry Nelson **Date:** 7/26/2004
Terry Nelson, Wildlife Biologist

Reviewed by: /s/ Pamela Jewkes **Date:** 7/27/2004
Pamela Jewkes, Forest Fisheries Biologist

I. INTRODUCTION

The purpose of this Wildlife Resources Report is to assess the potential affects of the proposed SUFCO 2004 Helicopter-Assisted Coal Exploration Drilling Project on wildlife species listed as threatened, endangered, candidate or proposed by the U.S. Fish and Wildlife Service (Service); wildlife species listed on the Intermountain Regional Forester's list of sensitive species; species identified as Management Indicator Species (MIS) by the Manti-La Sal National Forest; and migratory bird species identified as priority species by the Utah Partners in Flight Avian Conservation Strategy (2002).

A. PROPOSED ACTION

1. Summary of the Proposed Action

Canyon Fuel Co. Company has submitted a plan to conduct coal exploration and reclamation activities. Six drill holes are proposed for coal exploration during summer 2004. Five of the holes are proposed on unleased federal portions of the proposed Muddy Coal Area (Forest Service Surface/Federal Coal). One hole is proposed on Utah School and Institutional Trust Lands Administration (SITLA) portions of the Muddy Coal tract (Forest Service Surface/SITLA Coal). The project would be completed during the summer and early fall season, 2004. Access to three of the proposed drill sites would be along existing FS roads. Helicopters would be used to fly drill equipment to the other 3 remote sites where there are no existing roads. Since, helicopter-drilling techniques are proposed, there would be minimum disturbance (<100 ft² per site).

The proponent's proposed action as defined in its 2003 coal exploration license proposal is to access National Forest system lands, construct temporary drilling pads, drill holes to acquire needed geologic data from six coal exploration holes and reclaim disturbed areas on Forest Service managed land, using helicopter-assisted drilling methods. The proposed helicopter-assisted drilling project is outlined below:

- The planned drilling method is wireline core drilling from the surface down through to the lowest coal horizon. Equipment will include two heli-portable skid-mounted core drilling rigs together with all necessary equipment such as drill rod trays, fuel tanks, water tanks, etc. The necessary equipment and vehicles include an 18,000 gallon frac tank, helicopter, jet fuel tank (trailer mounted), 4000 gallon water truck, two or three fifth-wheel flatbed trucks trailers used to haul drill equipment, four pick-up trucks, a covered tool supply trailer, and a geophysical logging truck.
- Hauling exploration equipment and transporting personnel to the staging area (see map) would be via frFDR 50007, 50044, and 50132 which traverses both the

Fishlake and Manti-La Sal National Forests. Road-use permits would be obtained from the Forest Service before operation start.


- Site preparation would include removal of some vegetation with hand tools as needed for placement of the drill rig and needed equipment. Surface disturbance would be minimal; less than 100 square feet per site.
- The finished size of the hole will be nominally 2 3/16 inch diameter. Three-inch surface casing will be inserted through the surface alluvium and certain other intervals depending on hole conditions. Upon completion, holes would be geophysically logged.
- Soils would be protected from potential contamination by placement of brattice or similar impermeable material placed beneath mechanical equipment
- Water for drilling operations and road maintenance would be obtained from Muddy Creek and/or Quitchumpah Creek. Necessary arrangements would be made with shareholders and the Utah Division of Water Rights through a temporary water exchange permit. Completed drill holes would be plugged with a cement or cement/bentonite slurry to their full depth in accordance with BLM and Forest Service standards.
- Reclamation would include removal of equipment and trash immediately after hole completion. Topsoil would be scarified with hand tools. The disturbed areas would be reseeded (same as 2003 seed mix) with seed mix approved by the FS. The total plan, including reclamation, should be completed in 8 to 10 weeks.
- One hole may be completed as water monitoring well. Nominal 1.0 to 1.5 inch well screen and steel casing would be installed to below the deepest mineable coal seam. The screen zone would be sand packed and sealed from overlying strata and the overlying hole annulus would be cemented to the surface. Well casing with a locking lid would be left at the surface extending above the surface approximately two feet. The wellhead would be properly identified with either a brass marker or a welded-on identification. Once the monitor well is no longer in use, it would be completely plugged with a cement or cement/bentonite slurry to the top. The wellhead would be removed at the surface.

2. Description of the Project Location

The general locations are in San Pete and Sevier Counties about 10 miles northwest of the town of Emery, Utah. The proposed project area and drill hole locations are shown on Map 1. The proposed drill holes, lease tract administrator, location, depth and proposed access routes are summarized in the following table:

Drill Site	Tract	Location: T20S, R5E	Access Route
A	SITLA	SW, SE, Sec. 32	By Air FR 50044
B	BLM	NW, NW, Sec. 33	By Air FR 50132
C	BLM	SE, SW, Sec. 29	FR 50132

D	BLM	NW, NW, Sec. 32	FR 50132
E	BLM	NE, SE, Sec. 29	FR 50132
F	BLM	SE, NW, Sec. 29	By Air FR 50132



B. SPECIES OF CONCERN

1. Threatened and Endangered (T&E) Wildlife Species

Endangered species are species that have been identified, and listed in the Federal Register, by the U.S. Fish and Wildlife Service (Service) as being in danger of extinction throughout all or a significant portion of its range. Threatened species are species that have been identified, and listed in the Federal Register, by the Service as likely to become an endangered species within the foreseeable future throughout all or a significant portion of its range.

Table 1 lists wildlife and fish species designated as threatened or endangered (T&E) by the Service that could occur in San Pete or Sevier County, Utah. T&E species that could occur in San Pete or Sevier County but do not have suitable habitat in and are not likely to occur in or near the proposed project area are also identified in Table 1, and will not be considered further in this wildlife Resources Report. There are no fish species identified as a threatened, endangered or candidate species for San Pete or Sevier County, and there are no proposed wildlife or fish species identified for San Pete or Sevier County.

Table 1. Listed and candidate wildlife and fish species that could occur in San Pete or Sevier County, Utah, and their potential for occurrence in the proposed project area and consideration in this Wildlife Resources Report.

SPECIES	SPECIES STATUS	SPECIES OCCURRENCE IN THE PROJECT AREAS AND CONSIDERATION IN THIS WILDLIFE REPORT
Bald Eagle <i>Haliaeetus leucocephalus</i>	Threatened San Pete and Sevier Counties	Considered. A bald eagle pair has been known to nest in Emery County approximately 20 miles northeast of the proposed project area. Bald eagles may occur incidentally in the proposed project area.
Western Yellow-billed Cuckoo <i>Coccyzus americanus occidentalis</i>	Candidate San Pete and Sevier Counties	Not Considered. The western yellow-billed cuckoo breeds in western U.S. states including Utah, and migrates to South America during winter. Cuckoos are riparian obligates. Nesting habitat is classified as dense lowland cottonwood/willow riparian forest characterized by a dense sub-canopy or shrub layer. In Utah, nesting habitats are found at elevations between 2,500 to 6,000 feet. They appear to require large tracts (100 to 200 acres) of contiguous riparian nesting habitat (Parrish et al. 1999). The proposed project is located in fairly dry pinyon/juniper, sagebrush, mohogany habitats at between 8,500 and 9,000 ft. elevation, there is no suitable habitat for this species in or near the project area.
Canada Lynx <i>Lynx canadensis</i>	Threatened San Pete County	Not Considered. The proposed project is located in open fairly dry pinyon/juniper, sagebrush, mohogany habitats, which does not provide suitable habitat for the Canada lynx.
Utah Prairie Dog <i>Cynomys parvidens</i>	Threatened San Pete and Sevier Counties	Not Considered. Utah prairie dogs are found in areas where there are deep, well-drained soils, burrows extend straight down for about 10-15 ft. and then branch into horizontal tunnels. They feed on insects (particularly cicadas), where available. Their preferred vegetative food type is alfalfa, but they generally prefer grasses over forbs and shrubs. Moist palatable forage must be available throughout the summer. The proposed project is located in fairly dry pinyon/juniper, sagebrush, mohogany habitats with mostly shallow soils over Castle Gate sandstone. No evidence of Utah prairie dogs was found in or near the project area.

2. Sensitive Wildlife and Fish Species

Sensitive species are species that are recognized by the Regional Forester as needing special management attention in order to prevent them from becoming threatened or endangered.

Table 2 lists the Intermountain Regional Forester’s list of sensitive wildlife species that could occur on the Manti Division of the Manti-La Sal National Forest (MLNF). Sensitive wildlife species that do not occur or have suitable habitat in or near the proposed project area are identified in Table 2 and will not be considered further in this Wildlife Resources Report.

Table 2. Sensitive wildlife and fish species that could occur on the Manti Division of the MLNF, and their potential occurrence in the proposed project area and consideration in this Wildlife Report.

SPECIES	SPECIES OCCURRENCE IN THE PROJECT AREAS AND CONSIDERATION IN THIS WILDLIFE REPORT
Spotted Bat <i>Euderma maculatum</i>	Considered. In Utah, the spotted bat likely occurs throughout the state. It is known to use a variety of vegetation types from approximately 2,500 to 9,500 feet, including riparian, desert shrub, ponderosa pine, montane forests, open pastures and meadows. Spotted bats roost alone in rock crevices high up on steep cliff faces. There are potentially suitable roosting cliffs near the proposed project area. Spotted bats may occasionally forage in the sagebrush/shrub habitat in the vicinity of the proposed project, and in the nearby ponderosa pine habitat.
Townsend’s Big-eared Bat <i>Plecotus townsendii pallescens</i>	Considered. In Utah, Townsend’s big-eared bats roost and hibernate in caves and mines, they also roost (but not hibernate) in buildings (Oliver 2000). These bats use juniper/pine forests, shrub/steppe grasslands, deciduous and mixed conifer forests. There is potentially suitable roost sites and forage habitat in or near the proposed project area.
Greater Sage Grouse <i>Centrocercus urophasianus</i>	Considered. Sage grouse are generally found where there are large tracts of sage brush habitat with a diverse and substantial understory of native grasses and forbs or in areas where there is a mosaic of sagebrush, grasslands, aspen. Wet meadows, springs, seeps, or other green areas within sagebrush shrublands are generally needed for the early brood-rearing period. There is suitable breeding habitat near the proposed project area.
Northern Goshawk <i>Accipiter gentilis</i>	Not Considered. Goshawks forage in fairly dense (generally greater than 40 percent canopy cover) conifer forests, and they nest in even denser stands (generally greater than 60 percent canopy cover), many nest and forage sights contain an aspen component. The proposed project is located in fairly dry pinyon/juniper, sagebrush, mohogany habitats. There is no suitable goshawk habitat in or near the project area.
Peregrine Falcon <i>Falco peregrinus</i>	Considered. Peregrine falcons may travel more than 18 miles from the nest site to hunt for food, however average foraging distance from the eyrie extends out to 10 miles, with 80 percent of peregrine falcon foraging occurring within a mile of the nest. The nearest known peregrine falcon eyrie is located approximately 3 ½ miles from the project area. Nesting peregrine falcons may forage in the vicinity of the proposed project.
Flammulated Owl <i>Otis flammeollus</i>	Not Considered. Flammulated owls prefer mature ponderosa pine/Douglas fir forests with open canopies, but they can be found in second growth ponderosa pine, aspen and mixed conifer forests that contain a ponderosa pine component. The proposed project is located in fairly dry pinyon/juniper, sagebrush, mohogany habitats, and will not alter or disturb flammulated owl habitat.
Three-toed woodpecker <i>Picoides tridactylus</i>	Not Considered. Three-toed woodpeckers are found in northern coniferous and mixed forest types up to 9,000 feet elevation. Forests containing spruce, grand fir, ponderosa pine, tamarack, and lodgepole pine are used. Nests may be found in spruce, tamarack, pine, cedar, and aspen trees. The proposed project is located in fairly dry pinyon/juniper, sagebrush, mohogany habitats, and will not alter or disturb three-toed woodpecker habitat.
Spotted Frog <i>Rana pretiosa</i>	Not Considered. Spotted frogs are most commonly found in cold, still, permanent water in such habitats as marshy edges of ponds or lakes, in algae-grown overflow pools of streams, and near flat water springs with emergent vegetation. The spotted frog may move considerable distances from water after breeding, often frequenting mixed conifer and subalpine forests, grasslands, and brushlands of sage and rabbitbrush. No spotted frogs have been found on the Manti – La Sal National Forest, and they are not known or thought to occur on the Forest.
Colorado Cutthroat Trout <i>Oncorhynchus clarki pleuriticus</i>	Not Considered. This species is generally limited to small headwater streams in remote areas where other trout species have not been introduced. They historically occurred in most waters of the upper Colorado River basin. No populations were discovered during 1992 Utah Department of Wildlife Resources surveys on the Ferron/Price district, however a non-pure population was recently found in Crandall Canyon. The proposed project would not impact streams known or suspected to contain Colorado cutthroats.

**Bonneville
Cutthroat Trout
*Oncorhynchus
clarki utah***

Not Considered. This trout requires cool, clear, well-oxygenated water and the presence of clean, well-sorted gravels with minimal fine sediments for successful spawning (Lentsch et al. 1997) There are no streams in the proposed project area that would provide suitable habitat for this species, and the project would not impact streams known or suspected to contain Bonneville cutthroats

3. Management Indicator Species (MIS)

Table 3 lists wildlife species identified as Management Indicator Species (MIS) by the Manti-La Sal National Forest (MLNF) that could occur on the Manti Division of the MLNF. MIS species that do not occur or have suitable habitat in or near the proposed project area are identified in Table 3 and will not be considered further in this Wildlife Resources Report.

Table 3. Management Indicator Species that could occur on the Manti Division of the Manti-La Sal National Forest.

Species Common name (Scientific name)	Species/Habitat Associations	Consideration of this species
Rocky Mountain Elk <i>Cervus canadensis</i>	Elk tend to occupy the higher elevation aspen and mixed conifer habitats from spring through early fall, and move to lower elevation mixed shrub, pinyon/juniper, and sagebrush habitats for winter	Considered. Elk are known to use the area during snow free months.
Mule Deer <i>Odocoilus hemionus</i>	Mule deer use most of the habitat types surrounding the proposed project area. Lower elevation pinyon/juniper and sagebrush habitats provide suitable winter range. Most mule deer winter range is located at the edge of National Forest system lands on BLM managed land.	Considered. Mule deer are found in and around the proposed project area.
Northern Goshawk <i>Accipiter gentilis</i>	Goshawks forage in fairly dense (generally greater than 40 percent canopy cover) conifer forests, and they nest in even denser stands (generally greater than 60 percent canopy cover). In Utah, many nest stands contain an aspen component.	Not Considered. The proposed project is located in fairly dry pinyon/ juniper, sagebrush, mohogany habitats. There is no suitable goshawk habitat in or near the project area.
Golden Eagle <i>Aquila chrysaetos</i>	Golden eagles are generally found in mountainous or hilly terrain, but also inhabit valleys and plains, especially during migration and winter. They generally nest on cliffs, however tree nests are not uncommon. They hunt over open country for small mammals, snakes, birds and carrion.	Considered. There are a number of golden eagle nest sites located within 2 miles of the proposed project area.
Macroinvertebrates (aquatic Insects)	Macroinvertebrates (aquatic insects) are ecological indicator species in aquatic habitats. Habitat requirements for aquatic macroinvertebrates vary with species, habitat requirements for any one species are very specific so macroinvertebrate indices can provide an indication of general stream health.	Not Considered. The proposed project is located in fairly dry pinyon/juniper, sagebrush, mohogany habitats; the project will not alter or disturb aquatic macroinvertebrate habitat.

4. Migratory Birds

Migratory bird conventions impose obligations on federal agencies for the conservation of migratory birds and their habitats. The Migratory Bird Treaty Act has implemented these conventions with respect to the United States, and Executive Order 13186 ensures that environmental analyses of Federal actions required by the NEPA or other established environmental review processes evaluate the effects of actions on migratory birds, with emphasis on species of concern.

The Utah Partners in Flight Avian Conservation Strategy identifies 20 non-game migratory land birds as priority species. Eleven of these species could be expected to occur on the Ferron/Price Ranger District of the Manti-La Sal National Forest. Table 4 lists these species, their habitat associations, and their consideration in the document.

Table 4. Neotropical migratory birds (NTMBs) listed as priority species by the Utah Partners in Flight Avian Conservation Strategy that could occur on the Manti Division of the Manti-La Sal National Forest.

Common name (Scientific name)	Species/Habitat Associations	Consideration of this species
Virginia's Warbler (<i>Vermivora virginiae</i>)	Preferred breeding habitat includes chaparral and open stands of pinyon/juniper, ponderosa pine and scrub oak, mountain mahogany thickets or other low brushy habitats on dry mountainsides. In Utah, the primary breeding habitat is oak, and secondary breeding habitat is pinyon/juniper at elevations ranging from 4,000 to 10,000 ft (Parrish et al 2002)	Considered. Virginia's warblers are known to occur on the Ferron/Price Ranger District of the Manti-La Sal NF, but they are not known to nest here
Gray Vireo (<i>Vireo vicinior</i>)	Preferred breeding habitat is on arid slopes dominated by mature pinyon/juniper woodlands. This species commonly occurs in suitable habitats in Colorado, Nevada and Arizona at elevations ranging from 3200 ft to 6800 ft, and they are known to nest in southwest Utah north to Sevier County. Gray vireos are not believed to nest on the Manti Division of the Manti-La Sal NF, but occur at lower elevations in Emery County, Utah (Walters and Sorenson 1983)	Not Considered. The proposed drill sites are located at 8,500 feet elevation and above, which is above the elevation range of the gray vireo
Bell's Vireo (<i>Vireo bellii arizonae</i>)	Preferred nesting habitat in Utah is cottonwood-willow dominated riparian areas. This species breeds in southwestern Utah in the Virgin River drainage, Zion NP, and Beaver Dam Wash (Wauer 1997). Bell's vireos are not known to nest on the Manti Division of the Manti-La Sal NF	Not Considered. The proposed project area does not contain suitable riparian nesting habitat for this species
Black Rosy-Finch (<i>Leucosticte atrata</i>)	Breeds above timberline in Alpine tundra using barren, rocky or grassy areas and cliffs among glaciers or at bases of snow fields. In Utah, the largest breeding populations occur in alpine habitats in the Wasatch and Uinta Mountains	Not Considered. The proposed project is located in sub-alpine habitats below the elevation breeding range of the black-rosy finch
Brewer's Sparrow (<i>Spizella breweri breweri</i>)	Breeding habitat is primarily shrubsteppe, but may also breed in high desert scrub (greasewood) habitats. Breeding habitats are usually dominated by big sagebrush (Parrish et al 2002)	Considered. The sage brush habitat surrounding the proposed project sites may provide suitable breeding habitat for the Brewer's sparrow
Black Swift (<i>Cypseloides niger</i>)	Black swifts nest in small colonies near and often behind waterfalls at elevations ranging from 6,000 ft to 11,500 ft (Parrish et al 2002). There are only 2 confirmed breeding locations Utah: the Bridal Veil Falls area and Aspen Grove area (Knorr 1962)	Not Considered. Black swifts have been seen on the Manti Division of the Manti-La Sal NF. However, the proposed project area does not contain suitable nesting habitat for this species
Broad-tailed Hummingbird (<i>Selasphorus platycercus</i>)	In Utah, the primary breeding habitat is lowland riparian. They have also been recorded as breeding in mountain riparian, aspen, ponderosa pine, Engelmann spruce, subalpine fir, and Douglas fir (Calder and Calder 1992). Nesting typically occurs at elevations ranging from 6,000 to 8,000 ft near streamside habitat	Not Considered. The proposed project area does not provide suitable breeding habitat for this species
Ferruginous Hawk (<i>Buteo regalis</i>)	Usually breeds in areas of flat and rolling terrain in grassland or shrub steppe habitat. Avoids high elevations, forest and narrow canyons. Occurs in grasslands, agricultural lands, sagebrush/saltbrush/greasewood shrub lands and the periphery of pinyon/juniper habitats	Not Considered. Ferruginous hawks are not likely to occur in the high elevation project area
Yellow-billed Cuckoo (<i>Coccyzus americanus</i>)	In Utah, the yellow-billed cuckoo is a rare breeder in large tracts (100-200 acres) of contiguous dense lowland riparian habitats. Over the last 10 years, there are only 3 breeding records in the state, none on the Manti Division of the Manti-La Sal NF (Parrish et al 2002)	Not Considered. There are no large tracts of riparian habitat in or near the proposed project area

**Black-throated
Gray Warbler**
(*Dendroica
nigrescens*)

Preferred breeding habitat includes dry oak slopes, pinyon, juniper, pinyon/juniper woodlands, open mixed woods, and dry coniferous and mixed conifer habitats with brushy understories, and in chapparal. It occurs from sea level up to 5400 ft elevation.

Not Considered. The proposed project is located above 8,000 feet elevation, which is above the elevation range of the black-throated gray warbler.

Sage Sparrow
(*Amphispiza belli
nevadensis*)

Uncommon permanent resident in Utah, occurs up to 8,000 ft elevation. Nests have been found in rabbitbrush, hopsage, saltbush, and big sage.

Considered. The sage brush habitat surrounding the proposed project sites may provide suitable breeding habitat for the sage sparrow.

II. TES, MIS and PRIORITY MIGRATORY BIRD SPECIES POTENTIALLY AFFECTED by the PROPOSED PROJECT

A. THREATENED AND ENDANGERED SPECIES

Bald Eagle

Bald eagle nests are typically located in multi-storied (uneven aged) coniferous forest stands that contain elements of old growth structure, and are located near bodies of water that support prey species. Nest trees are generally one of the largest trees in the stand, which provides good visibility and a clear flight path to and from the nest (Stalmaster 1987). Bald Eagles typically construct large, conspicuous stick nests in sizeable trees.

Prey species commonly include fish, waterfowl, jackrabbits, and carrion; results of food-habit studies have indicated that bald eagle diets included: 56 percent fish, 28 percent birds, 14 percent mammals, and 2 percent miscellaneous sources (Stalmaster 1987).

Bald eagles spend over 90 percent of the daylight hours perching. Important perch sites generally have 3 fundamental elements: a direct view of potential food sources, located within 50 meters of water, and are located in areas isolated from human disturbance (Stalmaster 1987).

Unlike nesting and perch sites, roosting sites are not necessarily located close to water; during breeding season, nesting adults often roost in the nest or at the nest tree (Stalmaster 1987). Roost sites generally provide thermal cover, and are isolated from human disturbance. Bald eagles often roost communally during winter.

During the winter, Bald Eagles tend to concentrate wherever food is available; food availability is probably the single most important factor affecting winter eagle distribution and abundance, but availability of night roosts and diurnal perches are also fundamental elements of bald eagle winter range. Eagles are often attracted to wintering concentrations of waterfowl. In some regions, such as Utah, carrion can also be an important food source. At wintering areas, Bald Eagles often roost in large groups.

These communal roosts are located in forested stands that provide protection from harsh weather.

There are only a few known nesting pairs of bald eagles in Utah. There is a bald eagle nest site located approximately 20 miles from the proposed project area, and located approximately 7 miles from Forest Service managed land. A nesting pair had been observed at this site during the nesting and fledgling period for several years prior to 1997. This nesting territory was not occupied in 2001 or 2002. The nest was blown out of the tree in the winter of 2003, and a pair built a new nest approximately ½ mile southeast of the old one, but did not nest successfully in 2003. The pair worked on the nest again in early 2004, but did not nest. A 1997 study by N. Boschen indicated that the pair did not forage on national forest system lands; nesting adults and fledglings were found to forage within a 5 mile radius of the nest tree (Boschen, 1997). No bald eagles are known to nest on Manti-La Sal NF managed lands. Most bald eagle sightings on the Forest have been at Joe's Valley Reservoir and Huntington Canyon during late fall and early winter prior to freeze over.

B. SENSITIVE SPECIES

Spotted Bat

The spotted bat ranges from Mexico through the western states to the southern border of British Columbia; it is probably widely distributed in low numbers throughout western North America (Toone 1994). And it probably occurs throughout Utah, but its distribution appears to be patchy. Hasenyager (1980) thought that "the range of the spotted bat in Utah could incorporate the southern third of the state and central portions of the west desert where suitable roosts exist, excluding the higher portions of the central mountain range." Habitat occupied by this bat ranges from low desert to montane coniferous forests normally below 8,000 feet in elevation (Watkins 1977). They have been found in a variety of habitat types including open ponderosa pine, desert shrub, pinyon/juniper, and open pasture and hay fields. In Utah, the spotted bat has been captured in several habitats: lowland riparian habitat (open meadows), desert shrub communities (sagebrush/rabbitbrush), ponderosa pine forest, montane grassland (grass/aspen), and montane forest and woodland (grass/spruce/aspen). This species has also been occasionally found in or on buildings in Utah towns and cities (Oliver 2000).

They typically roost singly in crevices in steep cliff faces. Cracks and crevices in limestone or sandstone cliffs provide important roosting sites (Spahr et al. 1991), especially where rocky cliffs occur in proximity to riparian areas. Day roosts and maternal roosts are typically within small (up to 6 cm) cracks and crevices in cliff faces (Toone 1994). The relative inaccessibility of cliff roosts may insulate spotted bats from human disturbance, but the species has been observed roosting (and foraging) near campgrounds (Toone 1994). Spotted bats are thought to feed mainly on moths high above the vegetation canopy. They forage alone after dark using echolocation, which is effective for fast flight feeding on tympanate moths (moths that can detect ultra-sonic

sounds). As is common with many bats, spotted bats may forage a considerable distance (up to 6 miles) from roost sites (Toone 1994).

Roosting habitat in the Wasatch Plateau region is likely to occur in numerous cliffs along the edges of the plateau and on canyon walls that cut through the plateau. It is likely that spotted bats forage in a variety of habitats on the Plateau that are located within 6 miles of suitable roost cliffs and at elevations lower than 9,500 ft. Various surveys on the MLNF have detected spotted bats in several major canyons (and their tributaries) on the east side of the plateau, including Muddy, Ferron, Straight, Cottonwood, and Huntington Canyons (Perkins and Peterson 1997, and Sherwin et al. 1997). These surveys also detected spotted bats near Joes Valley Reservoir and Trail Mountain.

Observations made during the 1997 surveys on the MLNF indicated that spotted bats tolerate at least moderate human disturbance while foraging. Surveys were conducted at several sites near roads with light to moderate vehicular traffic (Crandall Canyon, Huntington Canyon, Straight Canyon), including tandem coal trucks. Spotted bats were observed foraging at low elevation sites, within 30 meters of the right-of-way. The fact that spotted bats were relatively common in active and previously mined areas may imply that subsidence caused cliff failures have not dramatically affected resident populations (Sherwin, et al. 1997).

Townsend's Big-eared Bat

Townsend's big-eared bats occur throughout North America, from British Columbia to southern Mexico; from California to South Dakota and western Texas and Oklahoma. They are widely distributed throughout the Intermountain Region, and they occur throughout Utah (Oliver 2000). They inhabit a wide variety of xeric and mesic habitats including: desert scrub, sagebrush, chaparral, deciduous and coniferous forests including, but not limited to pinyon/juniper, ponderosa pine, spruce/fir, redwood, mixed hardwood/conifer, and oak woodlands (Pierson et al. 1999), and their distribution is strongly correlated with the availability of caves or cave-like roosting habitat such as mines, buildings with cave-like attics, diversion tunnels or bridges (Pierson et al. 1999). They require relatively spacious, relatively cool cave-like roost sites; generally at least 30 meters in length, and at least 2 meters high with temperatures ranging from -2.0 to 13.0° C (Pierson et al. 1999).

These bats are relatively sedentary, and do not migrate long distances; generally seasonal movements are less than 32 km (Pierson et al. 1999). Detections in Utah have ranged from 3,300 feet to 9,520 feet (Oliver 2000). In Utah, night roosts are found in mines and caves; day roosts and maternity roosts are found in mines, caves and buildings (Oliver 2000).

Townsend's big-eared bats are insectivorous; a lepidopteran specialist eating mostly moths (Pierson et al. 1999). They forage after dark using echolocation on the wing (Sphar et al. 1991); a late flyer, emerging from the roost primarily after dark; well after sunset (Pierson et al. 1999).

Breeding occurs at winter sites between October and February, and parturition occurs in late spring and early summer. Each female usually gives birth to a single offspring. Females and young roost in communal nurseries, which range in size from 12 to 200 individuals. The offspring fly at three weeks and are weaned in six to eight weeks. Nurseries break up by August.

During winter, these bats roost singly or in small clusters in hibernacula from October to February. They don't migrate, but will move to different roost locations within hibernacula and may even move to different hibernacula during a winter in response to temperature changes.

Most of the bat surveys conducted on the MLNF that employed the use of mist nets or bat detectors have not revealed Townsend's big-eared bats (Perkins and Peterson 1997, and Sherwin et al. 1997). This is not unusual, as these bats are most commonly located during direct surveys of roosts (Oliver 2000).

There is potentially suitable Townsend's big-eared bat foraging habitat in and around the proposed project area.

Greater Sage Grouse

Sage grouse are sagebrush ecosystem obligates; they occur in mosaics of sagebrush, grasslands, and aspen, and are associated with both tall and short species of sagebrush in foothills, sagebrush shrublands, and mountain slopes. They do not occur in pinyon-juniper woodlands or in shadscale shrublands (Paige and Ritter 1999). At one time sage grouse were found in virtually all areas where sage brush (especially *Artemisia tridentata*) occurred in Western North America. It is hypothesized that the sage grouse breeding population circa 1800 was 1.1 million birds. Today, the estimated breeding population is 0.2 million (Parrish et al. 2002).

In Utah, sage grouse inhabit sagebrush habitat of the Colorado Plateau and Great Basin geographic regions from 6,000 to 9,000 ft. elevation. During spring, they use sagebrush habitats for breeding, feeding, roosting, nesting and rearing young (Connelly et al. 2000). Large, relatively continuous sagebrush stands, often exceeding 50 sq. mi., are needed to provide all habitat characteristics used by sage grouse; summer home ranges may be as small as 1 to 2.5 square miles, and annual home ranges may be as large as 577 square miles (Page and Ritter 1999).

Sage grouse males appear to form breeding leks opportunistically at sites within or adjacent to potential nesting habitat. Leks are typically established in openings within large sagebrush stands; openings include old lakebeds, low sagebrush flats, ridge tops, burn areas, and other open areas within sagebrush stands (Connelly et al. 2000). Most nests are placed under sagebrush in stands that provide higher than average canopies and lateral cover (Connelly et al. 2000). Nest sites also generally contain taller and denser grass cover than average. As sage brush habitats dry out during summer sage grouse use

a wider variety of habitats including meadow and riparian habitats. Hens with broods move to areas that support succulent vegetation including forbs (Parrish et al. 2002). Sites used by broods have been reported to have twice as much forb cover as independent sites (Connelly et al. 2000).

There suitable sage grouse habitat near the proposed project area.

Peregrine Falcon

The peregrine falcon is cosmopolitan, ranging from coast to coast in North America. Pesticide accumulation in the mid 1900s drove the peregrine to the verge of extinction, and by 1965 fewer than 20 pairs were known west of the Great Plains. In 1990 there were 326 known pairs in the southwest region (Rodriguez 2002). The peregrine falcon was federally listed as an endangered species in 1970, and again in 1984. With the help of reintroductions and pesticide controls (primarily banning DDT, which caused eggshell thinning and drastically low reproduction), the peregrine falcon population increased sufficiently to be de-listed in 2000.

Peregrine falcon preferred nesting habitat is on cliff faces with recesses or protected shelves, although reintroduced birds regularly nest on man-made structures such as towers and high-rise buildings. A wide variety of habitats are used for foraging, including riparian woodlands, open country near rivers and marshes, coniferous and deciduous forest edges, shrublands, and prairies. They prey on a wide variety of birds including pigeons, shorebirds, waterfowl, grouse and other small to mediums sized terrestrial birds. Peregrine falcons may travel up to 18 miles from their nest site to forage for food, however a 10 mile radius around the nest is an average hunting area, and 80% of foraging occurs within a mile of the nest (Spahr et al. 1991). The nearest known peregrine falcon eyrie is located approximately 3 ½ miles from the project area. Nesting peregrine falcons may forage in the vicinity of the proposed project.

C. MANAGEMENT INDICATOR SPECIES

Rocky Mountain Elk

Elk occurred within the mountainous regions of Utah historically. However, due to unlimited hunting, elk populations in the state diminished until 1898 when elk hunting was prohibited. Elk transplants were initiated in 1912 and continued until 1925. Today elk again occur within the mountainous regions of the state, and elk populations have increased dramatically over the last 20 years. They are once again considered a big game species in Utah.

Elk habitat includes semi-open forest and mountain meadows in the summer. They move to foothills, plains and valleys in winter. Rocky Mountain elk use uneven-aged, mature forest stands that include old growth characteristics, herbaceous openings, and water.

Dense brush understory is used for escape and thermal cover. They are herbivorous, and feed in riparian areas, meadows, and on herbaceous and brush stages of forest habitats. They graze and browse, eating grasses, forbs, tender twigs, and leaves of shrubs and trees, fungi, some mast, and aquatic vegetation.


A number of studies have shown that elk use has declined in areas adjacent to roads. The width of the area avoided has varied from 0.25 to 1.8 miles, depending on the amount and kind of traffic, quality of road, and density of cover adjacent to the road (Thomas and Toweill 1982). In general elk could be expected to move an average of approximately 0.5 miles from roadways that are being used.

The rut occurs from late August to November. Gestation period is about 255 days, and calving takes place during late spring and early summer in areas that provide dense cover with brushy vegetation near openings, available water, and seclusion from human impacts.

On the Wasatch Plateau, elk tend to occupy the higher elevation aspen and mixed conifer habitats from spring through early fall, and then move to lower elevation mixed shrub, pinyon/juniper, and sagebrush habitats for winter range. Elk generally occupy winter range from about the beginning of December through mid-April, but this varies depending on the severity of the winter. On the Plateau, parturition (calving) takes place roughly from the first part of May through early July, generally in aspen dominated habitat. Protection of winter range and calving habitat is considered a key factor in the maintenance of elk populations. It is important that higher nutritional demands during calving be met to improve the chances of calving success, cow recovery, and early calf growth. Therefore, available forage within calving habitat is especially important. Available forage within winter range is also important to increase chances of survival during this harsh season.

The elk population (composition and size) on the Manti-La Sal NF, for the most part, depends on the number and type of tags (Bull, Cow or Spike) issued by the Utah Department of Wildlife Resources (UDWR) each year, and on weather cycles and patterns. Graph 2 illustrates the results of UDWRs Manti Elk Census from 1992 through 2004.

Graph 2. The elk population (composition and size) from 1992 through 2004 within the Manti Elk Census unit (UDWR 2004).



The elk population for the Manti Elk Census in 2004 was slightly below the average population count for the 12 years of population information.

Mule Deer

Mule deer occur throughout the mountains and valleys of eastern Utah. Their populations throughout Utah have historically fluctuated, periodically affected by drought and severe winter weather. Populations in eastern Utah declined in the early to mid 1990s, but showed signs of recovery in the late 1990s. The decline was attributed to severe drought conditions from 1988 through 1992, which was followed by a severe winter in 1992-93. Other factors contributing to fluctuating mule deer populations include predators, habitat changes, and competition with elk.


Mule deer occupy several habitat types throughout the west including coniferous forests, desert shrubs, chaparral, and grassland with shrubs; they occur in early to intermediate successional stages of most forest, woodland, and brush habitats. Mule deer prefer a mosaic of various aged vegetation that provides woody cover, meadow and shrubby openings, and free water. Vegetation cover is critical for thermal regulation in winter and summer, and to provide escape cover. They browse and graze, and prefer tender new growth of various shrubs, many forbs, and a few grasses.

Human activity and traffic on roads are known to displace deer from the area of disturbance. The distance deer move away from disturbance areas depends on topographical features and the amount of vegetation cover in the area, but to average distance is approximately 660 feet from disturbance areas.

Rutting season occurs in late fall through early winter. Gestation is between 195 and 212 days, and fawns are born from early April to midsummer, with some geographic variation. Fawning peaks generally occur from late April through mid June. Fawning occurs in moderately dense shrublands and forests, dense herbaceous stands, and high elevation riparian and mountain shrub habitats that have available water and abundant forage.

The deer population on the Manti-La Sal NF, for the most part, depends on weather cycles and patterns. Graph 3 illustrates the results of UDWRs Manti deer population estimates from 1999 through 2003.

Graph 3. Population estimate of the deer population on the Manti Division of the Manti-La Sal National Forest from 1999 through 2003.



There is an upward trend in the deer population on the Manti over the 5 years of population information.


Golden Eagle

Golden eagles usually nest on cliffs overlooking large open expanses of grass-shrub or shrub steppe habitat, but tree nesting occurs in portions of their breeding range, including Utah. Nesting and brooding season generally extends from mid February to mid July. There is extensive cliff habitat along the eastern margin of the Wasatch Plateau and in canyons incising the Plateau. There are also extensive grassland and mountain brush habitats for foraging. Golden eagles primarily prey on small mammals including ground squirrels, prairie dogs, jack rabbits and cottontails.

Preferred golden eagle prey habitat includes edge along high mountain brush habitat, high/mid elevation perennial forb habitat, and high elevation perennial grassland habitat. Preferred golden eagle winter habitat includes large expanses of sagebrush.

There are a number of golden eagle nest sites located within the proposed project area; none of these nest sites were active in 2004. There are two golden eagle nest sites located less one mile from an area where project related activity could occur within the project area; neither of these nest sites have been active since surveys were began in 1998. The number of known golden eagle nests on the Forest has increased as new nests are found; therefore looking at the number of known active nests over the years would probably not give an accurate impression of the golden eagle population on the Forest since monitoring began in 1998. A better indication of how the golden eagle population is doing on the Forest would be the percent of monitored nest sites that were active each year, which is illustrated in Graph 5.

Graph 5. The percent of monitored golden eagle nest sites that were active on the Manti Division of the Manti-La Sal National Forest from 1998 through 2004.



The average percent of active nests over the 7 years of surveys is approximately 11.2%. Nesting activity was well below average in 2002, 2003 and 2004; there was somewhat of a rebound in 2004. There has not been a dramatic change in golden eagle nesting and foraging habitat attributed to management activity on the Forest over these 7 years of surveys. At least some of the change in golden eagle nesting activity during the last seven years is likely attributed to annual moisture.

D. PRIORITY MIGRATORY BIRD SPECIES

Virginia's Warbler

Virginia's warblers prefer scrub hillsides with a well developed herbaceous or woody understory. In Utah, preferred nesting habitat is lower elevation dense Gambel's oak stands. They are also known to nest in habitats with shrubby understories including: mountain mahogany, riparian areas, ponderosa pine, Douglas fir, aspen, and pinyon/juniper woodlands. Nests are typically embedded or covered with dead or decaying leaves and grasses in areas of dense brush. Virginia's warblers begin arriving in Utah in early May, and begin their breeding cycle from mid-May to early June (Parrish et al. 2002). They are a single brood nester. Pairs begin nesting by early June, and young fledge approximately 3 weeks later.

Their breeding range is almost exclusively in the southwestern United States. Historical nesting records for Utah include: Salt Lake County, Summit County, San Juan County, Utah County, Kane County, Garfield County, Daggett County, Beaver County, Weber County, and the Uinta Basin; in Utah, nesting elevation ranges from 4,000 to 10,000 ft. There has been no confirmed nesting in Emery County or on the Manti Division of the Manti-La Sal National Forest (Parrish et al. 2002).

Brewer's Sparrow

The subspecies of Brewer's sparrow that occurs in Utah is primarily a Great Basin species, but also occurs in shrubsteppe and high desert shrub (greasewood) habitats. They generally nest in habitats dominated by big sagebrush (*artemisa tridentata*), but occasionally use other shrubs. Nests are usually located in sagebrush patches that are taller and denser than surrounding habitat; with less herbaceous cover and more bare ground. They are primarily insectivorous during breeding season, consume mostly grass and weed seed in winter.

They generally arrive in Utah in mid April and depart in mid October (Parrish et al. 2002). Pair form shortly after arrival and nesting begins when weather permits. Hatching begins in late May and peaks in mid June (Parrish et al. 2002)

Brewer's sparrow populations are declining range wide, however in Utah their population appears to be stable and possibly increasing (Parrish et al. 2002).

Sage Sparrow

The sage sparrow is considered a shrubsteppe-obligate species, and are closely associated with big sagebrush (*A. tridentate*) throughout most of their distribution, but also uses bitterbrush, rabbitbrush, greasewood, tumbleweed, or bunch grasses. They nest primarily in shrubs, but nests have also been found in bunch grass and on the ground under shrubs. They prefer taller shrubs with larger canopies that provide cover. They are categorized as ground-foraging omnivores during nesting season and ground-gleaning granivores during

nonbreeding season (Parrish et al. 2002); nestlings are primarily fed spiders, butterflies, moths, true bugs and leafhoppers. They are known to occur up to 8,000 ft. in elevation.

III. AFFECTED ENVIRONMENT

The proposed project is located on a relatively high elevation plateau on the Castle Gate sandstone formation. There are a variety of habitats on this plateau including: pinyon/juniper, mixed conifer dominated by ponderosa pine, mixed conifer and aspen, mountain brush, sage brush and perennial grassland habitats. There are 6 drill sites in the proposed project plan: 5 drill holes are located in sagebrush dominated habitat and one drill hole is located in mountain brush habitat that includes sagebrush/rabbit brush, service berry and mahogany.

IV. ANALYSIS OF EFFECTS

This analysis of effects is based on the existing conditions within the project planning area. The analysis reviews the potential “direct and indirect effects” of the proposed SUFCO 2004 Helicopter-Assisted Coal Exploration Drilling Project on threatened, endangered and sensitive (TES) species, management indicator species (MIS), and priority migratory bird species. This report also states the expected “cumulative effects” that would potentially accrue to TES, MIS and priority migratory bird species if proposed project actions add cumulatively to other past, present or reasonably foreseeable future actions to impact the species of concern.

The past, present or reasonably foreseeable future actions that may add incrementally to impacts of the proposed SUFCO 2004 Helicopter-Assisted Coal Exploration Drilling Project include:

- Other exploration drilling activity
- Disbursed recreational activity
- Road construction and maintenance

A. Threatened and Endangered Species

Bald Eagle

Direct and Indirect Effects: There are no landscape characteristics in the vicinity of the proposed project that would attract bald eagles to the area; there are no water bodies that would provide suitable bald eagle forage habitat in or near the project area. The project area is not known or expected to be used by nesting, wintering or foraging bald eagles. However, bald eagles may occur incidentally while in transition during migration or dispersal during late fall or early winter months. These occurrences would only be

incidental and of short duration, and the proposed project would not alter bald eagle habitat. Therefore, the proposed project is not likely to directly or indirectly affect the bald eagle.

Cumulative Effects: Since the proposed project is not likely to exert direct or indirect affects on the bald eagle, no cumulative affects will accrue to this species because of the SUFCO 2004 Helicopter-Assisted Coal Exploration Drilling Project.

B. Sensitive Species

Spotted Bat

Direct and Indirect Effects: There are numerous cliff faces that could provide suitable spotted bat roost habitat within 2 miles of the proposed project area. The nearest suitable roost habitat is located approximately ½ mile from the nearest drill site. Activity during project implementation would not likely disturb roosting bats, and the project would not directly or indirectly impact spotted bat roost habitat.

The project would be implemented over a short period of time (7 plus days at each drill site) over small segments of the landscape that potentially provides suitable spotted bat forage habitat. However, since project activity would occur during daylight hours, it would not impact the nighttime foraging spotted bat. The proposed project would not appreciably directly or indirectly impact spotted bat foraging habitat.

Cumulative effects: Since the proposed project would not likely exert appreciable direct or indirect impacts on the spotted bat, no appreciable cumulative affects would accrue to this species because of the SUFCO 2004 Helicopter-Assisted Coal Exploration Drilling Project.

Townsend's big-eared Bat

Direct and Indirect Effects: There are a number of alcoves and cave like structures located within 2 miles of the proposed project area. Activity during project implementation would not likely disturb roosting bats; the project would not directly or indirectly impact Townsend's big-eared bat roost habitat.

The project will be implemented for a short period of time (7 plus days at each drill site) over small segments of the landscape that potentially provides suitable Townsend's big-eared bat forage habitat. However, since project activity would occur during daylight hours, it would not impact this nighttime foraging species. The proposed project would not appreciably directly or indirectly impact Townsend's big-eared bat foraging habitat.

Cumulative effects: Since the proposed project would not likely exert appreciable direct or indirect affects on the Townsend's big-eared bat, no appreciable cumulative affects

would accrue to this species because of the SUFCO 2004 Helicopter-Assisted Coal Exploration Drilling Project.

Greater Sage Grouse

Direct and Indirect Effects: The proposed project would occur outside the greater sage grouse lekking and breeding season, the project would not modify lekking or breeding habitat, and the project would not occur in brood rearing habitat. Therefore, the proposed project would not likely appreciably directly or indirectly impact the greater sage grouse.

Cumulative effects: Since the proposed project would not likely exert appreciable direct or indirect effects on the greater sage grouse, no appreciable cumulative effects would accrue to this species because of the SUFCO 2004 Helicopter-Assisted Coal Exploration Drilling Project.

Peregrine Falcon

Direct and Indirect Effects: The nearest known peregrine falcon eyrie is located approximately 3 ½ miles from the project area. Falcons may travel more than 18 miles from the nest site to hunt for food, however a 10 mile radius around the nest is an average hunting area, with 80% of foraging occurring within a mile of the nest. Nesting peregrine falcons may forage in the vicinity of the proposed project. Project implementation would not occur during the peregrine nesting period, and would only temporarily impact localized areas within potential forage habitat; therefore the proposed project would not likely appreciably directly or indirectly impact the peregrine falcon.

Cumulative effects: Since the proposed project would not likely exert appreciable direct or indirect effects on the peregrine falcon, no appreciable cumulative effects would accrue to this species because of the SUFCO 2004 Helicopter-Assisted Coal Exploration Drilling Project.

C. Management Indicator Species

Rocky Mountain Elk and Mule Deer

Direct Effects: Exploration holes would not be drilled simultaneously, but would be drilled consecutively one after the other. Each hole would take approximately 7 days to drill and cause relatively little habitat disturbance at each drill site. Potential direct impacts would occur over relatively small segments of the landscape for short periods of time. Drilling will occur during the time frame when deer and elk would be present on the plateau, but would occur outside the prime calving and fawning season for these species; therefore potential direct impacts to these species would be minor (would not impact the deer and elk populations in the area). Potential direct impacts could include

causing deer and elk to move away from areas where drilling is occurring; traffic along roadways associated with drilling activity may also cause disturbance.

Indirect Effects: Due to the short duration of the proposed project and the relatively small area of disturbance, there are not expected to be appreciable indirect impacts associated with the project.

Cumulative effects: Impacts on deer and elk from the proposed project may add cumulatively to impacts associated with disbursed recreational activity in the Pines and Big Ridge areas west of Emery, Utah. Potential impacts from disbursed recreational activity are variable; however the combined affects of these two activities is not expected to prevent deer or elk from using the general landscape of this area of the Forest.

Potential impacts from the proposed project are not expected to overlap temporally with other exploration drilling activities or road maintenance projects; therefore the proposed SUFCO 2004 Helicopter-Assisted Coal Exploration Drilling Project will not add cumulatively with those activities.

Golden Eagle

Direct and Indirect Effects: The nearest known golden eagle nest site is located approximately $\frac{3}{4}$ of a mile from one of the proposed drill sites, and there are a number of golden eagle nest sites located within 2 miles of the proposed project area; none of these nests were active in 2004. The proposed project will not directly affect these nest sites or any other golden eagle nest habitat. Golden eagles may forage in the vicinity of the proposed project; therefore the project could directly impact foraging eagles. These direct impacts may include diverting foraging eagle from the vicinity of project activity during drilling operations. The proposed project is not likely indirectly impact the golden eagle.

Cumulative Effects: The direct impacts from the proposed project may add cumulatively to impacts associated with disbursed recreational activity in the area. Potential impacts from disbursed recreational activity in the area are variable, however impacts are not expected to lead to mortality or reduced productivity. The cumulative affects of these activities is not expected to prevent golden eagles from using the general landscape of this area of the Forest.

D. PRIORITY MIGRATORY BIRD SPECIES

Virginia's Warbler

Direct and Indirect Effects: Virginia's warblers are not known to nest in San Pete or Sevier Counties, Utah. However, some of the pinyon/juniper/brush habitat near the proposed drill sites may provide characteristics of suitable nesting habitat. The proposed project would not remove suitable nesting habitat, and the project will not be

implemented during the nesting period for this species. Therefore there is not likely to be appreciable direct or indirect affects on this species.

Cumulative Effects: Since the proposed project is not likely to exert appreciable direct or indirect impacts on the Virginia's warbler, cumulative affects are not likely to accrue to this species as a result of the proposed SUFCO 2004 Helicopter-Assisted Coal Exploration Drilling Project.

Brewer's Sparrow

Direct and Indirect Effects: There is suitable Brewer's sparrow nesting habitat in the proposed project area. The proposed project is not expected to appreciably alter or remove suitable nesting habitat for this species. Project activity would not occur during this species breeding period; therefore there would be no impacts on nesting Brewer's sparrows. There would not be appreciable direct impacts to the Brewer's sparrow, and the project is not expected to cause any indirect impacts to this species.

Cumulative Effects: Since the proposed project is not likely to cause appreciable direct or indirect impacts on the Brewer's sparrow, no appreciable cumulative effects would accrue to this species as a result of the SUFCO 2004 Helicopter-Assisted Coal Exploration Drilling Project.

Sage Sparrow

Direct and Indirect Effects: There is suitable sage sparrow nesting habitat in the proposed project area; however the proposed project is not expected to appreciably alter or remove suitable nesting habitat for this species. Project activity would not occur during this species breeding period; therefore there would be no direct impacts on nesting sage sparrows. There would not be appreciable direct impacts to the sage sparrow, and the project is not expected to cause any indirect impacts to this species.

Cumulative Effects: Since the proposed project is not likely to cause appreciable direct or indirect impacts on the Brewer's sparrow, no appreciable cumulative effects would accrue to this species as a result of the SUFCO 2004 Helicopter-Assisted Coal Exploration Drilling Project.

V. DOCUMENTATION

Boschen, Nelson. 1997. Bald Eagles in Southeast Utah: 1997 Nesting Season. Moab Bureau of Land Management.

- Connelly, J.W., M.A. Schroeder, A.R. Sands, and C.E. Braun. 2000. Guidelines to manage sage grouse populations and their habitats. *Wildlife Society Bulletin*. 28(4): 967-985.
- Durrant, Stephen, D. 1952. *Mammals of Utah - Taxonomy and Distribution*. University of Kansas Publications, Museum of Natural History, Volume 6. pp. 549.
- Hasenyager, R. N. 1980. *Bats of Utah*. Utah Division of Wildlife Resources. Publication Number 80-15.
- Hynes, H.B.N. 1972. *The Ecology of Running Waters*. University of Toronto Press. Toronto, Canada.
- Oliver, G.V. 2000. *The Bats of Utah: A Literature Review*. Utah Division of Wildlife Resources, Salt Lake City, Utah.
- Paige, C., and S.A. Ritter. 1999. *Birds in a Sagebrush Sea: Managing sagebrush habitats for Bird Communities*. Partners in Flight Western Working Group, Boise, Idaho.
- Parrish, J.R., F.P. Howe, R.E. Norvell. 2002. *Utah Partners in Flight Avian Conservation Strategy Version 2.0*. Utah Partners in Flight Program, Utah Division of Wildlife Resources, 1594 West North Temple, Salt Lake City, Utah 84116, UDWR Publication Number 02-27.
- Pierson, E.D., M.C. Wackenhut, J.S. Altenbach, P. Bradley, P. Call, D.L. Genter, C.E.Harris, B. L. Keller, B. Lengus, L. Lewis, B. Luce, K.W. Navo, J.M. Perkins, S. Smith, and L. Welch. 1999. *Species Conservation Assessment and Strategy for Townsend's Big-eared Bat (*Corynorhinus townsendii townsendii* and *Corynorhinus townsendii pallescens*)*. Idaho Conservation Effort, Idaho Department of Fish and Game, Boise, Idaho.
- Perkins, J.M. and J.R. Peterson. 1997. *Bat survey for the SUFCO Mine, Emery County, Utah*. 8pp.
- Rodriguez, R.L. 2002. *Life History and Analysis of Endangered, Threatened, Candidate, Sensitive, and Management Indicator Species of the Fishlake National forest*. Version 2.0.
- Sherwin, R.E., D.S. Rogers, and C.A. Johansson. 1997. *Assessment of spotted bat (*Euderma maculatum*) and Townsend's big-eared bat (*Corynorhinus townsendii*) in the proposed Cottonwood Canyon lease area*. Manti La Sal National Forest, Emery County, Utah. Conducted for Energy West Mining Co. 18pp + append.
- Thomas, J.W., and D.E. Toweill. 1982. *Elk of North America: Ecology and Management*. Stackpole Books, Harrisburg, Pa.

- Toone, R.A. 1994. General Inventory for Bats in the Abajo and La Sal Mountains, Manti-La Sal National Forest, with Emphasis on the Spotted Bat (*Euderma maculatum*) and the Townsend's Big-eared Bat (*Plecotus townsendii*). Heritage Program Utah Department of Natural Resources, Salt Lake City, Utah.
- UDWR. 2004. Results of the Utah Department of Wildlife Resource – Manti Elk Census 92-04. Field Office, Price, Utah.
- USDA, U.S. Forest Service. 1986. Manti-La Sal National Forest Land and Resource Management Plan: Forest Plan Amendment, Appendix A. MLSNF, Price, Utah.
- USDI, U.S. Fish and Wildlife Service. 2002a. Bonytail (*Gila elegans*) Recovery Goals: Amendment and Supplement to the Bonytail Chub Recovery Plan. U.S. Fish and Wildlife Service, Mountain-Prairie Region (6), Denver, Colorado.
- USDI, U.S. Fish and Wildlife Service. 2002b. Humback Chub (*Gila cypha*) Recovery Goals: Amendment and Supplement to the Humback Chub Recovery Plan. U.S. Fish and Wildlife Service, Mountain-Prairie Region (6), Denver, Colorado.
- USDI, U.S. Fish and Wildlife Service. 2002c. Colorado Pikeminnow (*Ptychocheilus lucius*) Recovery Goals: Amendment and Supplement to the Colorado Squawfish Recovery Plan. U.S. Fish and Wildlife Service, Mountain-Prairie Region (6), Denver, Colorado.
- USDI, U.S. Fish and Wildlife Service. 2002d. Razorback sucker (*Xyrauchen texanus*) Recovery Goals: Amendment and Supplement to the Razorback Sucker Recovery Plan. U.S. Fish and Wildlife Service, Mountain-Prairie Region (6), Denver, Colorado.
- USDI, U.S. Fish and Wildlife Service. 2002e. Utah Field Office Guidelines for Raptor Protection from Human and Land Use Disturbances. Utah Field Office, Salt Lake City.
- USDI, Fish and Wildlife Service. 2001a. Final designation of critical habitat for the Mexican spotted owl. Federal Register 66(22): 8530-8553.
- USDI, U.S. Fish and Wildlife Service. 2001b. Status Review for Bonneville Cutthroat Trout (*Oncorhynchus clarki utah*). Regions 1 and 6 Portland, Oregon and Denver, Colorado.
- Watkins, L. C. 1977. Mammalian Species: *Euderma maculatum*. The American Society of Mammalogists. 77:1-4.

**Muddy Creek Technical Report
Wildlife**

Table of Contents

Table of Contents	2
List of Tables.....	5
List of Figures	5
List of Appendices	5
1.0 Introduction	1
1.1 Statement of Project Objectives	1
1.2 Statement of the Issues with Evaluation Criteria.....	1
1.3 Description of the Alternatives Evaluated.....	2
1.3.1 Alternative 1 - No Action Alternative	2
1.3.2 Alternative 2 - Standard Lease Terms and Conditions.....	2
1.3.3 Alternative 3 - Standard Lease Terms and Conditions and Special Stipulations	2
1.3.4 Alternative 4 – Standard Lease Terms and Conditions and Special Stipulations That Address Other Significant Issues	2
2.0 Methods.....	2
2.1 Contacts Made.....	2
2.2 Sources and Descriptions of Existing Information.....	3
2.3 Data Collection and Analysis Methodology.....	4
2.4 Description of Inventories and Data Collected by the Consultant	6
2.4.1 TEPS Wildlife Surveys.....	6
2.4.1.1 Bald eagle	6
2.4.1.2 Colorado River cutthroat trout.....	6
2.4.1.3 Northern goshawk	8
2.4.1.3.1 Survey Methods.....	8
2.4.1.3.2 Survey Results.....	8
2.4.1.4 Flammulated owl	9
2.4.1.4.1 Survey Methods.....	9
2.4.1.4.2 Survey Results	9
2.4.1.5 Three-toed woodpecker	10
2.4.1.5.1 Survey Methods.....	10
2.4.1.5.2 Survey Results	10
2.4.1.6 Spotted bats	11
2.4.1.6.1 Survey Methods.....	11
2.4.1.6.2 Survey Results	11
2.4.1.7 Western big-eared bats	11
2.4.1.7.1 Survey Methods.....	11
2.4.1.7.2 Survey Results	12
2.4.1.8 Additional Bat Surveys.....	12
2.4.1.8.1 Cooperative Challenge Cost Share Project.....	12
2.4.1.8.2 SUFCO Mine Bat Survey	12
2.4.2 Management Indicator Species.....	12
2.4.2.1 Golden eagle (UPDATE)	12
2.4.2.1.1 Survey Methods.....	13
2.4.2.1.2 Survey Results	13
2.4.2.2 Mule Deer.....	13
2.4.2.3 Elk	14
2.4.2.4 Blue Grouse	14
2.4.2.5 Macroinvertebrates	14
2.4.2.5.1 Year 2001 Surveys	15

2.4.2.5.2 Year 2002 Surveys	16
2.4.2.5.3 Year 2003 Surveys	18
2.4.2.5.4 Survey Summary: 2001- 2003	19
2.4.3 Species of High Federal Interest.....	22
2.4.3.1 Migratory Birds	22
2.4.3.1.1 Survey Methods.....	22
2.4.3.1.2 Survey Results	23
2.4.4 Other Wildlife Species	24
2.4.4.1 Sage-grouse	24
2.4.4.1.1 Strutting Ground Surveys	24
2.4.4.1.2 Grouse Sign Surveys	24
2.4.4.1.3 Survey Results	24
2.4.4.2 Amphibians.....	25
2.4.4.2.1 Survey Methods.....	25
2.4.4.2.2 Survey Results	25
2.4.4.3 Reptiles	26
2.4.4.4 Small mammals	26
2.4.4.5 Non-game birds	28
3.0 Results and Discussion	29
3.1 Description of the Affected Environment	29
3.1.1 Aquatic and Terrestrial Habitat	29
3.1.1.1 Aquatic Habitat.....	30
3.1.1.2 Terrestrial Habitat.....	30
3.1.2 TEPS.....	30
TEPS Fish.....	32
3.1.2.1 Bonytail	32
3.1.2.2 Colorado Pikeminnow	32
3.1.2.3 Humpback Chub.....	32
3.1.2.4 Razorback Sucker.....	32
3.1.2.5 Colorado River Cutthroat Trout	33
TEPS Birds.....	33
3.1.2.6 Bald Eagle	33
3.1.2.7 Mexican Spotted Owl.....	33
3.1.2.8 Northern Goshawk.....	34
3.1.2.9 Flammulated Owl	34
3.1.2.10 Peregrine Falcon.....	35
3.1.2.11 Three-Toed Woodpecker.....	35
3.1.2.12 Greater Sage-Grouse	35
TEPS Mammals.....	36
3.1.2.13 Canada Lynx.....	36
3.1.2.14 Spotted Bat	36
3.1.2.15 Western Big-Eared Bat.....	36
TEPS Amphibians	37
3.1.2.16 Spotted Frog	37
3.1.3 Management Indicator Species.....	37
3.1.3.1 Golden Eagle	37
3.1.3.2 Mule Deer.....	37
3.1.3.3 Elk	38
3.1.3.4 Aquatic Macroinvertebrates	38
3.1.4 Species of High Federal Interest.....	39
3.1.4.1 Migratory Birds	39

- 3.1.5 Other Species.....40
 - 3.1.5.1 Fishes.....40
 - 3.1.5.1.1 Brook trout.....40
 - 3.1.5.1.2 Rainbow trout.....40
 - 3.1.5.2 Blue Grouse.....40
 - 3.1.5.3 Amphibians.....40
 - 3.1.5.4 Reptiles.....41
 - 3.1.5.5 Small Mammals.....41
 - 3.1.5.6 Non-Game Birds.....41
- 3.2 Detailed Technical Assessment/Description of the Potential Effects.....41
 - 3.2.1 Wildlife Issue 1.....41
 - 3.2.1.1 Alternative 1: No-Action.....41
 - 3.2.1.2 Alternative 2: Standard Lease Terms and Conditions Only.....41
 - 3.2.1.2.1 Fisheries.....42
 - 3.2.1.2.2 Macroinvertebrates.....43
 - 3.2.1.2.3 Birds.....43
 - 3.2.1.2.4 Mammals.....44
 - 3.2.1.2.5 Amphibians.....44
 - 3.2.1.2.6 Reptiles.....44
 - 3.2.1.3 Alternative 3: Standard Lease Terms and Conditions and Special Stipulations.....44
 - 3.2.1.3.1 Aquatic and Terrestrial Wildlife.....44
 - 3.2.2 Wildlife Issue 2.....45
 - 3.2.2.1 Alternative 1: No-Action.....45
 - 3.2.2.2 Alternative 2: Standard Lease Terms and Conditions Only.....45
 - 3.2.2.2.1 Fisheries.....45
 - 3.2.2.2.2 Macroinvertebrates.....46
 - 3.2.2.3 Alternative 3: Standard Lease Terms and Conditions and Special Stipulations.....46
 - 3.2.2.3.1 Fish and Macroinvertebrates.....46
 - 3.2.3 Wildlife Issue 3.....47
 - 3.2.3.1 Alternative 1: No-Action.....47
 - 3.2.3.2 Alternative 2: Standard Lease Terms and Conditions Only.....47
 - 3.2.3.2.1 TEPS Fishes.....47
 - 3.2.3.2.2 TEPS Birds.....48
 - 3.2.3.2.3 TEPS Mammals.....48
 - 3.2.3.2.4 MIS.....48
 - 3.2.3.2.5 Species of High Federal Interest.....48
 - 3.2.3.2.6 Other Species.....49
 - 3.2.3.3 Alternative 3: Standard Lease Terms and Conditions and Special Stipulations.....49
 - 3.2.3.3.1 Aquatic and Terrestrial Wildlife.....49
 - 3.2.4 Wildlife Issue 4.....49
 - 3.2.4.1 Alternative 1: No-Action.....49
 - 3.2.4.2 Alternative 2: Standard Lease Terms and Conditions Only.....49
 - 3.2.4.2.1 TEPS Fishes.....50
 - 3.2.4.2.2 TEPS Birds.....50
 - 3.2.4.2.3 TEPS Mammals.....50
 - 3.2.4.2.4 MIS.....50
 - 3.2.4.2.5 Species of High Federal Interest.....50
 - 3.2.4.2.6 Other Species.....50
 - 3.2.4.3 Alternative 3: Standard Lease Terms and Conditions and Special Stipulations.....50
 - 3.2.4.3.1 Aquatic and Terrestrial Wildlife.....50
- 3.3 Mitigation and Monitoring Recommendations.....51

3.4 Cumulative Effects	51
4.0 Literature Cited and Contacts	51
5.0 List of Preparers with Qualifications of Preparers	56

List of Tables

Table 1. Wildlife survey methodology for the Muddy analysis area, Manti-La Sal National Forest.	4
Table 2. Fisheries surveys within the Muddy analysis area in the Manti-La Sal National Forest. .	7
Table 3. Stream habitat data for fisheries surveys conducted within the Muddy analysis area, Manti-La Sal National Forest, 2001 – 2002.	7
Table 4. Golden Eagle Surveys Conducted in the Muddy analysis area by UDWR, 1998-2003..	13
Table 5. Macroinvertebrate data from the Muddy Analysis Area. June and August 2001.	15
Table 6. Macroinvertebrate data from the Muddy Analysis Area. May and September 2002.....	17
Table 7. Macroinvertebrate data from the Muddy Analysis Area. May and September 2003.....	18
Table 8. Macroinvertebrate data from the Muddy Analysis Area. Three-year averages by season (2001-2003).	21
Table 9. Migratory birds of high federal interest.....	22
Table 10. Federally listed threatened and endangered and candidate species and Forest Service sensitive species potentially occurring on the MLNF in Emery, Sanpete, and/or Sevier counties.....	31
Table 11. Habitat disturbance associated with exploratory drilling in the Muddy analysis area. .	47

List of Figures

Figures 1-4. Summary of quantitative macroinvertebrate surveys conducted in the Muddy analysis area.....	39
--	----

LIST OF APPENDICES

Appendix A. Wildlife Survey Figures.....	58
Appendix B. Macro-invertebrate taxa collected in the Muddy analysis area, 2001-2003.....	59
Appendix C. Wildlife species observed in the Muddy analysis area, 2001 - 2003.	62
Appendix D. Small mammals.....	64
Appendix E. Non-game birds.....	80

Muddy Creek Technical Report

Wildlife

1.0 Introduction

1.1 Statement of Project Objectives

Passage of the Utah Schools and Lands Exchange Act of 1998 included the exchange of lands to resolve issues associated with creation of the Escalante-Grand Staircase National Monument. To balance land values exchanged under that act, the coal estates on several tracts of federal coal underlying the Manti-LaSal National Forest (MLNF) were conveyed to the State of Utah School and Institutional Trust Lands Administration (SITLA). This conveyance is temporal and the ownership of the coal will revert back to the federal government once a specific tonnage is produced or a specified royalty value is collected.

On the conveyed coal estates, SITLA has sole authority to lease the coal. Under the Surface Mine Control and Reclamation Act of 1977 and Utah Coal Rules, Forest Service must consent to the mine plan prior to mine development and can impose requirements for the protection of non-coal resources. The Forest Service decisions, as federal actions, are subject to the requirements of the National Environmental Policy Act of 1969 (NEPA), requiring environmental analysis and appropriate NEPA documents.

On the remaining federal coal estates within the Muddy Creek tract on National Forest System land, the U.S. Department of Interior, Bureau of Land Management (BLM) is the leasing authority. Under the Mineral Leasing Act of 1920, as amended by the Federal Coal Leasing Amendments Act of 1975, leases can only be issued by the BLM with consent from the Forest Service with conditions for protection of non-mineral resources. As federal actions subject to NEPA, both the BLM leasing decisions and the Forest Service consent decisions must be based on an environmental analysis and appropriate NEPA document.

This wildlife technical report is the result of three years of study of the Muddy Creek tract by Cirrus Ecological Solutions, LC (Cirrus), which included field studies, data acquisition, and data analyses and summaries. This technical report will form the basis for an analysis of impacts to wildlife in the project area in the subsequent Environmental Impact Statement (EIS) planned for the Muddy Creek tract on MLNF.

1.2 Statement of the Issues with Evaluation Criteria

The following wildlife issues and evaluation criteria were provided by the Forest Service in the scope of work for the Manti-LaSal Coal Tract Evaluations:

Wildlife Issue 1: Any changes in water flow and quality in perennial drainages and reservoirs or to riparian vegetation/wetlands could affect habitat for terrestrial and aquatic species.

Evaluation Criteria: Description of Potential Effect to Affected Habitat Amount and Quality.

Wildlife Issue 2: Subsidence of perennial streams could cause changes in stream morphology and aquatic habitat.

Evaluation Criteria: Description of changes to ratio of habitat types (pools, riffles, runs, glides, and cascades); changes in streambed sediments (spawning habitat); changes in bank stability.

Wildlife Issue 3: Exploration drilling and construction of mine vent holes could temporarily disrupt use of summer habitat by terrestrial species.

Evaluation Criteria: Area and Duration of Avoidance by Affected Species.

Wildlife Issue 4: Construction and operation of mine facilities and haul roads and coal traffic could remove habitat and associated noise/activity could displace dispersed wildlife (avoidance) including threatened, endangered, proposed and sensitive species.

Evaluation Criteria: Area of habitat removed or changed, type of habitat lost, duration of loss, area avoided, percent of available habitat effective habitat remaining, adequacy of remaining habitat to support wildlife populations.

1.3 Description of the Alternatives Evaluated

1.3.1 Alternative 1 - No Action Alternative

Under the No Action Alternative, no mining would take place on the Muddy Creek tract. For this technical report, the No Action Alternative represents the baseline for estimating the effects of the action alternatives on wildlife in the project area. Further analysis of the No Action Alternative has been deferred until the EIS for this project is initiated.

1.3.2 Alternative 2 - Standard Lease Terms and Conditions

Under this alternative, the Muddy Creek tract would be leased and mined with BLM standard lease terms and conditions (USDI-BLM undated). No special coal lease stipulations would be included in the lease, and longwall mining would be allowed throughout the tract which could result in subsidence of perennial drainages, escarpments, and surface facilities. This alternative emphasizes maximum coal production assuming maximum economic production with no specific restrictions for protection of surface resources from the effects of subsidence and is expected to result in the greatest amount of environmental impact. A more complete description of Alternative 2 can be found in the Conceptual Mine Plan for the Muddy Creek Tract located in the Detailed Description of Alternatives.

1.3.3 Alternative 3 - Standard Lease Terms and Conditions and Special Stipulations

Under Alternative 3, the Muddy Creek tract would be leased and mined with BLM standard lease terms and conditions (USDI-BLM undated) and Manti-LaSal National Forest's special stipulations (Forest Service 2003a). This alternative emphasizes protection of surface resources. Subsidence of perennial streams, escarpments, and surface facilities would not be allowed. There would, however, be no specific prohibition on subsidence of roads, trails, or range improvements. This is the most restrictive action alternative and would likely result in the least environmental damage. A more complete description of Alternative 3 can be found in the Detailed Description of Alternatives.

1.3.4 Alternative 4 – Standard Lease Terms and Conditions and Special Stipulations That Address Other Significant Issues

Under this alternative, the Muddy Creek tract would be leased and mined with BLM standard lease terms and conditions, as well as special stipulations to balance and address significant social, economic, or environmental issues or opportunities identified during analysis of Alternatives 1-3. No major potential impacts were identified for Alternative 3; therefore, Alternative 4 is not analyzed in this technical report.

2.0 Methods

2.1 Contacts Made

The following resource specialists were contacted over the contract period to obtain data, species lists, and/or discuss survey methods and results:

Manti La-Sal National Forest, USDA Forest Service

- Rod Player, Wildlife Biologist, Price Ranger District, Price, UT
- Kara Staab, Former Wildlife Biologist, Ferron Ranger District, UT
- Jeff Jewkes, Wildlife Biologist, Ferron Ranger District, MLNF
- Rob Davies, Former Fisheries Biologist, Price Ranger District, Price, UT
- Pamela Jewkes, Fisheries Biologist, Ferron Ranger District, MLNF
- Dale Harber, Contracting Officer Representative, MLNF, Price, UT

Utah Division of Wildlife Resources (UDWR)

- Ron Hodson, Former Wildlife Biologist, Southeastern Region, Price, UT
Current Wildlife Manager, Northern Region, Ogden, UT

- Chris Colt, Wildlife Biologist, Habitat Program Manager, Price, UT
- Craig Walker, Aquatic Biologist, Southeastern Region, Price, UT
- Louis Berg, Former Regional Aquatic Program Manager, Southeastern Region, Price, UT
- Amy Seglund, Sensitive Species Biologist, Southeastern Region, Price, UT

USDI Fish and Wildlife Service

- Laura Romin, T&E Species Biologist, Salt Lake Field Office, Salt Lake City, UT

Division of Oil, Gas, Mining

- Mark Mesch, Department of Utah Abandoned Mine Reclamation, Salt Lake City, UT

Utah State University (USU)

- Mark Vinson, Director, BLM BugLab & Research Assistant Professor, Department of Aquatic, Watershed, and Earth Resources, Logan, UT
- Jeff Ostermiller, Graduate Research Assistant, Aquatic Ecology Lab, Logan, UT

2.2 Sources and Descriptions of Existing Information

- UTM coordinates for bald eagle nest near Castledale. Received from the UDWR, Southeastern District.
- Fisheries survey data and sample locations in the analysis area. Received from the UDWR, Southeastern District.
- Report for UDWR Project Number F-44-R containing data on fisheries surveys in the Muddy Drainage. (Hart and Berg 2003).
- Location of goshawk nesting territories in the vicinity of the analysis area. Received from the Forest Service, Ferron Ranger District.
- Bat survey report for the SUFCO Mine, Emery County, Utah. (Perkins and Peterson, 1997).
- General inventory report for spotted bats on the Wasatch Plateau, MLNF. (Toone 1993).
- Raptor survey data conducted by UDWR over the Pines and Muddy coal tracts. Digital coverage data clipped to the analysis area received from the UDWR Southeastern District.
- Digital coverage data for mule deer winter and summer range was acquired from the UDWR GIS Data website (<http://dwrcdc.nr.utah.gov/ucdc/DownloadGIS/>) and received clipped to the analysis area from the Southeastern District.
- Utah big game annual report - 2001. Publication Number 01-30. UDWR.
- Digital coverage data for elk winter and summer range was acquired from the UDWR GIS Data website (<http://dwrcdc.nr.utah.gov/ucdc/DownloadGIS/>) and received clipped to the analysis area from the Southeastern District.
- Elk population data received from the UDWR Northern Region.
- Predicted elk calving data model received from the Forest Service, Ferron Ranger District.
- Digital coverage data for blue grouse potential habitat was acquired from the UDWR GIS Data website (<http://dwrcdc.nr.utah.gov/ucdc/DownloadGIS/>).
- List of species of high federal interest received from the FWS, Salt Lake Field Office.

- Sage-grouse data collected by UDWR was acquired from the UDWR Southwestern Region. Digital coverage data of sage-grouse habitat was acquired from the UDWR GIS Data website.
- Potential presence of species of high federal interest, small-mammals, and non-game birds was predicted by consulting the following resources, in addition to the UDWR raptor data listed above:
 - Fauna of Southeastern Utah and life requisites regarding their ecosystems (Dalton et al. 1990). Publication No. 90-11.
 - The Birder's Handbook (Erlich et al. 1988).
 - Inventory of Sensitive Species and Ecosystems in Utah. Inventory of Sensitive Vertebrate and Invertebrate Species: A Progress Report (UDWR 1997).
 - Colorado GAP Analysis website (<http://ndis.nrel.colostate.edu/cogap/>). (CDOW 2001).
 - Utah Conservation Data Center. Utah Division of Wildlife Resources, UCDC website, species information and Utah distribution maps (<http://www.utahcdc.usu.edu/ucdc>).
- Land and Resource Management Plan for the MLNF. 1986 and 2003 amendments.
- Vegetative coverage for the Manti-La Sal National Forest: Division and Sanpitch Divisions downloaded from the Forest Geographic Data website (<http://www.fs.fed.us/r4/mantilasal/downloads/>).
- Final Environmental Impact Statement for the Pines Tract Project (Forest Service 1999).

2.3 Data Collection and Analysis Methodology

Wildlife surveys and/or habitat assessments were conducted as part of the contract stipulations for the coal tract evaluation project. Data was collected for the following categories of terrestrial and aquatic wildlife: federally listed threatened and endangered species, and Forest Service sensitive species (TEPS), management indicator species (MIS), species of high federal interest, sage-grouse, amphibians and reptiles, small mammals, and non-game birds. Surveys were conducted between 2001 and 2003 in the Muddy Creek coal tract and within a 2-mile buffer surrounding the tract. This entire area is referred to as the analysis area throughout this document. Table 1 summarizes the methods associated with data collection and analysis by species. More detailed discussion of wildlife inventory methods and results is included in section 2.4 below.

Table 1. Wildlife survey methodology for the Muddy analysis area, Manti-La Sal National Forest.		
Species	Data Collection	Data Analysis
TEPS		
Bald eagle (<i>Haliaeetus leucocephalus</i>) (Threatened)	Existing data acquired from UDWR. Incidental observations recorded by Cirrus. No formal survey was required.	Species presence or absence in the analysis area determined with the use of GIS. No digital coverage was created because no nests were found.
Cutthroat trout (<i>Oncorhynchus clarki</i>) (FS Sensitive)	Existing survey data acquired from the UDWR. No formal survey by Cirrus was required.	Species presence or absence in the analysis area determined with the use of GIS. Digital coverage of surveyed reaches created from written descriptions and/or UTMs with ESRI ArcView software.
Northern goshawk (<i>Accipiter gentilis</i>) (FS Sensitive)	Presence/absence surveys conducted by Cirrus over two field seasons. Region 4 survey protocol for northern goshawks used (USDA-FS 1993).	Digital coverage of species presence (based on vocal responses and goshawk observations) and survey points created with Microsoft Excel and ESRI ArcView software.
Flammulated owl (<i>Otus flammeolus</i>) (FS Sensitive)	Presence/absence surveys conducted by Cirrus over two field seasons. UDWR Northern Region forest owl inventory protocol used (UDWR 1992).	Digital coverage of species presence (based on vocal responses) and survey points created with Microsoft Excel and ESRI ArcView software.

Table 1. Wildlife survey methodology for the Muddy analysis area, Manti-La Sal National Forest.

Species	Data Collection	Data Analysis
Three-toed woodpecker (<i>Picoides ridactytus</i>) (FS Sensitive)	Presence/absence surveys conducted by Cirrus over one field season. UDWR and UNHP (1992) woodpecker survey protocol used.	Digital coverage of species presence (based on vocal responses and woodpecker observations) and survey points created with Microsoft Excel and ESRI ArcView software.
Spotted bat (<i>Euderma maculatum</i>) (FS Sensitive)	Structural habitat searches were conducted and incidental sightings were recorded by Cirrus over two field seasons. Survey data from the Utah Natural Heritage Program and the SUFCO mine was acquired.	Digital coverage of species presence (based on audible vocalizations) created with Microsoft Excel and ESRI ArcView software.
Western big-eared bat (<i>Corynorhinus townsendii pallescens</i>) (FS Sensitive)	Conducted habitat assessment surveys using information on mine status. Structural habitat searches also conducted. Survey data from the Utah Natural Heritage Program and the SUFCO mine was acquired.	Survey results discussed in a narrative.
MIS		
Golden eagle (<i>Aquila chrysaetos</i>)	Aerial survey for golden eagle nest sites conducted by Cirrus and UDWR in 2002. Additional survey data for the analysis area acquired from UDWR.	Digital coverage of nest locations created by UDWR and clipped to the project area by Cirrus with Microsoft Access, and ESRI ArcView software.
Mule deer (<i>Odocoileus hemionus</i>) and elk (<i>Cervus elaphus</i>)	Existing survey and habitat modeling data acquired from UDWR. No formal survey by Cirrus was required.	Digital coverage of winter and summer range created by UDWR and clipped to the project area by Cirrus with Microsoft Access and ESRI ArcView software.
Blue grouse ¹ (<i>Dendragapus obscurus</i>)	Incidental sightings recorded by Cirrus. No formal survey was required. Habitat modeling data acquired from UDWR.	Digital coverage of incidental sightings created by Cirrus with Microsoft Excel and ESRI ArcView software.
Aquatic macro-invertebrates	Baseline data surveys were conducted for three years during the spring and fall by Cirrus. Macroinvertebrate samples and stream habitat data were collected. The USU/BLM National Aquatic Monitoring Center stream invertebrate sampling protocol was used (Hawkins et. al 1998).	Macroinvertebrate samples were analyzed by the National Aquatic Monitoring Center, Department of Aquatic, Watershed, and Earth Resources, Utah State University. Species abundance, diversity, and biotic health indices were generated. Digital coverage of survey areas created with Microsoft Excel and ESRI ArcView.
Species of High Federal Interest		
Migratory birds ²	Presence/absence surveys were conducted in suitable habitat by Cirrus over one field season. Incidental observations also made. Raptor nest data acquired from UDWR.	Digital coverage of observed territories created with Microsoft Excel and ESRI ArcView software.
Other Wildlife Species		
Sage-grouse (<i>Centrocercus urophasianus</i>)	Presence/absence surveys were conducted in suitable habitat by Cirrus over one field season. Incidental observations also made. Lek counts were made by Cirrus and UDWR.	Digital coverage of species presence and sign, survey areas, and lek sites created with Microsoft Excel and ESRI ArcView.
Amphibians	Presence/absence surveys were conducted in suitable habitat by Cirrus over two field seasons.	Digital coverage of suitable habitat and species presence created with Microsoft Excel and ESRI ArcView.
Reptiles	Incidental sightings recorded by Cirrus. No formal surveys required since construction	Species observations summarized in text.

Species	Data Collection	Data Analysis
	of mining facilities and roads was not proposed.	
Small mammals	Incidental sightings recorded by Cirrus. No formal surveys required. UDWR reports acquired.	Probability of occurrence analysis conducted using existing literature and other resources. Results summarized in tabular format.
Non-game birds	Incidental sightings recorded by Cirrus. No formal surveys required. UDWR reports acquired.	Probability of occurrence analysis conducted using existing literature and other resources. Results summarized in tabular format.

¹Note that the blue grouse is no longer a MIS. It was replaced in June 2003 by the Northern goshawk in an amendment to the MLNF Forest Plan.
²Migratory bird species of High Federal Interest are shown in Table 4.
³Note that the greater sage-grouse was added to the Region 4 sensitive species list in December 2003.

2.4 Description of Inventories and Data Collected by the Consultant

A description of field surveys and other forms of data acquisition, including survey methods and results, is discussed below in sections 2.4.1 - 2.4.4. Order of species described follows that outlined above in Table 1. Summary figures and tables, where applicable, are included in Appendices A, B, D, and E. In addition to the required survey data, general species lists were generated to document incidental wildlife sightings in the analysis area and are included in Appendix C.

2.4.1 TEPS Wildlife Surveys

2.4.1.1 Bald eagle

Data on bald eagle nest sites was acquired from the UDWR and is reported below.

There are no known bald eagle nests present on the Muddy tract or elsewhere on the Manti-La Sal National Forest. The closest nest is on private land about 18 miles east of the northeastern boundary of the analysis area, near the town of Castledale. It is unlikely that individuals from this eagle pair would utilize portions of the analysis area for foraging, since suitable habitat is available closer to the nest site. Five bald eagle individuals (3 adults and 2 juveniles) were observed in November 2003 along Cowboy Creek, presumably during fall migration. No other observations of this species were made during field visits between March and November, 2001-2003.

2.4.1.2 Colorado River cutthroat trout

Fish survey data was requested from the UDWR for perennial streams located within the Muddy analysis area. Surveys were conducted by UDWR personnel using standard electrofishing procedures. Streams surveyed included Muddy Creek, South and North Forks of Muddy Creek, and the North Fork of Quitchupah Creek. Fish surveys in other streams within the analysis area were not conducted. Cutthroat trout, believed to be of the Colorado River subspecies, were recorded during the most recent survey efforts in Muddy Creek and South Fork of Muddy Creek. Cutthroat trout were also observed incidentally in the North Fork of Muddy Creek, but electrofishing surveys have not been conducted there to date. Cutthroat trout were not observed within the North Fork of Quitchupah Creek. Results of the fisheries surveys are recorded by stream reach in Table 2. Cutthroat trout collected were assumed native. Rainbow trout and brook trout are introduced (non-native). Locations of sampled stream reaches were mapped and are depicted in Figure A-1, Appendix A. A digital coverage of the surveyed reaches in Figure A-1 is provided in conjunction with this report.

Region 1/4 aquatic habitat inventory data was not collected for fish-bearing streams by UDWR and was not part of the Cirrus proposed work plan for RFP 10-00-064. However, baseline habitat data was collected by UDWR at the sampled reaches and is reported below in Table 3. Habitat quality is summarized in Table 2.

Anthropogenic activities have led to the deterioration of riparian habitats and streams. Landslides, bank erosion, and sedimentation continue to impact streams and consequently fish habitat. The deterioration of riparian areas has reduced their capacity to provide shade to streams and to trap and retain sediments and pollutants from upslope areas. All these factors are contributing to the deterioration of fisheries resources.

Fish species not observed during surveys but predicted to occur or formerly observed in Muddy Creek include flannelmouth sucker, bluehead sucker and speckled dace. Fish species potentially present in Quitchupah Creek include flannelmouth sucker, leatherside chub, mountain sucker, and speckled dace.

Waterbody	Survey Date	Species Observed	Average Fish Size (Range)	Habitat Quality
Muddy Creek	July 2002	Cutthroat trout (most abundant) Rainbow trout Brook trout	Cutthroat: 198 mm (91-296 mm) Rainbow: 297 mm (only 1 collected) Brook: 135 mm (only 1 collected)	Moderate to high quality trout habitat.
South Fork Muddy	July 2002	Cutthroat trout (most abundant) Rainbow trout	Cutthroat: 188 mm (100-278 mm) Rainbow: 281 mm (only 1 collected)	Moderate to high quality trout habitat.
North Fork Muddy ¹	Never formally surveyed	Cutthroat trout observed in July 2002 about 2 miles above the confluence with Muddy Creek. Unidentified trout species observed in stream in summer 2002 and 2003.		Moderate to high quality trout habitat.
North Fork of Quitchupah Creek (one location)	October 2001	No fish observed at either location.	No fish collected	Unsuitable above Forest Road 007 because of erosion, siltation, and low water flows. Potentially suitable below road due to suitable flow and riparian area.

¹ Survey planned for this waterbody in summer 2004.
Data source: State of Utah Natural Resources, Division of Wildlife Resources, Southeastern Region. Louis Berg, Regional Aquatic Program Manager. Berg 2002a, 2002b, and 2002c, and Hart and Berg 2003.

Waterbody	Avg. Stream Width (ft)	Avg. Stream Depth (ft)	Substrate Rating	Cover Rating
Muddy Creek	24.6	0.57	Excellent >75% gravel/cobble/ boulder, <25% sand/silt	Poor <25% of stream shaded
South Fork Muddy	14.9	0.50	Excellent >75% gravel/cobble/ boulder, <25% sand/silt	Poor <25% of stream shaded
North Fork of Quitchupah Creek (at FS Road 007 crossing)	3.5	0.30	Good >50% gravel/cobble/ boulder, <50% sand/silt	Fair-good >25% of stream shaded, possibly >50% of stream shaded

2.4.1.3 Northern goshawk

Surveys for northern goshawks were conducted in suitable foraging and nesting habitat in the analysis area from June 21 to July 19 in 2001, July 9 to August 1 in 2002, and July 14 to July 17 in 2003. The survey periods were selected to coincide with portions of the nestling period in 2001 and the post-fledgling dependency period in 2002-2003. Suitable goshawk habitat was defined as gentle to moderate slopes (0-30%) containing mature conifer stands with closed canopies and open understories. Aspen stringers near perennial streams were also considered suitable nesting habitat, regardless of the grade of the adjacent slopes.

2.4.1.3.1 Survey Methods

Survey methods outlined in the draft Region 4 survey protocol for northern goshawks (USDA-FS 1993) were used to determine the presence of goshawks in the analysis area. The method employed the use of a conspecific broadcast calling tape of an adult northern goshawk alarm call and a portable tape player with external speakers capable of broadcasting at 90-100 decibels at one meter from the source. The taped alarm calls were played back as repeated sequences of 10 seconds of alarm calls followed by a 30-second observation period. At each broadcast station, the observer played a total of three sets of the 10 seconds of alarm call/30 seconds of observation sequence, rotating 120 degrees after each set. During all observation periods, the surveyor scanned the area vertically and horizontally for goshawks. Calling surveys were conducted during daylight hours, primarily from sunrise until early afternoon (about 14:00). No surveys were conducted in inclement weather (i.e., rain or wind).

Survey routes were determined using transects drawn on USGS topographic maps over areas containing suitable goshawk nesting and foraging habitat. Suitable habitat was determined by examining Forest Service vegetation coverage maps and aerial photos, and by field reconnaissance. Survey calling stations were located at approximately 900 foot intervals along transects to promote effective coverage of suitable habitat.

A digital coverage of the goshawk calling points and positive responses was created in Microsoft Excel. This coverage is being provided in conjunction with this report. The coverage was mapped in ArcView and overlaid onto the existing Forest Service vegetation layer to better assess distribution of positive responses in relation to habitat type (see Figure A-2, Appendix A). Results of the surveys are discussed below.

2.4.1.3.2 Survey Results

In 2001, 110 calling points were surveyed for northern goshawks. In 2002, these calling points were resurveyed and 102 new points were surveyed because of the additional potential habitat identified from vegetation maps or during field visits. In addition, goshawk alarm calls were broadcast at 59 of the stations surveyed for three-toed woodpecker between June 8 and June 23, 2002, to see if a response could be prompted earlier in the breeding season. Of these stations, 52 were resurveyed later in the year as part of the regular goshawk survey effort. In 2003, stations visited for the first time in 2002 were revisited, except where habitat was deemed unsuitable, so that each station with suitable habitat was visited twice over the 3-year study period.

Positive responses were received at three calling stations in the Muddy analysis area in 2001, two calling stations in 2002, and one calling station in 2003. One of the responses in 2002 was at a station where a response was also received in 2001. Goshawk responses were associated with mixed conifer/aspen forest at elevations above 8,650 feet. Positive responses were either silent fly-overs or vocalizations of adult birds. Of the responses in 2001, two were silent fly-overs and the third bird was flushed from a tree. Two of these responses were most likely from the same individual as the goshawk flew from the direction of the last positive calling station within fifteen minutes of the original response. Of the two responses in 2002, one goshawk flew in and vocalized and perched on a tree, the other one vocalized but did not fly in. In 2003, the response consisted of a single alarm call. Four of the responses were in the forested patches west of the tract, inside the buffer, and two were inside the tract. The number of responses observed does not represent the number of individuals. The responses were likely from individuals of one (or possibly two) goshawk pair or family unit. No physical nests were observed but it was assumed that there was at least one active nest in the area. In addition to the surveys, one incidental visual observation was made in September 2003, when an adult goshawk was seen flying over Black Fork Creek then disappearing into the forest canopy on the other side of the creek, near its confluence with the South Fork Muddy Creek.

In summary, two years of surveys were completed for all suitable goshawk habitat within the analysis area during the 3-year survey effort. Positive responses were received at five calling stations, including 2 responses at one of the

stations, for a total of 6 responses. No nests or juveniles were found but it is assumed that there was at least one active nest in the survey area.

2.4.1.4 Flammulated owl

Surveys for flammulated owls were conducted in suitable foraging and nesting habitat in the analysis area from June 20 to July 19 in 2001, June 7 to June 26 in 2002, and on June 25, 2003. Survey periods were selected to correspond with the breeding season when male owls were most likely to vocalize to signal their occupied territory. Suitable owl habitat was originally defined as pine woodlands, especially ponderosa pine. After owls were heard in alternative habitat types in the survey area, this definition was expanded to include mixed conifer forest stands that included an aspen and pine component.

2.4.1.4.1 Survey Methods

A forest owl inventory protocol (UDWR 1992), developed for use in the Northern Region and received from the Utah Department of Natural Resources, was followed to determine the presence of flammulated owls in the analysis area. The survey method employed the use of a conspecific broadcast calling tape of male flammulated owl vocalizations. The same equipment used for the goshawk surveys was used for owl surveys. Surveys consisted of broadcasting repeated sequences of a 30-second adult owl call followed by a 15-second period of silence. Twenty minutes were spent at each survey station. After listening for owls for 3 minutes, calls were broadcast in four directions, rotating 90 degrees every 4 minutes, approximately. Broadcast calling surveys began one-half hour after dusk, and continued throughout the night, as late as 3:30 am. Surveys were aborted or not conducted if it was rainy or windy.

Survey routes were determined using transects and isolated points, drawn on USGS topographic maps over areas containing suitable flammulated owl nesting and foraging habitat. Suitable habitat was determined by examining Forest Service vegetation coverage maps and aerial photos, and by field reconnaissance. Survey calling stations were located at approximately 0.5-mile intervals along transects, and 0.5 miles apart when isolated, to promote effective coverage of suitable habitat.

A digital coverage of the flammulated owl calling points and positive responses was created in Microsoft Excel. This coverage is being provided in conjunction with this report. The coverage was mapped in ArcView and overlaid onto the existing Forest Service vegetation layer to better assess distribution of positive responses in relation to habitat type (see Figure A-3, Appendix A). Results of the surveys are discussed below.

2.4.1.4.2 Survey Results

In 2001, surveys were conducted at 38 calling points. In 2002, calling points that had no response in 2001 were resurveyed, and 15 new points were surveyed to cover additional potential habitat. These new points were resurveyed in 2003 unless a positive response was recorded in 2002.

Owls responded to broadcast calls at a total of 26 stations in the analysis area over the 3-year survey period. Twelve flammulated owl responses were elicited from surveys conducted in 2001, twenty owl responses in 2002, and one in 2003. A flammulated owl was also heard incidentally along Cowboy Canyon in May 2003, at a location where no response had been recorded during the 2001 and 2002 surveys. At some of the stations, more than one owl responded to calls during the survey visit. The habitat surrounding the stations where positive responses were recorded can be categorized into ponderosa pine, limber pine and aspen, and mixed conifer and aspen forest types. The number of responses does not indicate the number of owl pairs on the tract, as unpaired male flammulated owls are thought to call more during the breeding season than paired owls. However, we can assume that numerous pairs are utilizing the survey area based on the high number of male responses and the presence of suitable habitat throughout the tract. In addition to flammulated owls, great-horned owls occasionally responded to the broadcast calls. Calling effort was aborted at these stations once the great-horned owl was heard. A saw-whet owl was also heard calling on the tract during a flammulated owl survey in 2001.

In summary, two years of surveys were completed for all suitable flammulated owl habitat within the study area during the 3-year survey effort. Positive responses were received at 26 calling stations in the survey area, for a total of 33 responses, as more than one owl responded at some of the stations. Some of these responses were likely from the same individuals as they were received at adjacent calling stations. It is assumed that several pairs of flammulated owls occur in the analysis area.

2.4.1.5 Three-toed woodpecker

Surveys for three-toed woodpeckers were conducted in suitable foraging and nesting habitat in the Muddy analysis area from June 21 to July 19 in 2001 and May 24 to June 23 in 2002. Survey periods were selected to coincide with the nest excavation period. Suitable woodpecker habitat was defined as spruce-fir forests, especially those that had been recently infested by bark beetles.

2.4.1.5.1 Survey Methods

A modification of the northern three-toed woodpecker inventory protocol, received by the Forest Service in 2001 (UDWR and UNHP 1992) was used to conduct surveys. The method employed the use of a con-specific broadcast calling tape of an adult three-toed woodpecker call plus three intermittent episodes of drumming and silence. The tape was played using a portable tape player with external speakers capable of broadcasting at 90-100 decibels at one meter from the source. The tape was played back as a repeated sequence of 10 seconds of calls and drumming followed by a 30-second observation period. At each survey station, the observer broadcast the call in all four cardinal directions for a total of eight minutes (2 minutes in each direction). During the observation periods, the surveyor scanned the area vertically and horizontally for woodpeckers and listened for drumming responses. Calling surveys were conducted during daylight hours, primarily from sunrise until early afternoon, when it became too warm (about 14:00). No surveys were conducted in inclement weather (i.e., rain or wind).

Survey routes were determined using transects drawn on USGS topographic maps over areas containing suitable three-toed woodpecker nesting and foraging habitat. Suitable habitat was determined by examining Forest Service vegetation coverage maps and aerial photos, and by field reconnaissance. The majority of transects traversed for goshawk surveys were used also for woodpecker surveys, because of the similarity of habitat requirements. When the same transects were used for both species, the survey calling stations were located at approximately 900 feet intervals. When transects were used to survey only for woodpeckers, the calling stations were located at intervals ranging between 0.15 and 0.25 miles, depending on terrain and forest density. If surveys were conducted for woodpeckers and goshawks during the same time interval, the woodpecker calls were always broadcast first.

A digital coverage of the three-toed woodpecker calling points and positive responses was created in Microsoft Excel. This coverage is being provided in conjunction with this report. The coverage was mapped in ArcView and overlaid onto the existing Forest Service vegetation layer to better assess distribution of positive responses in relation to habitat type (see Figure A-4, Appendix A). Results of the surveys are discussed below.

2.4.1.5.2 Survey Results

In 2001, 98 calling stations were surveyed, which comprised the best habitat in the tract. In 2002, 96 new calling stations were surveyed, which contained a variety of habitat including poor (canyon edges), marginal, and suitable habitats. In addition, 32 calling stations visited in 2001 were resurveyed in 2002 because they were originally surveyed after July 15, towards the end of the nest excavation period.

Fifteen woodpecker responses resulted at twelve calling stations in 2001. Six of these responses were paired, and one was incidental, occurring in the same location as a prior positive response. One woodpecker responded during the survey effort in 2002, and an additional one was observed incidentally during a goshawk survey in mid-July, when an adult female was observed drumming and flying. No nest sites were found. Positive identifications were made visually and aurally and were associated with dense forested habitats above 8,800 feet that contained patches of snags. Woodpecker responses consisted of drumming or a combination of flying to a few different trees and snags and drumming. All of the responses were in the spruce beetle infested forest habitat within the survey buffer, west of the tract boundary. Although, for a given year, it is difficult to determine the exact number of individuals observed during surveys, it can be assumed that there were at least three, and potentially more, woodpecker pairs nesting on the tract. This assumption is made based on the number of pairs observed, timing of and distance between positive survey responses, and home range size.

In summary, all suitable three-toed woodpecker habitat was surveyed at least once between 2001 and 2002. Responses were received at 13 calling stations in the Muddy analysis area, for a total of 16 responses, as both birds in a pair responded at three stations. Additionally, a female was observed incidentally in the area during a goshawk survey. It is assumed that three or more nesting pairs were present during the survey period.

2.4.1.6 Spotted bats

Surveys for spotted bats were conducted by Cirrus in potential roosting habitat in the Muddy analysis area in 2001 and 2002. Spotted bats have been recorded in a variety of habitats, including open ponderosa pine, desert shrub, pinyon-juniper, and open pastures and hay fields. Foraging occurs in riparian areas and open meadows with wet seeps or wetlands. Roosting habitat is more restrictive, being confined to rock crevices or overhangs associated with large cliff faces. Roosting habitat for spotted bats is abundant in vertical cracks of the sandstone cliff faces of the steep canyons in the tract. The riparian habitat and forest edges in the tract also provide potential foraging opportunities.

2.4.1.6.1 Survey Methods

Surveys for spotted bats in 2001 and 2002 consisted of structural searches of rock crevices or overhangs of cliffs that potentially support roost sites. Due to the hazards of steep cliff terrain, searches were confined to areas accessible safely by foot. No rock-climbing or repelling gear was used. The following structures were searched for bat roosts in the analysis area: Muddy Canyon east of Box Canyon, Greens Canyon, the East Fork of Box Canyon, the head of Box Canyon, and the North Fork of Quitcupah Canyon.

In addition to roost site searches, incidental observations of spotted bats were recorded. Spotted bat vocalizations are audible. The only other audible bat species in Utah, Allen's big-eared bat and the big-free-tailed bat, do not occur within the range of the project area (Oliver 2001). Therefore, when audible bat detections were made, it was assumed that the species heard was the spotted bat. UTM coordinates were recorded for these observations and were used to create the digital coverage provided in conjunction with this report.

2.4.1.6.2 Survey Results

No roosting sites or sign of bats were found during structural searches in the analysis area.

Numerous spotted bats were identified in the survey area by audible vocalizations or a combination of vocalizations and visual detection. Observations were made primarily in conjunction with nighttime flammulated owl survey efforts. A total of 36 spotted bat observations were recorded. Observations were associated with the rocky cliff habitat and ponderosa pine along the east fork and main stem of Box Canyon and along Greens and Cowboy Canyons. Bats were also observed foraging in the limber pine habitat near Julius Flat Reservoir and above the North Fork of Muddy Creek, and in the limber pine/Douglas fir habitat along the jeep trail running west and south of Cowboy Creek. Bat observations were mapped in ArcView and overlaid onto the existing Forest Service vegetation layer to better assess their distribution in relation to habitat type for the analysis area (see Figure A-5, Appendix A).

See section 2.4.1.8 for results of additional bat surveys conducted by other parties.

2.4.1.7 Western big-eared bats

Surveys for western big-eared bat (also known as Townsend's big-eared bat; *Corynorhinus townsendii*), were conducted by Cirrus in potential hibernacula and roosting habitat in the Muddy analysis area in 2001 and 2002. Western big-eared bats have been recorded in juniper/pine forests, shrub/steppe grasslands, deciduous forests, and mixed coniferous forests. They roost in hibernacula within caves, abandoned mine shafts, and occasionally in old buildings. Winter hibernacula for big-eared bats is very scarce in the analysis area due to the scarcity of suitable caves and open mine shafts. Cliff overhangs and shelter caves eroded in the sandstone cliffs could provide potential summer roosting habitat.

2.4.1.7.1 Survey Methods

Before surveys were initiated, the status of mines in the area was determined. One inactive mine, The Richie Mine, is present in the survey area. This mine was visited and determined closed because of lack of visible openings. The Link Canyon mine, just south of the buffer boundary is closed, with the exception of a small magazine, roughly 4 by 4 by 8 feet. One active mine, the Sufco mine, is present within and west of the Muddy analysis area, with its operating facilities southwest of the tract buffer boundary.

Surveys for western big-eared bats in 2001 and 2002 consisted of structural searches of rock overhangs and magazines that potentially support roost sites or hibernacula. These surveys were conducted concurrently with those for spotted bats and included searches of accessible structures in Muddy Canyon east of Box Canyon, Greens Canyon, the East Fork and head of Box Canyon, and the North Fork of Quitcupah Canyon. The Richie Mine and tramway were visited to determine the status of the mine and look for potential hibernacula. A rock overhang exists at the head of Box Canyon and was surveyed for sign of bat roosts. In addition, the magazine at Link Canyon Mine

was surveyed for roosting bats. Because big-eared bats are highly susceptible to temperature changes and disturbance, they would not occupy an active mine, thus the Sufco mine was not considered for the survey effort.

2.4.1.7.2 Survey Results

No roosting sites, potential hibernacula, or bat sign were found during structural searches in the tract. No substantial caves were observed on the tract and no other structures were considered potentially suitable for western big-eared bat hibernacula. No mine openings were found at the Richie Mine site after an extensive search of the area, and no bat sign was observed. This mine was considered closed. No bats or bat sign were observed in the magazine in Link Canyon Mine. Because of the disturbance of this opening by cattle and vehicular traffic, it is unlikely that western big-eared bats would roost there. Since no big-eared bats and little suitable habitat were observed, a digital coverage for this species was not created.

See section 2.4.1.8 for results of additional bat surveys conducted by other parties.

2.4.1.8 Additional Bat Surveys

2.4.1.8.1 Cooperative Challenge Cost Share Project

A general inventory was conducted for spotted bats in selected areas on the Ferron Ranger District in 1992 as part of a cooperative challenge cost share project between the MLNF and Utah Natural Heritage Program (Toone 1993). The survey methods employed consisted of listening for spotted bat audible echolocation sounds and categorizing them as a "bat pass" or a "foraging buzz", represented by the rate of echolocation heard, and mist netting. Survey locations were by Quichupah Creek, just south of the Muddy tract boundary.

No bats were captured in mist nets at Quichupah Creek, but audible bat detections were made on two occasions during the netting period. Bat activity was noted as low at this site.

2.4.1.8.2 SUFCO Mine Bat Survey

Surveys for spotted and western-big-eared bats were conducted in 1997 in Link, Muddy Creek, and Box Canyons as part of the SUFCO and Dugout Canyon Mine's permit requirements (Perkins and Peterson 1997). Four different survey methods were employed, including structure searches, mist netting, bat detectors, and audible bat transects. The survey area overlapped with the eastern portion of the Muddy tract and buffer, therefore, the SUFCO survey results are likely indicative of the species composition in the analysis area.

Results of the structure searches yielded no bats or bat sign in any shelter caves or in the Link Canyon magazine. No habitat suitable for western big-eared bats was observed. Mist netting resulted in the capture of California myotis (*Myotis californicus*) and Yuma myotis (*M. yumanensus*) in Link Canyon, and no species at Muddy Creek. The bat detector surveys resulted in the detection of spotted bats, as well as numerous other bat species including California myotis, Yuma myotis, big brown bat (*Eptesicus fuscus*), silver-haired bat (*Lasiycteris noctivagrans*), small-footed myotis (*M. ciliolabrum*), long-eared myotis (*M. evotis*), little brown bat (*M. lucifusus*), western pipistrelle (*Pipistrellus hesperus*), and an unidentified *Myotis* species. No western big-eared bats were detected. Transect surveys resulted in detection of spotted bat calls in nearly all stations in lower Box Canyon and throughout Muddy Canyon. No calls were heard in the upper reaches of the three canyons or in Link Canyon.

The results of the surveys conducted by Perkins and Peterson (1997) suggest that these canyons do not contain suitable habitat for western big-eared bats; suitable structures for day roosting and hibernacula are absent. Cliff habitat below the rims of Muddy Creek Canyon and the lower reaches of Box Canyon surveyed by Perkins and Peterson (1997) appear to provide ample habitat for spotted bats.

2.4.2 Management Indicator Species

2.4.2.1 Golden eagle (UPDATE)

Surveys for golden eagles were conducted in May 2002 in suitable nesting habitat in the Muddy analysis area with the assistance of UDWR. Additional golden eagle survey data was acquired for this area from UDWR for the period between 1998 and 2003. Suitable habitat was defined primarily as tall cliffs and occasionally trees.

2.4.2.1.1 Survey Methods

Aerial helicopter surveys were conducted annually during the eagle breeding season by UDWR. Survey methods included flight transects over and adjacent to suitable cliff habitat and some forested habitat. When nests were observed, the UTM's of the location were recorded, and the type (species) and status (active, inactive, tended) of the nest were determined, if possible. Cirrus participated with the survey effort in 2002.

A digital coverage of the flight lines and nest locations was created in Microsoft Access. A map was created for the analysis area which depicts the location and status of golden eagle nests for each year surveyed (see Figure A-6, Appendix A). Results of the surveys are discussed below.

2.4.2.1.2 Survey Results

In 2002, 12 golden eagle nests were known in the analysis area, of which 11 were surveyed. Of these, none were active, two were tended, seven were inactive, one was dilapidated, one was not found, and one was not surveyed. No eagles were seen during this survey effort. In 2003, eight of the known nests were surveyed. Of these, none were reported as active, but two were tended. No new golden eagle nests were identified during this survey effort and no golden eagles were observed. Results from surveys between 1998 and 2003 are depicted below in Table 4. Of all the golden eagle nests in the analysis area, one has been active at least once over the last six years and seven additional nests have been tended at least once.

Incidental sightings of golden eagle individuals were made during summer 2002. One observation was made of an eagle soaring over the North Fork Muddy Creek near the junction with the South Fork. The other observation was of a golden eagle perching and vocalizing on a tree and then soaring over the main stem of the Muddy Creek about one mile below the confluence of the South and North Forks. One individual was seen flying over the North Fork of Muddy Creek in July 2003 and several eagle observations were made over the main stem of Muddy Creek in 2002 and 2003.

Species	Nest Status	2003	2002	2001	2000	1999	1998
Golden eagle	Active	0	0	0	0	0	1
	Tended	2	2	2	2	1	1
	Inactive	6	7	7	5	4	4
	Dilapidated	0	1	0	0	1	1
	Not found	0	1	1	0	1	0
	Not surveyed	4	1	0	2	2	0
	Total		12	12	10	9	9

2.4.2.2 Mule Deer

Data on mule deer was acquired for the Muddy analysis area from UDWR.

The Muddy analysis area contains winter and summer range for mule deer. The value of this range is classified as high summer and high winter. The range combined covers over 90 percent of the analysis area. The extent of these ranges within the analysis area is depicted in Figure A-7 in Appendix A and is provided as a digital coverage in conjunction with this report.

No true migration routes have been identified by the UDWR or Forest Service in the analysis area. Since the area contains both winter and summer range, movements are not extensive, and follow the seasons. All areas are used, and the animals move from place to place as necessary.

Fawning areas also have not been identified, studied, reported, or mapped by these agencies. It is assumed that fawning potentially occurs in all suitable habitats. In the analysis area, suitable fawning habitat coincides with coniferous forests, mixed aspen coniferous forest, young aspen stands, and mountain brush and mahogany cover

types, with the latter being preferred because of the protective cover it affords. These habitats occur in the western and southern portions of the analysis area. Deer fawning habitat overlaps elk calving habitat to some degree.

2.4.2.3 Elk

Data on elk was acquired for the Muddy analysis area from UDWR.

The Muddy analysis area contains winter and summer range for elk. The value of this range is classified as critical summer and critical winter. The range combined covers over 90 percent of the analysis area. The extent of these ranges within the analysis area is depicted in Figure A-8 in Appendix A and is provided as a digital coverage in conjunction with this report.

No true migration routes have been identified or mapped by the UDWR or Forest Service in the analysis area. Since the area contains both winter and summer range, movements are not extensive, occur between ranges, and follow the seasons. All areas are used, and the animals move from place to place as necessary. In some years they move greater distances than others, depending on the weather and available resources.

Calving areas also have not been identified, studied, reported, or mapped by these agencies. It is assumed that calving potentially occurs in all suitable habitats. In the analysis area, suitable calving habitat coincides with aspen forests and mountain brush and mahogany cover types. Predicted calving areas occur in the southwestern portion of the analysis areas, as far north as Julius Flat Reservoir, and as far south as the North Fork of Quitchupah Creek (Hodson 2004). Potential calving habitat was modeled by the MLNF. Modeled habitat coincided with the aforementioned predicted habitat, but also occurred in the northwest portion of the analysis area. In total, modeled habitat occurred in about 10 percent of the analysis area, of which roughly 2.5 and 7.5 percent occurred in the tract and 2-mile buffer, respectively. Modeled calving habitat was associated primarily with aspen and aspen mixed conifer habitat types that were close to water (Jewkes 2004b).

2.4.2.4 Blue Grouse

No formal surveys were required for this species. Incidental observations were made while traversing potential habitat during goshawk, woodpecker, and amphibian surveys. Suitable habitat was defined as open stands of conifers or aspen with brushy understory. Potentially suitable habitat for blue grouse is present, primarily in the western portion of the Muddy analysis area.

In 2001, one adult was observed just off of Road 044 in the Greens Hollow area. In 2002, one adult blue grouse and five chicks were observed not far from this location. The surrounding habitat was a mix of small aspen and mountain shrubs. A group of four grouse consisting of one adult and three subadults was observed in 2002 above Cowboy Creek. The birds were flushed from underneath some shrubs. The area where they were observed was an opening containing grass and scattered shrub patches. Aspen and conifer patches were adjacent to this opening. In September 2003, four adults were observed at the edge of a clearing (campsite) lined by young aspen and shrubs, near Julius Flat Reservoir, and two adults were observed near Brush Reservoir (UTMs not recorded).

A digital coverage of the locations of grouse observations was created with Microsoft Excel. This coverage is depicted in Figure A-9 in Appendix A and is provided in conjunction with this report. A digital coverage of year-round blue grouse habitat based on known use areas in the late 1980s was created by UDWR in 2000. The area mapped as grouse habitat does not overlay the analysis area. Therefore, this coverage is not provided.

2.4.2.5 Macroinvertebrates

Surveys for aquatic macroinvertebrates were conducted in late spring and late summer/early fall in 2001, 2002, and 2003. Sampling took place in small perennial streams in the Muddy analysis area. Stream levels were much higher in the spring than in the summer or fall, as the streams received additional water from snowmelt and runoff in the spring, and only minimal additional water from rainfall throughout the summer.

Sampling methods outlined in the field protocol developed by Utah State University and the BLM National Aquatic Monitoring Center (Hawkins et. al 1998) were used to determine the abundance and diversity of stream invertebrate assemblages. A 500-micron mesh surber net was used to collect samples, and invertebrates were immediately preserved in a solution of 75% ethanol and 10% formaldehyde. Where possible, two invertebrate samples were taken from each site: a constant area sample and a qualitative sample. The constant area sample was a compilation of eight 0.09m² fixed-area samples taken from four different habitat units (e.g. riffles and runs). The qualitative sample consisted of a single 10-minute sample taken from all major habitat types in approximate proportion to their

occurrence. When water levels were too low to get an adequate quantitative sample, only a qualitative sample was taken. Physical habitat data was also recorded, for use in the computation of biotic indices.

In the Muddy analysis area, four sites were sampled that would be potentially impacted by mining activities. All four sites were within the tract boundary. In addition, a control site located outside the zone of potential impact was sampled. Site selection was based on the same criteria used to select water quality monitoring stations (relatively straight perennial stream reaches in narrow channels that were above or below reaches with bedrock substrate). Therefore, the same sites used to monitor water quality were selected for invertebrate sampling. The criteria used for selecting the control site were as follows: the stream must be 1) outside the buffer zone, 2) a perennial stream, 3) minimally impacted by human and natural disturbances, and 4) most closely reflected the conditions at the four sample sites. The site that most closely met these criteria was an unnamed stream near White Mountain Cabin, which is located on the westernmost side of the tract buffer. This site marginally met criterion number 1, in that it is just inside the buffer zone (within 0.3 miles of the boundary). However, this site is a good representation of a perennial stream in the area, and is outside of the zone of subsidence that could result from proposed mining activities. Furthermore, the stream source is well outside the buffer boundary (approximately 0.43 miles from buffer boundary and 1 mile from the sampling location). It was very difficult finding a control site that had similar substrate, adjacent vegetation communities, hydrology, and shape to the four sample sites. The selected site was the closest match, and was approved by the Forest Service hydrologist, Katherine Foster.

A digital coverage of macroinvertebrate sampling stations was created in Microsoft Excel. This coverage is depicted in Figure A-10 in Appendix A and is provided in conjunction with this report.

2.4.2.5.1 Year 2001 Surveys

Aquatic invertebrate sampling was conducted during June and August 2001 in perennial stream reaches in Greens Canyon (Site 1), Cowboy Creek (Site 2 [lower] and Site 3 [upper]), Greens Hollow (Site 4), and an unnamed stream near White Mountain Cabin (Control Site). A total of 98 invertebrate taxa were identified in the 17 samples collected in 2001. Taxa from five functional feeding groups (shredders, scrapers, collector filterers, collector gatherers, and predators) were collected, with collector gatherers representing the highest number of taxa and individuals collected. The five dominant taxa collected consisted of Baetis, Turbellaria, Orthocladinae, Pericoma, and Nemouridae, and the dominant families included Chironomidae, Baetidae, Psychodidae, and Nemouridae (not all invertebrates were identified to family). A complete list of taxa collected is included in Appendix B. Results of the 2001 survey effort are summarized below in Table 5. (Vinson 2002a.)

Table 5. Macroinvertebrate data from the Muddy Analysis Area. June and August 2001.										
	Site 1 (QN)	Site 1 (QL)	Site 2 (QN)	Site 2 (QL)	Site 3 (QN)	Site 3 (QL)	Site 4 (QN)	Site 4 (QL)	Ctrl (QN)	Ctrl (QL)
<i>June 2001</i>										
Mean Abundance ¹	--	979	732	330	1614	785	1133	564	5052	1908
Taxa Richness (# distinct taxa)	--	26	34	24	33	27	40	33	38	40
Mean EPT ² Abundance ¹	--	786	435	186	468	176	240	50	2329	877
EPT Taxa Richness (# EPT taxa)	--	11	13	11	9	9	9	5	19	18
Number of Unique Families	--	16	15	16	18	16	20	15	18	18
Shannon Diversity Index ³	--	1.73	2.73	2.33	2.56	2.28	2.69	2.43	2.68	2.74
Simpson's Diversity Index ⁴	--	0.30	0.09	0.13	0.12	0.16	0.12	0.15	0.10	0.09
Evenness ⁵	--	0.51	0.68	0.70	0.60	0.58	0.55	0.55	0.70	0.69
Hilsenoff Biotic Index ⁶	--	4.08	3.35	2.05	4.86	5.50	4.33	5.02	3.06	3.01
Richness-pollution intolerant taxa	--	7	4	5	2	4	2	1	8	9

Table 5. Macroinvertebrate data from the Muddy Analysis Area. June and August 2001.

	Site 1 (QN)	Site 1 (QL)	Site 2 (QN)	Site 2 (QL)	Site 3 (QN)	Site 3 (QL)	Site 4 (QN)	Site 4 (QL)	Ctrl (QN)	Ctrl (QL)
Richness-pollution tolerant taxa	--	1	1	0	1	1	1	1	1	1
	Site 1 (QN)	Site 1 (QL)	Site 2 (QN)	Site 2 (QL)	Site 3 (QN)	Site 3 (QL)	Site 4 (QN)	Site 4 (QL)	Ctrl (QN)	Ctrl (QL)
<i>August 2001</i>										
Mean Abundance ¹	--	163	3004	1654	--	664	1801	959	25029	17550
Taxa Richness (# distinct taxa)	--	22	25	29	--	20	25	22	33	27
Mean EPT ² Abundance ¹	--	42	2326	724	--	37	303	102	11409	7598
EPT Taxa Richness (# EPT taxa)	--	8	11	9	--	6	6	6	16	14
Number of Unique Families	--	14	11	12	--	10	12	12	17	15
Shannon Diversity Index ³	--	2.33	1.96	2.24	--	1.36	2.27	1.98	2.48	2.44
Simpson's Diversity Index ⁴	--	0.16	0.27	0.17	--	0.44	0.18	0.20	0.11	0.12
Evenness ⁵	--	0.58	0.45	0.57	--	0.44	0.54	0.65	0.72	0.69
Hilsenoff Biotic Index ⁶	--	4.16	2.68	4.38	--	1.63	4.68	3.74	2.43	2.30
Richness-pollution intolerant taxa	--	5	3	4	--	2	2	2	9	6
Richness-pollution tolerant taxa	--	1	1	1	--	1	1	1	1	1

QN = quantitative sample, QL = qualitative sample, Ctrl = control site.

Note: data in this table replaces that provided in Table 4 in the Cirrus Wildlife Surveys Year 2001-2002 Progress Report (February 2003).

¹ Mean Abundance is reported as number per square meter for quantitative samples and number per sample of unknown area for qualitative samples.

² EPT = Invertebrates from the orders Ephemeroptera, Plecoptera, and Trichoptera. These orders are commonly considered sensitive to pollution.

³ Shannon Diversity Index is a measure of community structure defined by the relationship between the number of distinct taxa and their relative abundances. Higher values indicate greater diversity.

⁴ Simpson's Diversity Index is also a measure of community structure defined by the proportion of taxa within the assemblage, giving little weight to rare taxa. Values range from 0 (low diversity) to 1 - 1/# taxa).

⁵ Evenness is a measure of the distribution of taxa within a community. Values range from 0 to 1, and approach zero as a single taxa becomes more dominant.

⁶ Hilsenoff Biotic Index values of 0-2 are considered clean, with little organic enrichment, 2-4 slightly enriched, 4-7 moderately enriched, and 7-10 polluted.

2.4.2.5.2 Year 2002 Surveys

Surveys for macroinvertebrates were conducted in May and September 2002. Sample periods were earlier and later in the season than in 2001 in order to sample during periods of greater water flow. In May, samples were collected at same sites sampled in 2001 with the exception of Site 1. Site 1 had no water in it in May 2002 so a new site, Site 1A, was placed upstream in Greens Canyon, about 100 meters above the point where the stream was no longer flowing. In September, samples were collected at the same sites sampled in May 2002 with the exception of Site 3. Site 3 had no water in it in September so a new site, Site 3A was placed downstream between Site 3 and Site 2 in

Cowboy Creek (referred to as Middle Cowboy Creek), in a location with sufficient flow (see Figure A-10, Appendix A).

Both types of samples (quantitative and qualitative) were taken at Site 1A during May and only a qualitative sample in September because of low flow. Flow was sufficient to take both types of samples at Site 2 during May and September. Both types of samples were taken at Site 3 in May but no samples were taken at this site in September due to lack of water. Site 3A was sampled instead, and only a qualitative sample was collected due to low flow. Both types of samples were taken at Site 4 in May and none were collected in September because the site had completely dried up. The Control Site had a larger volume of water than the other sites thus both types of samples were easily obtained during both sample periods.

A total of 86 invertebrate taxa were identified in the 16 samples collected in 2002. Taxa from five functional feeding groups (shredders, scrapers, collector filterers, collector gatherers, and predators) were collected, with collector gatherers representing the highest number of taxa and individuals collected. The five dominant taxa collected consisted of Turbellaria, Baetidae, Orthoclaadiinae, Chironominae, and Pericoma, and the dominant families included Chironomidae, Baetidae, Psychodidae, and Tipulidae (not all invertebrates were identified to family). A complete list of taxa collected is included in Appendix B. Results of the 2002 survey effort are summarized below in Table 6. (Vinson 2002b.)

Table 6. Macroinvertebrate data from the Muddy Analysis Area. May and September 2002.										
	Site 1A (QN)	Site 1A (QL)	Site 2 (QN)	Site 2 (QL)	Site 3 (QN)	Site 3 (QL)	Site 4 (QN)	Site 4 (QL)	Ctrl (QN)	Ctrl (QL)
May 2002										
Mean Abundance ¹	199	351	1966	1149	689	765	139	337	6117	2167
Taxa Richness (# distinct taxa)	15	28	22	28	17	13	10	17	23	28
Mean EPT ² Abundance ¹	28	202	796	300	19	11	1	12	5222	1573
EPT Taxa Richness (# EPT taxa)	5	11	9	11	4	1	1	4	11	13
Number of Unique Families	10	13	11	14	9	7	5	10	16	15
Shannon Diversity Index ³	1.48	2.43	1.96	2.30	1.87	1.43	0.92	1.87	2.04	2.25
Simpson's Diversity Index ⁴	0.36	0.14	0.18	0.15	0.22	0.36	0.61	0.24	0.18	0.17
Evenness ⁵	0.53	0.62	0.73	0.64	0.65	0.57	0.43	0.58	0.70	0.59
Hilsenoff Biotic Index ⁶	5.83	4.64	4.80	5.40	5.77	2.83	5.84	5.86	3.62	3.64
Biotic Condition Index (BCI) ⁷	56	63	59	56	49	47	52	49	125	123
Richness-pollution intolerant taxa	1	3	2	2	1	0	0	1	7	7
Richness - pollution tolerant taxa	1	1	1	1	1	1	1	1	1	1
	Site 1A (QN)	Site 1A (QL)	Site 2 (QN)	Site 2 (QL)	Site 3A (QN)	Site 3A (QL)	Site 4 (QN)	Site 4 (QL)	Ctrl (QN)	Ctrl (QL)
September 2002										
Mean Abundance ¹	--	264	1156	1374	--	1622	--	--	21540	13201
Taxa Richness (# distinct taxa)	--	17	32	36	--	28	--	--	30	29

	Site 1A (QN)	Site 1A (QL)	Site 2 (QN)	Site 2 (QL)	Site 3 (QN)	Site 3 (QL)	Site 4 (QN)	Site 4 (QL)	Ctrl (QN)	Ctrl (QL)
Mean EPT ² Abundance ¹	--	2	116	457	--	506	--	--	7889	1824
EPT Taxa Richness (# EPT taxa)	--	2	7	9	--	8	--	--	13	12
Number of Unique Families	--	8	16	18	--	10	--	--	20	17
Shannon Diversity Index ³	--	1.60	2.48	2.53	--	2.26	--	--	2.56	2.44
Simpson's Diversity Index ⁴	--	0.31	0.14	0.12	--	0.20	--	--	0.11	0.13
Evenness ⁵	--	0.58	0.57	0.64	--	0.48	--	--	0.67	0.66
Hilsenoff Biotic Index ⁶	--	0.48	4.63	3.82	--	4.18	--	--	1.92	2.93
Biotic Condition Index (BCI) ⁷	--	49	59	60	--	60	--	--	105	107
Richness-pollution intolerant taxa	--	0	5	3	--	3	--	--	7	8
Richness - pollution tolerant taxa	--	1	1	1	--	1	--	--	1	1

QN = quantitative sample, QL = qualitative sample, Ctrl = control site.
¹⁻⁶ See definitions in Table 5.
⁷ Biotic Condition Index = an index of stream quality, as defined in Vinson 2004.

2.4.2.5.3 Year 2003 Surveys

Surveys for macroinvertebrates were conducted in May and September 2003, at the same sites sampled in September 2002. Both types of samples (quantitative and qualitative) were taken at all sites during May. During September, only qualitative samples were taken at sites 1A and 2, because of low flow, and no samples were taken at site 4, which was dry. Site 3A and the Control Site had sufficient flow in September to take both types of samples.

A total of 87 invertebrate taxa were identified in the 16 samples collected in 2003. Taxa from five functional feeding groups (shredders, scrapers, collector filterers, collector gatherers, and predators) were collected, with collector gatherers representing the highest number of taxa and individuals collected. The five dominant taxa collected consisted of Orthocladiinae, Baetis, Turbellaria, Pericoma, and Chironominae, with Trombidiformes being very abundant also, and the dominant families included Chironomidae, Baetidae, Simuliidae, and Psychodidae (not all invertebrates were identified to family). A complete list of taxa collected is included in Appendix B. Results of the 2003 survey effort are summarized below in Table 7. (Vinson 2004.)

	Site 1A (QN)	Site 1A (QL)	Site 2 (QN)	Site 2 (QL)	Site 3A (QN)	Site 3A (QL)	Site 4 (QN)	Site 4 (QL)	Ctrl (QN)	Ctrl (QL)
May 2003										
Mean Abundance ¹	6426	659	1634	663	2784	1674	2690	896	6338	4454
Taxa Richness (# distinct taxa)	19	24	30	27	37	39	21	20	35	42
Mean EPT ² Abundance ¹	77	34	139	89	193	124	24	0	2161	2080
EPT Taxa Richness (# EPT taxa)	5	4	9	8	10	13	3	0	16	19

Table 7. Macroinvertebrate data from the Muddy Analysis Area. May and September 2003.										
	Site 1A (QN)	Site 1A (QL)	Site 2 (QN)	Site 2 (QL)	Site 3A (QN)	Site 3A (QL)	Site 4 (QN)	Site 4 (QL)	Ctrl (QN)	Ctrl (QL)
Number of Unique Families	10	11	16	14	17	20	10	8	19	19
Shannon Diversity Index ³	1.69	1.79	1.67	1.95	2.07	1.90	1.21	1.38	2.46	2.36
Simpson's Diversity Index ⁴	0.25	0.30	0.39	0.26	0.28	0.26	0.47	0.38	0.12	0.15
Evenness ⁵	0.67	0.46	0.37	0.47	0.38	0.50	0.49	0.54	0.69	0.61
Hilsenoff Biotic Index ⁶	5.27	3.86	4.62	3.76	4.35	3.58	6.84	6.72	3.25	2.96
Biotic Condition Index (BCI) ⁷	108	99	102	110	95	95	82	79	113	110
Richness-pollution intolerant taxa	1	0	4	2	4	6	0	0	9	11
Richness-pollution tolerant taxa	1	1	1	1	1	1	1	1	1	1
	Site 1A (QN)	Site 1A (QL)	Site 2 (QN)	Site 2 (QL)	Site 3A (QN)	Site 3A (QL)	Site 4 (QN)	Site 4 (QL)	Ctrl (QN)	Ctrl (QL)
September 2003										
Mean Abundance ¹	--	279	--	309	1277	365	--	--	20168	14626
Taxa Richness (# distinct taxa)	--	26	--	21	28	21	--	--	38	35
Mean EPT ² Abundance ¹	--	197	--	233	797	235	--	--	4917	4705
EPT Taxa Richness (# EPT taxa)	--	9	--	7	10	7	--	--	20	18
Number of Unique Families	--	13	--	10	13	9	--	--	20	20
Shannon Diversity Index ³	--	1.99	--	1.38	1.85	1.47	--	--	2.55	2.50
Simpson's Diversity Index ⁴	--	0.23	--	0.46	0.30	0.39	--	--	0.11	0.11
Evenness ⁵	--	0.52	--	0.39	0.43	0.47	--	--	0.68	0.71
Hilsenoff Biotic Index ⁶	--	3.16	--	2.11	2.36	2.69	--	--	2.62	3.21
Biotic Condition Index (BCI) ⁷	--	118	--	111	93	96	--	--	108	104
Richness-pollution intolerant taxa	--	5	--	3	3	3	--	--	12	12
Richness-pollution tolerant taxa	--	1	--	0	1	1	--	--	1	1
QN = quantitative sample, QL = qualitative sample, Ctrl = control site. 1-7 See definitions in Tables 5 and 6.										

2.4.2.5.4 Survey Summary: 2001- 2003

A total of 126 invertebrate taxa were identified in the 49 samples collected over the 3-year sampling period (2001-2003). Taxa from five functional feeding groups (shredders, scrapers, collector filterers, collector gatherers, and predators) were collected, with collector gatherers representing the highest number of taxa and individuals collected for each year of sampling. The five dominant taxa collected consisted of Turbellaria, Orthocladinae, Baetis, Pericoma, and Chironominae, and the dominant families included Chironomidae, Baetidae, Psychodidae, and

Nemouridae (not all invertebrates were identified to family). A complete list of taxa collected is included in Appendix B. Average results for the 3 years of surveys are summarized below by season in Table 8. (Vinson 2004.)

Table 8. Macroinvertebrate data from the Muddy Analysis Area. Three-year averages by season (2001-2003).

	Season	Site 1 (QN)	Site 1 (QL)	Site 1A (QN)	Site 1A (QL)	Site 2 (QN)	Site 2 (QL)	Site 3 (QN)	Site 3 (QL)	Site 3A (QN)	Site 3A (QL)	Site 4 (QN)	Site 4 (QL)	Ctrl (QN)	Ctrl (QL)
Number of samples (# years) ⁸	Spring	0	1	2	2	3	3	2	2	1	1	3	3	3	3
	Falls	0	1	0	2	2	3	0	1	1	2	1	1	3	3
Mean Abundance ¹	Spring	--	979	3313	505	1444	714	1152	775	2784	1674	1321	599	5836	2843
	Fall	--	163	--	272	2080	1112	--	664	1277	994	1801	959	22246	15126
Taxa Richness (# distinct taxa)	Spring	--	26	17	26	29	26	25	20	37	39	24	23	32	37
	Fall	--	22	--	22	29	29	--	20	28	25	25	22	34	30
Mean EPT ² Abundance ¹	Spring	--	786	53	118	457	192	244	94	193	124	88	21	3237	1510
	Fall	--	42	--	100	1221	471	--	37	797	371	303	102	8072	4709
EPT Taxa Richness (# EPT taxa)	Spring	--	11	5	8	10	10	7	5	10	13	4	3	15	17
	Fall	--	8	--	6	9	8	--	6	10	8	6	6	16	15
Number of Unique Families	Spring	--	16	10	12	14	15	14	12	17	20	12	11	18	17
	Fall	--	14	--	11	14	13	--	10	13	10	12	12	19	17
Shannon Diversity Index ³	Spring	--	1.73	1.58	2.11	2.12	2.19	2.22	1.85	2.07	1.90	1.61	1.89	2.39	2.45
	Fall	--	2.33	--	1.79	2.22	2.05	--	1.36	1.85	1.87	2.27	1.98	2.53	2.46
Simpson's Diversity Index ⁴	Spring	--	0.30	0.31	0.22	0.22	0.18	0.17	0.26	0.28	0.26	0.40	0.26	0.13	0.14
	Fall	--	0.16	--	0.27	0.20	0.25	--	0.44	0.30	0.29	0.18	0.20	0.11	0.12
Evenness ⁵	Spring	--	0.51	0.60	0.54	0.59	0.60	0.62	0.57	0.38	0.50	0.49	0.56	0.70	0.63
	Fall	--	0.58	--	0.55	0.51	0.54	--	0.44	0.43	0.47	0.54	0.65	0.69	0.68
Biotic Condition Index (BCI) ^{7,9}	Spring	--	--	82	81	81	83	49	47	95	95	67	64	119	117
	Fall	--	--	--	84	59	86	--	--	93	78	--	--	107	106
Hilsenoff Biotic Index ⁶	Spring	--	4.08	5.55	4.25	4.26	3.74	5.32	4.17	4.35	3.58	5.67	5.87	3.31	3.20
	Fall	--	4.16	--	1.82	3.66	3.44	--	1.63	2.36	3.44	4.68	3.74	2.32	2.81
Richness-pollution intolerant taxa	Spring	--	7.0	1.0	1.5	3.3	3.0	1.5	2.0	4.0	6.0	0.7	0.7	7.7	9.0
	Fall	--	5.0	--	2.5	4.0	3.3	--	2.0	3.0	3.0	2.0	2.0	9.3	8.7
Richness-pollution tolerant taxa	Spring	--	1.0	1.0	1.0	1.0	0.7	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0
	Fall	--	1.0	--	1.0	1.0	0.7	--	1.0	1.0	1.0	1.0	1.0	1.0	1.0

QN = quantitative sample, QL = qualitative sample, Ctrl = control site

¹⁻⁷ See definitions in Tables 5 and 6

⁸ Site 1 was only sampled in 2001, site 1A in 2002 and 2003, site 3 in 2001 and May 2002, and site 3A in September 2002 and in 2003. Additionally, site 4 was dry in September 2002 and 2003. Quantitative samples could not be collected at some of the sites due to insufficient flow or lack of water.

⁹BCI was only calculated in 2002 and 2003. Total alkalinity and sulfate were used to calculate this index and were not measured in 2001.

2.4.3 Species of High Federal Interest

2.4.3.1 Migratory Birds

The species of high federal interest for the Uinta-Southwestern Utah coal production region of Utah and Colorado include 22 species of migratory birds. These species are listed below in Table 9 along with the general habitat types required, elevation range, and predicted occurrence.

Species	Elevation Range (ft) ¹	Breeding Habitats ²	Occurrence Expected ³
Western bluebird	3,000-8,000	Open, riparian, or burned woodlands	Possible
Sandhill crane	3,000-10,000	Wetlands, freshwater margins	Possible during migration
Long-billed curlew	3,000-5,000	Prairies, grassy meadows near water	No
Bald eagle ⁴	3,000-8,000	Rivers, lakes, reservoirs	Observed
Golden eagle ⁴	3,000-14,000	Open mountain habitat	Observed
Peregrine falcon	3,000-10,000	Open forest and mountain habitat	Observed
Prairie falcon	3,000-14,000	Open mountain habitat, prairies	Observed
Cooper's hawk	3,000-10,000	Riparian woodlands, conifer, decid.	Observed
Ferruginous hawk	3,000-9,500	Grasslands, shrub-steppe	Possible
Great blue heron	3,000-9,000	Lakes, rivers, marshes	Possible
Merlin	3,000-9,000	Conifer, riparian woodlands, prairie	Possible
Scott's oriole	3,000-5,500 ⁶	Riparian woodlands, pinyon/juniper	Possible
Osprey	3,000-10,000	Rivers, riparian, lake	Possible
Burrowing owl	3,000-9,000	Grasslands, prairie, savanna	Possible
Flammulated owl ⁴	6,000-10,000	Pine forest, mixed conifer/aspens	Observed
Mexican spotted owl ⁵	5,500-9,000	Wooded steep-walled canyons	No
Band-tailed pigeon	5,000-9,000	Coniferous forests, pine, woodlands	Possible
Williamson's sapsucker	5,500-11,000	Montane conifer and aspen forests	Observed
Black swift	7,500-14,000	Montane forests, cliffs, waterfalls	Possible
Grace's warbler	5,000-7,500	Montane pine forests - southern UT	No
Lewis's woodpecker	3,000-8,000	Pine, mixed conifer, P/J, deciduous	No
Pileated woodpecker	- -	Conifer and deciduous forests	No

¹ Elevation range data is general - from Colorado GAP. Source: CDOW 2001.

² Sources: Ehrlich et al. 1988; NatureServe Explorer 2002; UCDC 2003; UDWR 1997.

³ Expected occurrence based on known distribution of species, known or predicted habitat in project area (Utah Gap Analysis 1997 and 1999), county record, elevation range of species, and habitat requirements. No = occurrence not expected; Possible = low possibility of species occurring, and Observed = species observed in the project area.

⁴ These species are addressed in more detail in additional sections of this report.

⁵ The Mexican spotted owl is being surveyed in the analysis area under a separate contract. No owls have been observed to date and the probability of occurrence is very low.

⁶ Dalton et al. 1990 record this species as usually occurring in submontane habitats from 5,500 to 8,500 ft.

2.4.3.1.1 Survey Methods

Migratory bird surveys were conducted in the following general habitat types, both concurrent and in addition to other required surveys: riparian, grassland and forblands, sagebrush, mixed conifer, aspen and aspen mixed conifer, ponderosa pine, mahogany and mountain brush, limber pine, and pinyon pine/juniper. Greater emphasis was given to surveying the forested and sagebrush habitat types.

Considerable time was spent traversing coniferous and mixed-coniferous forests and riparian habitats during the breeding season while surveying for sensitive species. Extensive time was also spent traversing shrub-steppe environments during the breeding season while surveying for sage-grouse. Time surveying in grassland habitat in 2001 to 2002 was limited to those areas adjacent to sagebrush habitat, and to the grasslands traversed while traveling to other habitats. In 2003, survey effort specifically included grasslands.

A digital coverage was created in Excel for observations of species of high federal interest not addressed elsewhere in this document. Only nests, young, or observations of adults exhibiting nesting/territorial behavior were mapped. Observations made outside of the breeding season were not mapped. This coverage is depicted in Figure A-11 in Appendix A and is provided in conjunction with this report. See Figures A-3 and A-6 for depictions of flammulated owl responses to surveys and golden eagle nest sites, respectively.

2.4.3.1.2 Survey Results

Of the migratory bird species listed above, five are not expected to occur in the analysis area, ten have a possibility of occurring, and seven were observed in the project area during 2001-2003. The species observed include bald eagle, golden eagle, peregrine falcon, prairie falcon, Cooper's hawk, flammulated owl, and Williamson's sapsucker. Observations of bald eagles, flammulated owls, and golden eagles are discussed in sections 2.4.1.1, 2.4.1.4, and 2.4.2.1, respectively.

A pair of peregrine falcons was observed during the survey effort in 2002. The falcons were encountered while walking along the rim of Muddy Creek Canyon within the 2-mile buffer on the west side of the tract (see Figure A-11, Appendix A). The pair was exhibiting territorial behavior, and it was presumed that a nest was nearby within the cliff faces. One peregrine falcon was observed circling above an inactive golden eagle nest during UDWR aerial surveys in 2003. No falcons were observed in 2001.

Three known prairie falcon nests occur in the tract buffer and have been surveyed with helicopters intermittently by UDWR since 1998 (see Figure A-11, Appendix A). All three nests were surveyed with the assistance of Cirrus in 2002. Two were inactive and one was not found. None of the surveyed nests have been active or tended since 2001, at which time one was active. Over the six-year survey period, one of the nests was active twice and another nest was tended twice. The third nest was never recorded as either active or tended. In 2001, a prairie falcon was observed by Cirrus during pedestrian surveys (location unknown). One prairie falcon was observed across the canyon near Buzzard Bench during the 2002 aerial survey effort. No prairie falcons were observed in 2003.

An adult Cooper's hawk was observed in 2002 during a goshawk survey along the North Fork of Quitcupah Creek, in the southwest portion of the analysis area. It started vocalizing and flew in after goshawk calls were played. Habitat was a drainage bottom with spruce, aspen, and a few limber pines.

One female Williamson's sapsucker was observed while conducting three-toed woodpecker surveys in 2002. It was observed first on a north-facing embankment above a tributary of the North Fork Muddy. The embankment contained a mixture of spruce, fir, and aspen and snags. The sapsucker was then observed entering a nest tree. The nest was in a 14 inch dbh aspen snag within an old beaver pond site.

Although coniferous and mixed-coniferous forest, riparian habitats, and canyon edges were frequently traversed during the breeding season, no observations of band-tailed pigeon or black swift were made. The majority of the forested portions of the analysis area above 9,000 feet, thus it is possible that this habitat is above the elevation usually used by band-tailed pigeons. Although band-tailed pigeons have been recorded using coniferous forests, they are most closely associated with Gambel oak-pinyon pine habitat types in Utah (NatureServe Explorer 2002), which are lacking in the project area. Habitat mapped by UDWR as suitable for this species is over 4 miles west of the Muddy buffer boundary (UCDC 2003). The elevation of the analysis area is within the range of that used by black swifts, and there was a report of a swift west of Joe's Valley Reservoir in 1998. However, the black swift is considered extremely rare in Utah, and its breeding habitat is limited to wet cliff ledges behind waterfalls, which have not been observed in the analysis area. Western bluebirds were not observed, although they could potentially occur in the riparian areas in the analysis area. Merlins were not observed in coniferous forest or riparian habitats, and there are no records of merlins in Emery, Sevier, or Sanpete counties, and parts of the analysis area are above the upper elevation range used by this species.

Great blue herons were not observed in the survey areas. However, this species could potentially use habitat at Julius Flat Reservoir, Brush Reservoir, or some of the beaver ponds and associated wetlands located in the western and northwestern portions of the tract buffer. Sandhill cranes could potentially use these areas too, but use would be associated with migration only, as breeding populations of this species are restricted to northeast Utah.

Survey efforts in shrub-steppe environments and grasslands resulted in no observations of burrowing owls or ferruginous hawk. Grasslands in the study area tend to be small and interspersed with shrubs and may not provide enough open habitat for these species.

Survey efforts in pinyon-juniper habitats and riparian woodlands resulted in no observations of the Scott's oriole. These habitats are extremely limited in the analysis area, thus reducing the likelihood of this species presence. Furthermore, it is likely that this species inhabits elevations lower than those present in the analysis area.

2.4.4 Other Wildlife Species

2.4.4.1 Sage-grouse

Surveys for the greater sage-grouse consisted of visits to known lek sites and searches for grouse sign in suitable habitat that could potentially be used during the breeding season. Grouse sign was in the form of tracks, fecal and cecal pellets. Suitable habitat was defined as plains, foothills, and mountain valleys, where the predominant shrub species is sagebrush, of short to medium stature. Suitable habitat for sage-grouse exists in both tracts, and although populations have greatly declined, this species was historically abundant in the area. Additionally, 48 sage-grouse were transplanted to the southern portion of the Muddy analysis area by UDWR between 1987 and 1990.

2.4.4.1.1 Strutting Ground Surveys

UDWR has been annually monitoring the strutting ground utilized by reintroduced grouse on the Muddy tract since 1991. This ground, referred to as Wildcat Knolls, has received use by 3 to 20 cocks on a given year with the lowest numbers observed in 2003. Cirrus personnel assisted with the survey in April 2002 and 2003 and also observed three hens in the area in 2002. This lek site is currently the only one in the analysis area known to be active. In 2003, UDWR and Cirrus personnel also surveyed additional areas identified as potential lek sites by Cirrus in 2002, along the east side of Box Canyon and near Pines Knolls. UDWR observed two cocks and two hens between Box Canyon and the East Fork of Box Canyon in early April, but the birds were not engaged in any lekking displays. No sage-grouse were observed near Pines Knolls. These two sites were revisited by Cirrus later in April but no grouse were observed.

2.4.4.1.2 Grouse Sign Surveys

Sagebrush habitat potentially suitable for sage-grouse was surveyed for sign by Cirrus. Priority was given to areas with historic grouse use and to those containing a good understory of grasses and forbs, although lesser quality habitat was also surveyed. Survey methods consisted of walking along closely spaced, parallel transects through sagebrush habitat and searching the ground for fecal and cecal pellets, feathers, and tracks. The majority of the survey effort took place in April 2002, although one area not visited at that time was surveyed in June 2002.

Digital coverages were generated to delineate the boundaries of survey areas, locations of abundant grouse sign, and the existing strutting ground within the tract. These coverages are depicted in Figure A-12, Appendix A, and are provided in conjunction with this report.

2.4.4.1.3 Survey Results

Surveys for sage-grouse sign within the Muddy analysis area took place from mid to late April in the following locations: Greens Hollow, The Pines, Julius Flat, the area from Wildcat Knolls north to the tip of Box Canyon, the area west of Box Canyon, and between Box Canyon and Greens Canyon. Additional surveys were conducted in early June between Box Canyon and East Fork Box Canyon.

There was no grouse sign observed at Greens Hollow or near Julius Flat reservoir, presumably because the sage in that area was very tall, with minimal understory grasses and forbs. Although the sagebrush was shorter between Greens Canyon and Box Canyon, only a few old piles of grouse pellets were found. As expected, abundant sage-grouse sign was found in the area around the Wildcat Knolls site, clear up to the intersection of FS roads 044 and 007. Further north from this intersection, on the west side of Box Canyon, there was very little grouse sign. One concentrated area of grouse use was found on the western portion of FS road 028 in The Pines. This area contained numerous patches of sagebrush that had been burned to increase cattle forage, and the grouse pellets were found on the edge between burned and unburned areas. The sage-grouse were probably using the taller sagebrush for roosting and the burned areas for foraging. The most extensive sign of sage-grouse outside of the Wildcat Knolls area was

between the East Fork and main fork of Box Canyon. Numerous piles of fecal and cecal pellets and a few feathers were found at the junction of and between FS roads 318 and 058. In addition, 12 roosting adults were flushed near the head of Box Canyon during this survey effort. (See Figure A-12, Appendix A.) South of this area, near Box Pond, SUFCO Mine personnel also reported seeing over a dozen sage-grouse (adults and chicks) in June, presumably looking for water.

2.4.4.2 Amphibians

Surveys for amphibians were conducted in suitable breeding habitats in the Muddy Creek analysis area in 2001, 2002, and 2003. Suitable habitat was defined as natural ponds and wetlands, and pooled habitat adjacent to streams. Man-made water holes and reservoirs containing emergent vegetation were also considered suitable habitat. Potential pond sites were identified from aerial photographs and through field reconnaissance.

2.4.4.2.1 Survey Methods

Amphibian encounter surveys were conducted in June and July 2001 within the analysis area by walking around the periphery of ponds and pools and scanning the area for amphibian adults, larvae, and/or eggs. Visual scans for amphibians were also made in streams where habitat conditions looked favorable. In addition, aural observations of the more vocal amphibian species were made during late afternoon and in the evening.

Additional amphibian habitat was assessed during the pond monitoring effort conducted for the Utah School and Institutional Trust Land Administration (SITLA) in September 2002. A total of 11 ponds were surveyed within the analysis area. Of these, five were natural basins and six were man-made. The majority of natural basins were dry during the survey effort. In addition to the survey parameters recorded for SITLA, the ponds were visually scanned for the presence of amphibians, and were assessed for habitat suitability (water depth, presence of emergent vegetation, and livestock disturbance).

Ponds that were dry during the 2001 and 2002 surveys and appeared to offer suitable habitat for amphibians were revisited in early spring of 2003, as soon as they became accessible, to survey for amphibians while these ponds still held water. Ponds where boreal toads had been observed in 2001 were also revisited in 2003, to attempt to confirm those sightings. Additional sites identified from aerial photos were also visited.

A digital coverage was generated for all natural ponds identified within the analysis area and for locations of identified amphibians. This coverage represents a comprehensive coverage of all natural ponds observed and/or surveyed for amphibians between 2001 and 2003. The natural pond coverage is depicted in Figure A-13, Appendix A and is provided in conjunction with this report. This coverage, used in combination with the stock pond coverage and wetland coverage (provided with the Surface and Ground Water and Vegetation Resources Technical Reports prepared for the Muddy Creek Tract, respectively), represents a relatively comprehensive coverage of all potential amphibian habitat in the analysis area.

2.4.4.2.2 Survey Results

Four species of amphibians were observed in the analysis area in 2001. Chorus frogs were found in a series of ponds in the western portion of the 2-mile buffer, from White Mountain Cabin to Julius Flat Reservoir and in one pond just inside the tract. Chorus frogs were also heard calling in late June at Julius Flat Reservoir. Tiger salamander larvae were abundant in a pond in The Pines, in the eastern portion of the 2-mile buffer, and were also present in four ponds in the western part of the buffer zone and in one pond just inside the tract. Boreal toads larvae were found in two ponds in the western part of the buffer zone, between White Mountain Cabin and Julius Flat Reservoir. Great Basin spadefoot toads (*Scaphiopus intermontanus*) were potentially heard calling in the southeast corner of the buffer zone from a cattle pond and a stream channel at the bottom of Box Canyon. However, the elevation of the analysis area may be above that used by this species. Very few amphibians were found inside of the tract boundary, proper, presumably because very little amphibian breeding habitat is present.

No amphibians were observed during the 2002 survey effort, presumably because it was conducted after the breeding season and the majority of the suitable habitat was dry. However, chorus frogs were heard calling at Julius Flat Reservoir. Ponds that were considered suitable amphibian habitat were recorded for future surveys during the breeding season.

In 2003, chorus frogs were observed at eight new ponds and at two ponds where they had already been observed in 2001, and tiger salamanders were observed at three new ponds and at one pond where they had been observed in

2001. All those ponds were located in the western part of the buffer zone. No boreal toads or Great Basin spadefoot toads were observed in 2003.

In summary, over the three-year survey period, chorus frogs were observed at fifteen ponds, tiger salamanders at nine ponds, and boreal toads at two ponds. Additionally, Great Basin spadefoot toads were possibly heard at two locations. Most of those ponds were located in the western portion of the buffer zone, with the exception of the two potential spadefoot toad observations and one salamander pond located in the southeast part of the buffer, and one pond located inside the tract, near its western edge.

2.4.4.3 Reptiles

No formal surveys for reptiles were conducted in the Muddy analysis area as no areas were identified that would be directly disturbed by mining facilities and mining roads, and facilities have already been built. However, Cirrus personnel traversed abundant habitat at all hours of the day, and performed informal searches on and under rocks and ledges in rock outcrops and sandstone formations. Five species of reptiles were observed incidentally during field visits in the analysis area (eastern fence lizard, western terrestrial garter snake, tree lizard, sagebrush lizard, and short-horned lizard), between 2001 and 2003. Overall, very few reptiles were observed.

2.4.4.4 Small mammals

No surveys were required for small mammals. Twenty-two species of mammals, including sixteen small mammals, were observed incidentally in the Muddy analysis area between 2001 and 2003 (See Appendix C). A few additional chipmunks, ground squirrels, and pocket gophers were observed but not identified to species. For the purpose of this analysis, small mammals include shrews, bats, small carnivores (Procyonidae, Bassariscidae, and Mustelidae, with the exception of the wolverine), rodents, and lagomorphs. Ungulates and large carnivores (Ursidae, Canidae, Felidae, and the wolverine) will not be addressed in this section.

Since no trapping was conducted, and since the nocturnal nature of many small mammals makes them difficult to observe, probability-of-occurrence analysis was conducted to determine what additional species could potentially occur in the analysis area. Factors used to determine probable occurrence included habitat requirements reported in the literature, habitat presence in the project area, and documented occurrence, through surveys, historic records, and incidental observations of individuals in or near the analysis area. Results of this analysis are discussed generally below, by habitat type, and depicted in Appendix D.

According to Dalton et al. (1990), 69 species of small mammals are likely present in the Wasatch Plateau area, where the analysis area is located. These species include 5 shrews, 15 bats, 9 small carnivores, 34 rodents, and 6 lagomorphs. However, based on the UDWR inventory of sensitive species in Utah (UDWR 1997), two of these species are not expected to occur in the project area: the red bat and the river otter. On the other hand, two additional species, the spotted bat and the grasshopper mouse, were observed by Cirrus personnel in the analysis area, and one more species, the Hopi chipmunk, could also occur there, based on predicted habitat maps found on the UDWR web site (Utah Gap Analysis 1997). This would bring the total number of small mammal species potentially occurring in the analysis area to 70 (5 shrew, 15 bats, 8 small carnivores, 36 rodents, and 6 lagomorphs), of which 16 were observed during the survey effort.

A comprehensive list of small mammal species and their habitat requirements and relative abundance in the Wasatch Plateau area is presented in Appendix C. County records of species occurrence, the presence of predicted suitable habitat, and the expected occurrence of individual species in the analysis area are also depicted. Species were included in the table if they were mentioned as occurring in the Wasatch Plateau area by Dalton et al. (1990), or if predicted habitat for these species was present on or near the analysis area, according to the maps on the UDWR web site (Utah Gap Analysis 1997), or if records of the species existed in one or more of the counties in which the analysis area was located. However, some of these species are not expected to occur in the Muddy analysis area proper. For instance, Utah prairie dogs are present in Sevier County but are not expected to occur in the analysis area due to lack of habitat. Predicted habitat for the dwarf shrew exists in the analysis area but this species is seemingly very rare, known from only four localities in Utah, and is not expected to occur in the Wasatch Plateau area.

Various habitat types are represented in the analysis area. General types include sagebrush, pinyon-juniper, mahogany and mountain brush, grassland and forbland, aspen and aspen-mixed conifers, mixed conifers, ponderosa pine, limber pine, rock outcrops and barren areas, and riparian areas, wetlands, and reservoirs. Some wildlife

species may occur in only one particular habitat type, while others may use a wide variety of habitats. Riparian and wetland areas are used by the highest number of wildlife species but represent only a very small proportion of the habitat in the analysis area. A general description of species that use each habitat type follows.

Sagebrush is one of the most widespread and abundant habitat types in the analysis area. Small mammal species using sagebrush on the Wasatch Plateau include the least chipmunk, Great Basin pocket mouse, long-tailed vole, meadow vole, black-tailed jackrabbit, white-tailed jackrabbit, desert cottontail, and mountain cottontail. Both species of jackrabbits and the mountain cottontail were observed in the analysis area. The desert cottontail occurs at lower elevations than the mountain cottontail, generally below 6,000 feet. Since the analysis area is above 6,000 feet, most cottontails present in the area are likely to be mountain cottontails.

Pinyon-juniper habitat type is rare in the analysis area. Small mammal species found in this habitat on the Wasatch Plateau include the cliff chipmunk (which was observed in the analysis area), Hopi chipmunk, Great Basin pocket mouse, canyon mouse, and pinyon mouse. The ringtail could also potentially occur, in Muddy Creek Canyon.

Mahogany and mountain brush represent the most abundant habitat type in the analysis area. Species associated with this habitat on the Wasatch Plateau include the spotted skunk, cliff chipmunk, brush mouse, canyon mouse, and mountain cottontail. The cliff chipmunk and mountain cottontail were observed in the analysis area.

Grassland/forbland habitats are relatively common in the analysis area, occurring primarily in patches adjacent to pinyon-juniper, and sagebrush habitat types. Small mammals found in these habitats on the Wasatch Plateau include the badger, yellow-bellied marmot, Uintah ground squirrel, least chipmunk, northern pocket gopher, plains pocket mouse, Great Basin pocket mouse, long-tailed vole, montane vole, meadow vole, western jumping mouse, black-tailed jackrabbit, and white-tailed jackrabbit. The badger, yellow-bellied marmot, Uintah ground squirrel, northern grasshopper mouse, both species of jackrabbits, and an unidentified pocket gopher species were observed in the analysis area.

The aspen and aspen-mixed conifer habitat type is common in the western half of the analysis area. Small mammal species using those habitats include the beaver, the porcupine, and the snowshoe hare, all of which were observed in the analysis area.

Mixed conifers (mostly Douglas fir, subalpine fir, Englemann's spruce) represent a moderate component of the analysis area and are associated primarily with the perennial drainages. Typical small mammal species inhabiting coniferous forests on the Wasatch Plateau include the northern flying squirrel, the red squirrel, the porcupine, and the snowshoe hare. The Uintah chipmunk is found in openings in coniferous forests or at forest edges. Most of these species were observed in the analysis area, with the exception of the northern flying squirrel.

Ponderosa pine represents a minor component in the analysis area, occurring primarily in the southeast portion. Limber pine is even less abundant, occurring only in small, localized areas. Small mammal species using these two habitats on the Wasatch Plateau primarily include generalist species that can be found in a variety of other habitats as well including the cliff chipmunk (observed in the analysis area), least chipmunk, northern pocket gopher, deer mouse, and long-tailed vole.

Rock outcrops and barren areas are rare in the analysis area. Several small mammals use these habitat features on the Wasatch Plateau. Species observed in the analysis area include the yellow-bellied marmot, golden-mantled ground squirrel, cliff chipmunk, Uintah chipmunk, bushy-tailed woodrat, and mountain cottontail. The spotted bat uses cliffs with rock crevices for roosting and was heard vocalizing at night. The ringtail, spotted skunk, rock squirrel, Hopi chipmunk, desert woodrat, canyon mouse, and pika could also occur in this habitat type.

Riparian areas, wetlands, and reservoirs are scarce in the analysis area but are used by the highest number of wildlife species. These habitats are present around and south of Julius Flat Reservoir, in Muddy Creek Canyon and associated tributaries, and along the North Fork of Quitcupah Creek. Typical riparian or wetland species found on the Wasatch Plateau include the masked shrew, northern water shrew, vagrant shrew, ringtail, raccoon, ermine, mink, beaver, western harvest mouse, western jumping mouse, long-tailed vole, meadow vole, water vole, and muskrat. Of these species, only the beaver was actually seen in the analysis area, but the other species are also expected to occur there. Most bat species also forage near water.

2.4.4.5 Non-game birds

No surveys were required for non-game birds. A list of bird species observed was compiled from incidental observations made during field visits to the study area in spring, summer, and fall 2001-2003. This list is presented in Appendix C. Cirrus personnel identified 90 species of birds in the Muddy analysis area, including 84 non-game birds and 6 game birds. Unidentified flycatchers and vireos were also observed.

A probability-of-occurrence analysis was conducted to determine what additional non-game bird species could potentially occur in the analysis area. Factors used to determine probable occurrence included habitat requirements reported in the literature, habitat presence in the project area, and documented occurrence, through surveys, historic records, and incidental observations of individuals in or near the analysis area. Results of this analysis are discussed generally below by habitat types described in section 2.4.4.4, and are depicted in Appendix E.

According to the information presented in Dalton et al. (1990), 201 species of non-game birds frequent the Wasatch Plateau area, excluding accidental species that are not normally found in the area. This includes 69 yearlong residents, 92 summer residents, 8 winter residents and 32 transients. Yearlong and summer residents are species that breed in the area (161 species total), winter residents breed in northern regions and only spend the winter in the area, and transients pass through the area during spring and/or fall migrations and do not remain in the area for any extended period of time. Nine of these species are unlikely to occur in the project area due to lack of suitable habitat or because the project area is outside of their range. On the other hand, six additional species were observed by Cirrus personnel in the analysis area or its vicinity, and five more could potentially be present there, based on the predicted habitat maps from the UDWR web site (Utah Gap Analysis 1997 and 1999). This brings the total number of non-game birds potentially present in the analysis area to 203, of which 84 were actually observed during the survey effort.

Some of these species, such as shorebirds associated with mudflats, are unlikely to be present on the coal tract itself but may occur locally in the buffer zone. Julius Flat Reservoir, located near the western edge of the Muddy tract buffer, has cobbly shores and does not offer any habitat for species foraging in the mud when water level is high. During late summer and fall, however, water level recedes and mudflats may become exposed. Migratory shorebird could potentially use the reservoir during fall migration. However, the only shorebird species we observed there was the spotted sandpiper.

A comprehensive list of non-game bird species and their habitat requirements, seasonal status, and relative abundance in the Wasatch Plateau area is presented in Appendix E. Upland game birds, waterfowl, and pigeons/doves were considered to be game birds and are not included in the table. Five species of upland game birds, 20 waterfowl species (10 of them transient) and 3 pigeon/dove species are also present in the Wasatch Plateau area.

Non-game bird species observed in sagebrush habitat in the analysis area included the turkey vulture, golden eagle, common poorwill, broad-tailed hummingbird, gray flycatcher, sage thrasher, and vesper sparrow. The Brewer's sparrow was also observed, even though this species was listed by Dalton et al. (1990) as not known to inhabit the Wasatch Plateau area. The rough-legged hawk is expected to occur in this habitat during winter but most of the analysis area is usually inaccessible in that season due to deep snow or mud making the roads undrivable.

Species observed in pinyon-juniper habitat included the gray flycatcher, ash-throated flycatcher, western scrub jay, pinyon jay, green-tailed towhee, and gray vireo (which was not listed by Dalton et al. (1990) as present in the Wasatch Plateau area). The bushtit, plain titmouse, blue-gray gnatcatcher, Bewick's wren, black-throated gray warbler, and Virginia's warbler could also occur in this habitat in the analysis area.

Species commonly occurring in mahogany and mountain brush on the Wasatch Plateau include the broad-tailed hummingbird, the dusky flycatcher, the western scrub jay, the black-billed magpie, the bushtit, the Virginia's warbler, the dark-eyed junco, and the spotted towhee. Of these, the broad-tailed hummingbird, western scrub jay, black-billed magpie, dark-eyed junco, and spotted towhee were observed in the analysis area, as well as unidentified flycatchers.

Species observed in grassland and forbland habitats in the analysis area included the turkey vulture, prairie falcon, short-eared owl, common poorwill, vesper sparrow, and western meadowlark, even though this last species was not listed as present on the Wasatch Plateau by Dalton et al. (1990). Other species potentially using these habitats in the analysis area include the northern harrier, rough-legged hawk (in winter), and horned lark. We surveyed some of the grasslands in the analysis area for burrowing owls but did not find any. Grasslands in the analysis area may not be open enough for this species, as most of them are small and interspersed with shrubs.

Aspen and aspen-mixed conifer habitats in the analysis area are used by the Cooper's hawk, northern goshawk, flammulated owl, broad-tailed hummingbird (near openings), northern flicker, downy woodpecker, hairy woodpecker, Williamson's sapsucker, red-naped sapsucker, olive-sided flycatcher, western wood-pewee, violet-green swallow, black-capped chickadee, red-breasted nuthatch, brown creeper, warbling vireo, yellow-rumped warbler, orange-crowned warbler, western tanager, and dark-eyed junco. The white-breasted nuthatch and solitary vireo were not observed but are also expected to occur in that habitat in the study area.

Mixed conifers provide habitat for the Cooper's hawk, northern goshawk, great horned owl, flammulated owl, northern saw-whet owl, broad-tailed hummingbird (near openings), three-toed woodpecker, hairy woodpecker, Williamson's sapsucker, red-naped sapsucker, olive-sided flycatcher, western wood-pewee, tree swallow, violet-green swallow, Steller's jay, Clark's nutcracker, gray jay, mountain chickadee, red-breasted nuthatch, brown creeper, ruby-crowned kinglet, hermit thrush, Townsend's solitaire, yellow-rumped warbler, western tanager, dark-eyed junco, chipping sparrow, pine siskin, and pine grosbeak. Additional species that were not observed but are expected to occur in this habitat include the cordilleran flycatcher, white-breasted nuthatch, golden-crowned kinglet, Swainson's thrush, Cassin's finch, and red crossbill. The Townsend's warbler could occur here during migrations.

Species observed in ponderosa pine included the flammulated owl, northern flicker, hairy woodpecker, brown creeper, and pine grosbeak. The pygmy nuthatch, solitary vireo, and Cassin's finch are also likely to be present in this habitat in the study area and the Williamson's sapsucker could occur in this habitat also (it was observed in the analysis area in a different habitat).

Species associated with rocky outcrops or barren areas included the golden eagle, peregrine falcon, prairie falcon, white-throated swift, canyon wren, and rock wren. The black rosy-finch and grey-crowned rosy-finch could also occur in the higher portions of the analysis area, in the western part of the Muddy buffer.

Of all habitats, riparian areas, wetlands, and reservoirs are used by the highest number of bird species. Some species use almost exclusively these habitats while others spend part of their time in other habitats. Most transient species use these habitats during migrations, including loons, grebes, shorebirds, waterfowl, gulls, and warblers. Two reservoirs are present in the Muddy buffer (Julius Flat Reservoir and Brush Reservoir), as well as various ponds and several perennial streams. Typical species using riparian areas, wetlands, or reservoirs in the analysis area include the sora rail, spotted sandpiper, cliff swallow, tree swallow, American dipper, MacGillivray's warbler, yellow warbler, and song sparrow. Five bald eagles were also observed along Cowboy Creek on the Muddy Tract in November 2003, as mentioned earlier in this report. The common loon, western grebe, pied-billed grebe, American white pelican, American coot, common snipe, great blue heron, California gull, belted kingfisher, barn swallow, bank swallow, northern rough-winged swallow, willow flycatcher, Wilson's warbler, and lazuli bunting were not observed but could also occur in the study area, as well as a variety of transient shorebirds.

3.0 Results and Discussion

This section provides background information necessary to assess potential impacts to terrestrial and aquatic wildlife that could occur as a result of implementing the action alternatives. The potential and/or known occurrence and habitat requirements for four categories of wildlife are discussed. The categories of wildlife addressed include the following: (1) federally listed and proposed endangered, threatened, and candidate species and Forest Service sensitive species (TEPS), (2) management indicator species (MIS), (3) species of high federal interest, and (4) other wildlife species not addressed in the previous categories, including fishes, blue grouse, amphibians, reptiles, small mammals, and non-game birds.

3.1 Description of the Affected Environment

3.1.1 Aquatic and Terrestrial Habitat

The analysis area is comprised of the Muddy coal tract (~8,645 acres) and the 2-mile buffer surrounding the tract (~28,205 acres), for an approximate total of 36,850 acres. The majority of the impacts associated with mining would be associated with the tract, although potential impacts could occur up to approximately 0.25 miles outside of the tract.

Ten wildlife habitat types are used in this analysis, of which one is aquatic and the rest terrestrial. The digital coverage of vegetation types for the MLNF (Forest Service 2002a) was used as a guide to define these types, and similar vegetation cover types in this coverage were consolidated. A brief description of habitat types used in this analysis follows. Further discussion of habitat types can be found in the Vegetation Resources Technical Report prepared for the Muddy Tract. Streams were not categorized in the MLNF digital coverage, so they are described separately under aquatic habitat.

3.1.1.1 Aquatic Habitat

The primary perennial streams in the analysis area include Muddy Creek, North and South Forks of Muddy Creek, Horse Creek, Meadow Gulch, Box Canyon, East Fork Box Canyon, The Box, and the North Fork of Quitcupah Creek. In addition, perennial flow is present intermittently in portions of Cowboy Creek, Greens Hollow, and Greens Canyon. Of these streams, only portions of Muddy Creek and its north and south forks, The Box and Box Canyon, Cowboy Creek, Greens Hollow, and Greens Canyon occur within the coal tract boundary. Most streams in the analysis area are intermittent and do not provide good quality fish habitat. Intermittent streams are unlikely to contain cutthroat trout or other fish species. A complete list of perennial streams in the analysis area is included in the Surface and Ground Water Technical Report prepared for the Muddy Creek Tract.

The analysis area contains a minor component of riparian habitat. This includes wetlands, dry and wet meadows, willow and tree dominated riparian areas, and lakes, ponds, and reservoirs. Springs and seeps are also present. Combined, these habitats comprises less than one percent of the analysis area. However, these habitats are important for a variety of wildlife species, as most wildlife use riparian areas for at least some part of their life cycle. The extent of wetlands in the analysis area is detailed in the Vegetation Resources Technical Report prepared for the Muddy Creek Tract.

3.1.1.2 Terrestrial Habitat

The analysis area contains a variety of terrestrial habitats, including sagebrush, pinyon-juniper, mahogany and mountain brush, grassland and forbland, aspen and aspen-mixed conifers, mixed conifers, ponderosa pine, limber pine, and rock outcrops and barren areas.

Mahogany and mountain brush constitute the most abundant habitat type, occurring throughout the analysis area (27 percent). Sagebrush is the second most widespread and abundant habitat type in the analysis area, comprising roughly 21 percent of the area. The aspen and aspen-mixed conifer habitat type is common on the western half of the analysis area, comprising roughly 16 percent of the area. Grassland/forbland habitats are relatively common in the analysis area (15 percent of the area), occurring primarily in patches adjacent to pinyon-juniper and sagebrush habitat types. Mixed conifers (mostly Douglas fir, subalpine fir, Englemann's spruce) represent a moderate component of the analysis area (about 8 percent) and are associated primarily with the perennial drainages. Ponderosa pine represents a minor component in the analysis area (about 4 percent), occurring primarily in the southeast portion, outside of the tract. The pinyon-juniper and limber pine habitat types are rare in the analysis area (about 2 percent each), occurring mostly outside the tract. Rock outcrops and barren areas are also rare in the analysis area (about 2 percent), being limited primarily to the canyon walls of the Muddy drainage.

3.1.2 TEPS

The FWS annual list of federally listed and proposed endangered, threatened, and candidate species and habitat in Utah by County (FWS 2002) indicates that nine threatened or endangered wildlife species of concern and one candidate for listing could potentially occur in Emery, Sanpete, and/or Sevier counties. The Intermountain Region list of proposed, endangered, threatened, and sensitive species known or suspected distribution by Forest (Forest Service 2003b) indicates that ten Forest Service Sensitive species could occur on the MLNF. These species and their status are depicted in Table 10.

Table 10. Federally listed threatened and endangered and candidate species and Forest Service sensitive species potentially occurring on the MLNF in Emery, Sanpete, and/or Sevier counties.

Species	Status
Fishes	
Bonytail (<i>Gila elegans</i>)	Endangered (Emery County)
Colorado Pikeminnow (<i>Ptychocheilus lucius</i>)	Endangered (Emery County)
Humpback Chub (<i>Gila cypha</i>)	Endangered (Emery County)
Razorback Sucker (<i>Xyrauchen texanus</i>)	Endangered (Emery County)
Colorado River Cutthroat Trout (<i>Oncorhynchus clarki pleuriticus</i>)	Sensitive
Bonneville Cutthroat Trout (<i>Oncorhynchus clarki utah</i>)	Sensitive
Birds	
Bald Eagle (<i>Haliaeetus leucocephalus</i>)	Threatened (Emery, Sanpete, and Sevier counties)
Mexican Spotted Owl (<i>Strix occidentalis lucida</i>)	Threatened (Emery County)
Southwestern Willow Flycatcher (<i>Empidonax trailii extimus</i>)	Endangered (Sevier County)
Western Yellow-Billed Cuckoo (<i>Coccyzus americanus occidentalis</i>)	Candidate (Emery, Sanpete, and Sevier counties)
Northern Goshawk (<i>Accipiter gentilis</i>)	Sensitive
Flammulated Owl (<i>Otus flammeolus</i>)	Sensitive
Peregrine Falcon (<i>Falco peregrinus</i>)	Sensitive
Three-Toed Woodpecker (<i>Picoides ridactytus</i>)	Sensitive
Greater Sage-Grouse (<i>Centrocercus urophasianus</i>)	Sensitive
Mammals	
Black-Footed Ferret (<i>Mustela nigripes</i>)	Endangered (Emery County)
Canada Lynx (<i>Lynx canadensis</i>)	Threatened (Sanpete County)
Utah Prairie Dog (<i>Cynomys parvidens</i>)	Threatened (Sanpete, and Sevier counties)
Spotted Bat (<i>Euderma maculatum</i>)	Sensitive
Western Big-Eared Bat (<i>Corynorhinus townsendii pallescens</i>)	Sensitive
Amphibians	
Spotted Frog (<i>Rana luteiventris</i>)	Sensitive

Of the species listed in Table 10, the yellow-billed cuckoo, southwestern willow flycatcher, black-footed ferret, Utah prairie dog, and Bonneville cutthroat trout are not predicted to occur in the analysis area and are not analyzed in this document. The remaining species could potentially occur in the analysis area and are addressed in this document.

The yellow-billed cuckoo is not addressed because the analysis area is above the elevational range of this species. The southwestern willow flycatcher is not addressed because the analysis area does not contain suitable habitat and the known distribution of this species does not overlap the Ferron Ranger District or other portions of the northern region of the MLNF (Utah Gap Analysis 1997). Furthermore, two years of surveys on the MLNF have failed to locate this species. The black-footed ferret is not addressed because predicted habitat does not occur in the analysis area (Utah Gap Analysis 1997), and this species is presumed extirpated from all but the eastern portion of Utah. The Utah prairie dog is not addressed because suitable habitat does not occur in the analysis area. Suitable habitat is present below the tract buffer, near the town of Emery, however, the last record of this species in this area was in

1929 (Utah Gap Analysis 1997). The Bonneville cutthroat trout is not addressed because the analysis area is outside of the geographical range for this species.

TEPS Fish

Habitat requirements and life history characteristics of the species present within the analysis area or in the vicinity of it are described below. Special emphasis is given to TEPS. Within the analysis area, cutthroat trout is the only species listed as sensitive by the FWS and the State of Utah. No other TEPS are present within the analysis area. However, Muddy Creek flows into the Colorado River, which provides habitat to four endemic endangered species, including the bonytail, Colorado pikeminnow, humpback chub, and razorback sucker. Habitat and life history characteristics of these federally listed species are discussed briefly, as their habitat range is adjacent to the analysis area and impacts to water quality in the Muddy Creek drainage could potentially affect water quality in the Colorado River.

The results of fish surveys conducted on perennial streams by the UDWR indicated that native cutthroat trout were present in Muddy Creek and the South Fork of Muddy Creek. Cutthroat were also observed incidentally on the North Fork of Muddy, but formal surveys have not yet occurred there. This cutthroat trout is thought to be of the Colorado River subspecies based on their known distribution in Utah. No fish were observed at the North Fork of Quitcupah Creek.

3.1.2.1 Bonytail

The bonytail is a member of the minnow family (Cyprinidae) similar to the humpback chub. The historic range of this species encompassed the mainstem and large tributaries of the Colorado River. The distribution and abundance of bonytail have been reduced greatly due to flow depletions, habitat loss and alteration, predation, and competition with exotic species. In hatcheries, spawning starts at temperatures of 20 °C. Eggs hatch 4 to 7 days after fertilization. Spawning is now rare in natural environments. However, they spawn during the spring and summer over gravel substrates, and they seem to prefer eddys and pools rather than swift currents. They are opportunistic feeders with an omnivorous diet that includes insects, zooplankton, algae, and higher plant matter (Sigler and Sigler 1996).

3.1.2.2 Colorado Pikeminnow

Native to the Colorado River system, the Colorado pikeminnow (formerly known as the Colorado squawfish) is the largest American minnow. This species occurs in warm, swift waters of large rivers in the Colorado Basin. However, they can tolerate a wide temperature range from 10°C in winter to more than 30°C in the summer. They are adapted to rivers with seasonally variable flow, high silt loads, and turbulence. Adults are migratory and inhabit pools and eddies near the main current while juveniles prefer backwater areas. Spawning occurs during spring and summer over riffle areas with grabble or cobble substrate. These fishes are primarily piscivorous, but small individuals also feed on insects and other invertebrates. This species has declined drastically due to stream alteration and habitat fragmentation caused by the construction of dams, irrigation dewatering, and the introduction of competitive and predatory non-native fishes. In addition, the size and number of backwaters and sloughs used for nursery and resting areas have decreased due to channelization below dams, and the natural cycle of flood and drought has been replaced by stable discharges and water levels (Sigler and Sigler 1996).

3.1.2.3 Humpback Chub

The humpback chub is a member of the Cyprinidae family, native to the upper Colorado River. Severe population declines of this species have occurred due to the alteration of streams, which have lead to changes in turbidity, volume, current velocity, and water temperature. In addition, this fish has also been affected by predation and competition with introduced fish species, pollution and eutrophication, parasitism, changes in food sources, and fishing pressure. Fast currents and deep water over substrates of sand, silt, boulder, and bedrock have been associated with this species. Spawning occurs during spring and summer in shallow, backwater areas, with cobble substrate. Juveniles remain in these waters until they are large enough to move into the white-water areas (Sigler and Sigler 1996).

3.1.2.4 Razorback Sucker

The razorback sucker is a species native to the Colorado River system that has been greatly impacted by competition and predation from nonnative fish species, as well as by changes in natural flow and temperature regimes. This fish feed on algae, zooplankton, and other aquatic invertebrates. They occur in medium to large rivers with swift turbulent waters, as well as in slow backwater habitats and impoundments. Spawning occurs from February to June. Limited numbers of this fish species persist (Sigler and Sigler 1996). The largest current concentration of razorback suckers can be found in the Upper Green River and lower Yampa River (Tyus 1987). They also occur in small numbers in the Grand Valley area of the Colorado River (Osmundson and Kaeding 1991).

3.1.2.5 Colorado River Cutthroat Trout

The following description is based on the summary of habitat requirements and life history characteristics presented by Lentsh and Converse (1997). The Colorado River cutthroat trout (CRCT) is a subspecies of the cutthroat trout that is native to the upper Colorado River drainage of Utah, Wyoming, Colorado, Arizona, and New Mexico. This species is rare within its historic range. Habitat loss, predation, competition with non-native species, and hybridization have contributed to its population decline.

Generally, CRCTs begin to spawn when spring floods start to recede in late spring and early summer. This behavior may be triggered by changes in water temperature. Fecundity varies with individual size; a 290-mm female can lay over 600 eggs. Water temperature, elevation, and climate variations determine fry emergence, which usually occurs in late summer. Maturity is reached approximately 3 years after.

There is limited information on habitat requirements for CRCT. This species spawns over gravel substrates with good water flows. Studies have provided evidence of a positive association between CRCT presence and the amount of large woody debris, depth, and low water velocity. However, many streams that present CRCT do not present these habitat characteristics. CRCT generally feed on macroinvertebrates. Adults can also feed on other fish and eat larger proportions of large macroinvertebrates and terrestrial insects than subadults.

Introduced species may outcompete CRCT, as this species did not evolve with other salmonids. The different life history traits of non-native salmonids also poses a competitive advantage of these species over the native trout. Brook trout reach larger sizes than CRCT by their first winter season as they spawn in the fall and fry emerge early in the spring. Furthermore, brook trout mature earlier and have the potential to produce a greater number of offspring during their life span.

TEPS Birds

3.1.2.6 Bald Eagle

In Utah, the bald eagle is primarily a winter resident, with only four known pairs of nesting eagles in the state, none of which occur on the MLNF. An eagle nest does occur on private land about 18 miles east of the northeastern boundary of the analysis area, near the town of Castledale. It is unlikely that individuals from this eagle pair would utilize portions of the analysis area for foraging, since suitable habitat is available closer to the nest site. Several hundred bald eagles winter in Utah, where they typically congregate in large groups at roost sites. Wintering eagles typically begin arriving in November, are most abundant in January and February, and begin migrating north in March. Bald eagles generally utilize cottonwoods and snags near open bodies of water as winter roosting sites, and feed opportunistically on live or dead fish, waterfowl, and mammals (Beck 1980).

Only one observation of bald eagles was recorded in the analysis area during the analysis period. Five bald eagle individuals (3 adults and 2 juveniles) were sighted in November 2003 along Cowboy Creek, presumably during fall migration. Winter roosting habitat is limited in the analysis area due to the high elevation and lack of roost trees. Potential roosting habitat occurs along the lower portion of Muddy Creek, in the tract buffer. Foraging habitat is present along Muddy Creek and its main tributaries, as well as in Julius Flats Reservoir. It is likely that these waterbodies are used in late fall and early winter, before they freeze over. In general, use of the permit area would be incidental and likely in connection with fall or spring migration.

3.1.2.7 Mexican Spotted Owl

Spotted owls in Utah are generally found in the pinyon-juniper zone, below the mixed conifer forests typical of owl habitat in Arizona and New Mexico. These birds select steep, narrow, cool canyons for roosting and nesting. These sites are characterized by small clumps of true fir and deciduous trees growing within cool canyons or on steep north-facing slopes. Ponderosa pine/gamble oak forests are also used if they exhibit characteristics of large cavity trees, broken tops, numerous snags, and heavy accumulations of down woody material. During the winter, the owls tend to move out of the canyons and onto mesa-tops, benches and warmer slopes (Wiley 1992). Spotted owls apparently use a wider array of habitat types for foraging than for nesting and roosting, including fairly open and non-contiguous forest, small openings, and pure ponderosa pine stands. Little is known about the habitat requirements for dispersal. Mexican spotted owls are generally absent from high elevations. (Rodriguez 1998.)

Potentially suitable canyon habitat for spotted owls occurs in the analysis area. However, the analysis area is north of the known distribution of this species in Utah and is above the elevation generally used by this species. Suitable habitat was surveyed in 2002 and 2003 by Arizona Biological Surveys. No spotted owl responses were detected in

the survey area. The likelihood of occurrence of this species in the analysis area is very low due to elevation and geographic range.

3.1.2.8 Northern Goshawk

The northern goshawk is a forest habitat generalist that uses a wide variety of forest ages, structural conditions, and successional stages. The goshawk preys on large-to-medium-sized birds and mammals which it captures on the ground, in trees, or in the air. Selected goshawk prey includes squirrels, chipmunks, woodpeckers, jays, rabbits, and grouse. Specific habitat attributes include snags, downed logs, woody debris, large trees, herbaceous and shrubby understories, and a mixture of various forest vegetative structural stages. (Reynolds et al. 1992.)

Three components of a goshawk's home range (total about 6,000 acres) have been identified: nest area, post fledging/family area (PFA), and foraging area. The nest area is approximately 30 acres and may include one or more nests. It is usually located on northerly aspects in drainages or canyons, and is often near streams. Nest areas contain one or more stands of large, old trees with a dense canopy cover. The PFA surrounds the nest area. It totals approximately 420 acres and most often, because of its size, includes a variety of forest types and conditions. Small openings, snags, downed logs, and woody debris are critical PFA attributes. Goshawk foraging areas are approximately 5,400 acres in size. Observations of foraging goshawks show that they hunt in many forest types and conditions. This opportunism suggests that the choice of foraging habitat may be as closely tied to prey availability as to habitat structure and composition. (Reynolds et al. 1992.)

Suitable habitat is present in the analysis area, but is limited primarily to the aspen and aspen mixed conifer cover types on the western portion of the tract buffer. Two years of surveys were conducted for goshawks in suitable habitat in the analysis area (see section 2.4.1.3). Goshawks were heard and/or seen at four calling stations. No nests or juveniles were found, but it is assumed that there was at least one active nest in the area, and likely two, based on the distance between responses and size of home ranges.

The Forest Service has been monitoring two goshawk nests near or in the analysis area. One occurs north of Julius Flats Reservoir, on the edge of the northern buffer boundary. The analysis area contains a portion of the nest stand, and is within the PFA and foraging area for this pair. The other nest occurs near Meadow Gulch, about one mile north of the northeast side of the buffer boundary. The analysis area is within the foraging area for this pair. None of the goshawk responses during the survey effort occurred within the home ranges of the known goshawk pairs, indicating that these goshawks were from a different pair or pairs.

The northern goshawk was added to the list of MIS for the MLNF in June 2003 as an amendment to the 1986 Forest Plan. This species replaced the blue grouse as a MIS. One of the standards and guidelines associated with goshawk management is monitoring of territory occupancy on a Forest-wide basis. Less than 20 percent decline in territory occupancy over a 3-year period is considered acceptable for this species. Monitoring efforts conducted since 1992 for the northern division of the MLNF indicate that territory occupancy has been at least 30 percent and thus within an acceptable range for this species (Jewkes 2004a). However, breeding bird trend studies for the state of Utah indicate that this species has been decreasing throughout its range since 1966 (Sauer et al. 2003).

3.1.2.9 Flammulated Owl

Flammulated owls occur in mixed pine forests, from pine mixed with oak and pinyon at lower elevations to pine mixed with spruce and fir at higher elevations. They have also been found in aspen, second-growth ponderosa pine, and mixed coniferous forest. Preferred habitat is mature ponderosa pine/Douglas-fir forests with open canopies. Large diameter (>20 inch dbh) dead trees with cavities at least as large as northern flicker cavities are important site characteristics. Flammulated owls are strictly nocturnal and feed almost exclusively on insects. Foraging occurs in large, open forest stands with space between the tree crowns to provide room for flight and hovering (Reynolds and Linkhart 1987). Territory size varies from 20 to 59 acres and is determined by age and patchiness of tree canopy. Flammulated owls are migratory in the northern part of their range. They arrive on their breeding territories in May and depart by mid-October, when they migrate south to central Mexico and Central America.

Preferred habitat is present in the southeastern portion of the analysis area. In addition, aspen stands to the west provide suitable habitat for this species. Two years of surveys were conducted for flammulated owls in suitable habitat in the analysis area (see section 2.4.1.4). Flammulated owls were heard and/or seen at 26 calling stations. No nests or juveniles were found, but based on the number of responses and small territory size, it is assumed that several pairs of flammulated owls occur in the analysis area.

3.1.2.10 Peregrine Falcon

Peregrine falcons are found in a wide variety of habitats in the Intermountain West. They prefer to nest on cliffs (generally at least 200 feet in height), usually in mountainous areas or in river canyons and gorges, although aeries are also known in metropolitan areas on structures such as towers and high-rise buildings (Bond et al. 1984). Peregrines prey almost exclusively on other species of birds, especially doves, pigeons, shorebirds, waterfowl, and passerines. They may forage up to 18 miles away from their aeries, although most hunting occurs within a 10-mile radius of the nest, and often over 80 percent of the foraging occurs within 1 mile of the aerie (Ehrlich et al. 1988). Peregrines overwinter in a wide range of habitats, but in the Intermountain West they appear to concentrate along large rivers and in wildlife refuges. Some birds may remain on their breeding territories throughout the year if there is an adequate food supply (Spahr et al. 1991). Aeries have been reported at elevations above 10,500 feet, although nesting above 8,000 feet is extremely rare (Bond et al. 1984).

Suitable nesting habitat is present in the analysis area, on the rock escarpments bordering parts of Muddy Creek and its tributaries. A pair of peregrine falcons was observed in 2002 near the rim of Muddy Creek Canyon in the eastern portion of the tract buffer. The pair was exhibiting territorial behavior thus it is presumed that a nest was nearby within the cliff faces. A peregrine falcon was observed circling above an inactive golden eagle nest during UDWR aerial surveys in 2003, less than one-half mile from the 2002 observation. No falcons were observed in 2001.

3.1.2.11 Three-Toed Woodpecker

Three-toed woodpeckers typically inhabit spruce/fir forests up to 9,000 feet, but where insect populations are high they may also occur in ponderosa or lodgepole pine forests. They are most apparent in years and locations where trees have high insect populations, and are attracted to areas with numerous dead trees from wildfires, insect epidemics, blow-down, or other die-off (Andrews and Righter 1992). The woodpeckers forage on wood-boring insects in dead trees, primarily spruce beetles (Peterson 1990). Soft snags are preferred for excavating nest cavities, although they occasionally excavate live trees. This species may make small movements off its breeding territory in the winter to find food but is generally a year-round resident.

There has been an ongoing outbreak of spruce beetle in the MLNF and subsequently, localized areas of spruce forest in the analysis area have been infected and contain suitable habitat for three-toed woodpeckers. Surveys for this woodpecker in the analysis area resulted in 16 individual responses at 13 separate locations (see section 2.4.1.5). Additionally, a female was observed incidentally in the area during a goshawk survey. All woodpecker observations were in the western portion of the tract buffer and associated with the aspen and aspen mixed conifer habitat type. It was assumed that three or more nesting pairs were present during the survey period.

3.1.2.12 Greater Sage-Grouse

The greater sage-grouse is an upland game bird that is entirely dependent upon sagebrush communities for all stages of its life cycle, with extensive areas of this habitat type required year-round. Sage-grouse have a high fidelity to their seasonal habitats (breeding, late brood-rearing, and wintering habitats), and females commonly return to the same areas to nest each year. Depending on geographic location, breeding activities occur from March through early summer. Most sage-grouse nests are located under sagebrush plants that provide overhead cover, with 15 to 30 percent canopy cover preferred. Late brood-rearing habitats, used from summer into fall, usually have less dense sagebrush canopy than nesting habitats and generally have a higher proportion of grasses and forbs in the understory. Riparian meadows, springs, and streams are also used during this time, especially in dry years, as these areas produce the forbs and insects necessary for juvenile birds. Because the diet of chicks consists of forbs and insects, diverse plant communities with abundant insect populations are especially important. During winter, sage-grouse feed almost exclusively on sagebrush leaves and buds, so exposure above the snow, rather than canopy cover, is critical. (BLM 2003.)

Sage-grouse were historically abundant in the analysis area, and one known lek site, the Wildcat Knoll strutting ground, is currently used. Forty-eight sage-grouse were transplanted to the southern portion of the analysis area by UDWR between 1987 and 1990. UDWR has been annually monitoring the Wildcat Knolls strutting ground since 1991, and grouse from the reintroduction effort use this site. The site has received use by 3 to 20 cocks on a given year, with the lowest numbers observed in 2003. Grouse sign was observed in additional portions of the analysis area that potentially provide suitable habitat for lek sites. These sites were surveyed during the breeding season, and although a few cocks and hens were observed between the Head of Box Canyon and East Fork Box Canyon, the birds were not engaged in any lekking displays. Abundant grouse sign was observed in that area and several adults were observed near the headwaters of Box Canyon later in the year. It is assumed the canyon edges are used for

roosting. Brood-rearing habitat is also present. In addition, several grouse were observed drinking from cattle troughs in this area. Grouse were also observed using Box Pond as a watering site.

Winter and brood-rearing habitat was mapped by the UDWR and overlaps approximately 39 percent of the analysis area. These areas coincide with locations where grouse and grouse sign were observed.

TEPS Mammals

3.1.2.13 Canada Lynx

Mature to late-successional spruce-fir forests provide suitable foraging habitat for Canada lynx in the southern portion of its range. These forests can support snowshoe hares, the primary prey species for lynx, as well as red squirrels, an important alternative prey species. Early successional stands with high densities of shrubs and seedlings are optimal for snowshoe hares, and subsequently important for lynx. Conifer-aspen forests, particularly those with dense regeneration or an extensive shrub and woody debris understory component, may also be important for prey species. The Canada lynx breeds from late winter to early spring, with denning beginning in late spring. Mature forest stands are used for denning, cover for kittens, and travel corridors. Denning habitat includes dense downed trees and root wads, or dense live vegetation (Koehler 1990, Mowat et al. 2000). For denning habitat to be functional, it must be in or adjacent to large areas of quality foraging habitat (Ruediger et al. 2000).

Reports of lynx in Utah indicate no sightings between 1961 and 1993 on the MLNF (Ruggiero et al. 1994). Recent records of lynx in Utah include a 2002 record from the Manti-La Sal National Forest (Forest Service 2002b). This record was from a hair sample collected in an isolated location near Joe's Valley. No additional lynx have been recorded in this area and it is likely that this individual was dispersing through the forest as opposed to having an established home range. Lynx are considered extremely rare in Utah and, of the few historic sightings that have occurred, the majority have been in the Uinta Mountains. Suitable habitat for lynx is limited due to the isolated nature and small size of forest patches on this part of the Forest, but could potentially occur in the western portion of the analysis area.

3.1.2.14 Spotted Bat

Spotted bats are found in relatively remote, undisturbed areas in a variety of habitats, including open ponderosa pine, desert shrub, pinyon-juniper, and open pasture and hay fields, and have been recorded at elevations as high as 9,500 feet. They roost alone in rock crevices on steep cliff faces and have been found hibernating in caves. Spotted bats are territorial and use echolocation to avoid each other while foraging. Their diet consists primarily of moths caught in flight after dark in open pine stands and over marshes (Wai-Ping and Fenton 1989). Information on seasonal movements is scarce, though spotted bats are thought to migrate south to hibernate.

Suitable roosting habitat for spotted bats is abundant in vertical cracks of the sandstone cliff faces of steep canyons in the analysis area. Riparian habitat and forest edges in this area also provide potential foraging opportunities. Several spotted bats were identified in the analysis area by audible vocalizations.

Auditory bat observations were associated with the rocky cliff habitat and ponderosa pine along the East Fork and main stem of Box Canyon and along Greens and Cowboy Canyons. Bats were also observed foraging in the limber pine habitat near Julius Flats Reservoir and above the North Fork of Muddy Creek, and in the limber pine/Douglas fir habitat along the jeep trail running west and south of Cowboy Creek. Spotted bats have also been identified in Muddy Creek Canyon and the lower end of Box Canyon with ANABAT detectors (Perkins and Peterson 1997).

3.1.2.15 Western Big-Eared Bat

Townsend's big-eared bats use juniper/pine forests, shrub/steppe grasslands, deciduous forests, and mixed coniferous forests from sea level to 10,000 feet. During winter they roost singly or in small clusters in caves, mine shafts, rocky outcrops, or occasionally in old buildings (Oliver 2000). They remain at these sites, called hibernacula, from October to February. They do not migrate, but will move to different roost locations within hibernacula during winter. In summer, females roost with their young in nursery roosts. Males and non-breeding females roost alone. Big-eared bats are sensitive to human disturbance and will abandon roost sites if disturbed. Townsend's big-eared bats are nocturnal insectivores and prey primarily on moths along forest edges.

No substantial caves have been observed in the analysis area and no other structures are considered potentially suitable for western big-eared bat hibernacula.

TEPS Amphibians

3.1.2.16 Spotted Frog

Columbia spotted frogs are found in areas where permanent, quiet water is present, such as marshy edges of ponds or lakes, algae-grown overflow pools of streams, emergent wetlands, and near springs. Emergent and submergent vegetation are considered important habitat features. Following the spring breeding season they may move considerable distances from water, often frequenting mixed conifer and subalpine forests, grasslands, and brushlands of sage and rabbitbrush if puddles, seeps or other water is available. However, in the Wasatch front, research indicates that spotted frogs travel only short distances between breeding and post-breeding habitats, with dispersal corridors typically being limited to aquatic or semi-aquatic habitats such as streams, intermittent drainages, and seeps, and that many breeding sites serve as year-round habitat (FWS 2002). Adult spotted frogs feed on invertebrates, generally within 0.5 meters of shore on dry days. During and immediately after rains, they may move away from permanent water to feed in wet vegetation or ephemeral puddles (Licht 1986). Spotted frogs hibernate during winter and emerge to breed when open water becomes available, generally during spring thaw.

Utah is in the southern portion of the spotted frog range in which two populations, the Wasatch Front and West Desert populations, are known to exist. These are believed to be relict populations, occurring in small patches of suitable habitat remaining since the last ice age (FWS 2002). Spotted frogs have not been located on the MLNF, although individuals were observed near Fairview, just west of the Forest. These frogs were likely from the southernmost range of the Wasatch Front population. Spotted frogs were not observed during survey efforts in the analysis area. Although potentially suitable habitat is present in localized areas, the analysis area is outside of the known and predicted range of this species, and it is unlikely that spotted frogs are present.

3.1.3 Management Indicator Species

Golden eagles, mule deer, elk, and aquatic macroinvertebrates are discussed in the following section. Although goshawks are also a MIS, they are addressed above as a Forest Service sensitive species in section 3.1.2. Since blue grouse are no longer a MIS for the MLNF, they are discussed briefly in section 3.1.4.

3.1.3.1 Golden Eagle

Golden eagles are typically found in open country, including shrublands, grasslands, canyons, and desert plains, as well as open coniferous forests in mountainous regions. Elevated nest sites, typically on cliff faces near hunting grounds, are the preferred breeding habitat. In the absence of suitable cliffs and rock outcrops, they have been known to nest in trees. Golden eagles feed mainly on small mammals, especially rabbits, marmots, and ground squirrels, but also eat insects, snakes, birds, juvenile ungulates, and carrion. Golden eagles typically mate for life. The breeding season generally begins in mid-January and continues through mid-September, though it can vary according to geographic region.

Suitable nesting habitat is present in the northern and northeastern portions of the analysis area on rock escarpments along Muddy Creek Canyon and some of its tributaries. Aerial surveys for eagles have been conducted by UDWR since 1998. Twelve golden eagle nest sites are known in the analysis area, of which one has been active and seven more tended at least once over the last six years.

3.1.3.2 Mule Deer

Mule deer are found in coniferous forests, shrub steppe, chaparral and grasslands with shrubs, from dry, open country to dense forests. They are often associated with early successional vegetation. They are known to utilize rocky brushy areas, open meadows, open pine forests, and burns. Mid to late seral range vegetation is used for forage. They browse on various grasses and forbs during the spring, summer, and fall, and on woody plants during the winter. Thermal and hiding cover is required year-round by elk. Thermal cover for deer generally consists of small conifers and shrubs on winter range, and deciduous or evergreen saplings or shrubs with high canopy closure on summer and spring-fall ranges. Water is also an important habitat component, especially on summer range. Fawning habitat for deer consists of low shrubs or small trees (2 to 6 feet tall) under a partially closed forest canopy. The fawning areas tend to be relatively small, close to water (less than 600 feet), and on mild slopes where succulent vegetation is abundant (Thomas et al. 1979).

The Muddy analysis area contains winter and summer range for mule deer. The value of this range is classified as high summer (~14,855 acres) and high winter (~18,860 acres). The range combined covers over 90 percent of the analysis area.

Annual winter counts of deer are not conducted by the UDWR. However, population data is modeled using harvest data for the entire Manti Deer Management Unit, which contains the analysis area. For the 2000 to 2001 hunting season, the post-hunting and spring populations in the Unit were predicted to be 3,603 and 5,436 individuals, respectively. The fawns/100 does were estimated at 69, fawns/100 adults at 60, and bucks/100 does at 14 for the post-hunting season. The deer population is far below the UDWR objectives for this unit and has been so for several years. The decline in deer populations is attributed to the drought and other natural environmental factors (UDWR 2001b).

3.1.3.3 Elk

Elk inhabit coniferous and mixed-coniferous forests as well as woodlands, chaparral, and grasslands in the Rocky Mountains. Mid to late seral range vegetation is used for forage. They rely on grasses for most of the year but also consume forbs in the summer and may browse on woody plants where grasses are unavailable, especially during winter months. Water is an important habitat component, particularly on summer range. During the summer elk spend the majority of their time in alpine and subalpine mountain meadows or in stream habitats. Thermal and security cover is required year-round by elk and generally consists of mature forest with large amounts of edge along grasslands or meadows. During the winter, elk movements are restricted by forage availability and snow conditions, and heat and energy are conserved in order to survive. Elk migrate altitudinally to lower elevations where snow depth is shallow and typically inhabit coniferous forests interspersed with riparian areas and south-facing slopes with sagebrush and shrubs, as well as aspen forests. Calving habitat for elk contains forage areas, hiding cover, and thermal cover within forest stands. Components of this habitat include shrubs or downed logs, gentle slopes, succulent forage, and a source of nearby water (less than 1,000 feet).

The Muddy analysis area contains winter and summer range for elk. The value of this range is classified as critical summer (~16,505 acres) and critical winter (~17215 acres). The range combined covers over 90 percent of the analysis area.

The winter aerial census for elk conducted in 2001 shows that the elk populations in the South Manti Sub-Unit of the Manti Management Unit to be slightly below UDWR objectives. However, elk numbers were purposely decreased, through increasing the number of cow tags issued, to compensate for the affects of the drought. A total number of 1,120 elk were counted on the South Muddy survey area and 449 in the North Muddy/Ferron survey area during this survey effort. Of these, 51 and 63 were bulls, respectively, and the remaining elk were antlerless. The calves/100 cows ratio was estimated at 29. A more recent winter census was conducted in January 2004 by UDWR, but summarized data is currently not available (UDWR 2001a).

3.1.3.4 Aquatic Macroinvertebrates


Macroinvertebrates are ecological indicators of the condition of aquatic habitats and the ability of these habitats to support fisheries. These species are affected by several environmental factors including water temperature, water quality, flow, and substrate type. Changes in aquatic habitats caused by management activities can lead to changes in the species composition and abundance of macroinvertebrates.

In general, higher abundance and diversity of macroinvertebrates are associated with cool water temperatures, substantial perennial flows, and diverse streambed substrate. Lower abundance and diversity are associated with ephemeral streams. In general, ephemeral streams present high water temperature, low flow, and streambeds with large amounts of fine sediment. Therefore, macroinvertebrate diversity and abundance within the analysis area is expected to be higher on perennial streams than in the ephemeral springs and drainages.

Aquatic invertebrate sampling was conducted in Greens Canyon (Sites 1 and 1A), Cowboy Creek (Sites 2, 3, and 3A), Greens Hollow (Site 4), and an unnamed stream near White Mountain Cabin (Control site). A total of 126 invertebrate taxa were identified in the 49 samples collected over the 3-year sampling period (2001 to 2003). Taxa from five functional feeding groups (shredders, scrapers, collector filterers, collector gatherers, and predators) were collected, with collector gatherers representing the highest number of taxa and individuals collected for each year of sampling. The five dominant taxa collected consisted of Turbellaria, Orthocladinae, Baetis, Pericoma, and Chironominae, and the dominant families included Chironomidae, Baetidae, Psychodidae, and Nematodeidae.

Mean macroinvertebrate abundance, richness, diversity, and biotic condition are depicted in Figures 1 through 4. The observed macroinvertebrate abundance was similar during spring and fall across the main sampling sites. Considerably higher invertebrate abundance was observed at the White Mountain Cabin control site. The lowest

Taxa Richness and Shannon Diversity Index (H) estimates were observed at the Greens Canyon site. Higher H values were estimated at the remaining sites. These estimates, including the control, were similar across sites and seasons. Estimates of the Biotic Condition Index (BCI) were near or above the level required by the Forest management plan ($BCI \geq 75$) with the exception of the site located at Upper Cowboy Creek (Site 3; average $BCI = 49$). In general, these estimates indicated that the streams surveyed were in fair to good condition. In addition, the Hilsenoff Biotic Index (HBI) indicated that the surveyed streams are slightly (HBI: 2-4) to moderately enriched (HBI: 4-7). The highest HBI estimates were observed at Upper Cowboy Creek (HBI=5.3) and Greens Hollow (HBI=5.7). A summary of results of the macroinvertebrate survey results is presented in Table 8 in section 2.4.2.5.



Figures 1-4. Summary of quantitative macroinvertebrate surveys conducted in the Muddy analysis area. Mean abundance (#/m²; Top-left), taxa richness (Top-right), Shannon diversity (Bottom-left), and biotic condition index (Bottom-right) for Greens Canyon (Site 1A), Lower Cowboy Creek (Site 2), Upper Cowboy Creek (Site 3), Middle Cowboy Creek (Site 3A), Greens Hollow (Site 4), and White Mountain Cabin (Control Site). Error bars represent one Standard Deviation.

3.1.4 Species of High Federal Interest

Species of high federal interest, as defined by the FWS, include several migratory birds. No other categories of wildlife were identified by the FWS.

3.1.4.1 Migratory Birds

Twenty-two migratory birds are on the list of species of high federal interest (see Table 9, section 2.4.3.1), of which seven are known to occur and ten could potentially occur in the analysis area. Species observed include the bald eagle, golden eagle, peregrine falcon, prairie falcon, Cooper's hawk, flammulated owl, and Williamson's sapsucker. Species that could potentially occur include the osprey, ferruginous hawk, merlin, western bluebird, Scott's oriole, burrowing owl, Mexican spotted owl, band-tailed pigeon, great blue heron, and black swift. The bald eagle, spotted owl, flammulated owl, peregrine falcon, and golden eagle are discussed in sections 3.1.2.6, 3.1.2.7, 3.1.2.9, 3.1.2.10, and 3.1.3.1, respectively.

Of the species observed or potentially occurring in the analysis area, golden eagles, peregrine falcons, prairie falcons, and Mexican spotted owls use cliffs for nesting. Black swifts also may use cliff habitats, such as a ledge or a crevice, but nests are usually near or behind waterfalls. Flammulated owls use abandoned woodpecker holes in snags, and merlins typically use abandoned hawk nests in trees, but may also use cliffs. Cooper's hawks, ferruginous hawks, band-tailed pigeons, Williamson's sapsuckers, and western bluebirds typically nest in trees. Burrowing owls nest in mammal burrows in grasslands and Scott's orioles use suspended nests attached to shrubs or small trees. Bald eagles are not known or expected to nest in the analysis area, although perennial streams may be used for foraging.

3.1.5 Other Species

3.1.5.1 Fishes

Rainbow and brook trout were also observed during fish surveys conducted by UDWR on perennial streams in the analysis area. Rainbow trout were observed in Muddy Creek and South Fork of Muddy Creek, while Brook trout were only observed in Muddy Creek. The streams where cutthroat trout were also observed during surveys and incidentally (i.e., Muddy Creek and its South and North forks) present moderate to high quality trout habitat. As described in section 2.4.1.2, no fish were observed in the North Fork of Quitchupah Creek. Erosion, siltation, and low water flows have led to the poor trout habitat in this stream. Speckled dace (*Rhinichthys oscolus*) have been observed on mainstem sections of Quitchupah Creek located outside of the analysis area.

3.1.5.1.1 Brook trout

Brook trout (*Salvelinus fontinalis*) is a sport fish native to the eastern United States and eastern Canada. This species has become established in many of Utah's cold higher-elevation lakes and streams. The diet of brook trout is based primarily on invertebrates, including insects and zooplankton; large individuals occasionally feed on small fishes. Spawning occurs in the fall over gravel substrate in lakes and streams. Hatching and emergence occurs approximately after two months. The successful reproduction of brook trout has led to overcrowding, and consequently, to a large number of stunted (small) individuals in streams in Utah. The overcrowding problem can be exacerbated by low fishing pressure in the brook trout's high elevation habitat. This species poses a threat to native cutthroat trout populations (Sigler and Sigler 1996).

3.1.5.1.2 Rainbow trout

Rainbow trout (*Oncorhynchus mykiss*) is a popular sport fish in Utah. This species is native to western North America but is not native to Utah. The popularity of this species in addition to their unsuccessful reproduction in the wild has led to the introduction of millions of rainbow trout to cold and cool waters throughout the state by the UDWR. The UDWR has also stocked an albino form of the rainbow trout into many Utah waters. Their diet is based primarily on invertebrates, including insects, worms, zooplankton, and insect larvae. Rainbow trout that reach larger sizes can switch to a piscivorous diet. The species spawns in streams over gravel substrate during the spring, and the eggs hatch in about one month. Fry emerge approximately two to three weeks after hatching. The presence of rainbow trout in streams that provide habitat to cutthroat poses a major threat to the native species. Similarities in the habitat and timing of spawning often lead to the production of rainbow - cutthroat hybrids, and thus, to the loss of genetic purity through hybridization (Sigler and Sigler 1996).

3.1.5.2 Blue Grouse

Blue grouse breed in open coniferous and aspen forests with a shrub understory or adjacent to shrublands. They spend the winter at higher elevations than summer habitat, primarily in Douglas-fir and lodgepole pine forests of various age classes and tree densities (Andrews and Righter 1992). They have also been known to winter in spruce forests in southwest Colorado. Grouse roost in large conifers with dense foliage. Grouse feed primarily on needles and buds of conifers in the winter (Douglas-fir often important) and berries, insects, flowers, and leaves in the summer.

Suitable habitat for blue grouse is present, but limited, in the analysis area. Grouse were observed at three separate locations in this area, and all observations were associated with or near small aspens and mountain shrubs. Potential brood-rearing habitat could occur within the forested portions of the analysis area. However, this habitat is not typical of that used by grouse, and the scant shrub component in the spruce-fir stands likely renders this habitat unsuitable. The forested portions of the project area are likely more suitable as summer habitat and potentially winter habitat.

3.1.5.3 Amphibians

Amphibian habitat is limited in the analysis area, consisting of wetlands, ponds (natural and stock), edges of lakes and reservoirs, springs and seeps, and pooled habitat adjacent to streams. Potentially suitable amphibian habitat surveyed during the analysis period resulted in observations of boreal toads, chorus frogs, tiger salamanders, and possibly Great Basin spadefoot toads. Chorus frogs were the most abundant species observed (see section 2.4.4.2). All life stages of chorus frogs and tiger salamanders (eggs, tadpoles, and adults) were observed in ponds. Chorus frogs were also heard at Julius Flat Reservoir. All boreal toad observations were of larvae in ponds. Great Basin spadefoot toads were potentially heard at a cattle pond and in a stream channel at the bottom of Box Canyon.

Other amphibian species that could potentially occur in the analysis area include the great plains toad, woodhouse's toad, and northern leopard frog. Spotted frogs are not expected to occur as far south as the analysis area.

3.1.5.4 Reptiles

Suitable habitat for several reptile species is present in the analysis area. Lizard, whiptail, and skink species primarily occur in desert and semi-desert areas with sandy or rocky soil and sparse vegetation, such as pinyon-juniper and sagebrush, but also occur in grasslands and the lower edge of the spruce-fir zone. Habitat for snake species is also variable, ranging from lowlands to high mountains, with some species having an affinity for riparian habitats, and others for more arid environments.

Five reptile species were incidentally observed in the analysis area: the eastern fence lizard, tree lizard, sagebrush lizard, short-horned lizard, and western terrestrial garter snake. Reptile species not observed but likely present include the common side-blotched lizard, gopher snake, night snake, striped whipsnake, and western rattlesnake. Other species possibly present include the Great Basin collared lizard, long-nosed leopard lizard, desert spiny lizard, Western whiptail, Western skink, Eastern racer, milk snake, Southwestern blackheaded snake, and ground snake.

3.1.5.5 Small Mammals

Seventy species of small mammals could potentially occur in the study area, including 5 shrews, 15 bats, 8 small carnivores, 36 rodents, and 6 lagomorphs. Of these, 16 were observed by Cirrus personnel (1 bat, 1 carnivore, 10 rodents, and 4 lagomorphs). All habitats in the analysis area are potentially used by at least some small mammals, with riparian habitats being used by the largest number of species.

3.1.5.6 Non-Game Birds

A total of 203 species of non-game birds could potentially occur in the study area. Of these, 84 were observed by Cirrus personnel. All habitats in the analysis area are potentially used by at least some non-game birds, with riparian habitats being used by the largest number of species. Non-game species that potentially use cliffs in the analysis area for nesting include, but are not limited to, the golden eagle, prairie falcon, peregrine falcon, red-tailed hawk, Mexican spotted owl, raven, white-throated and black swifts, cliff swallows, canyon wren.

3.2 Detailed Technical Assessment/Description of the Potential Effects

This section presents an assessment and description of potential impacts to aquatic and terrestrial wildlife resources. The section is organized by issue statement, with Alternatives 1 through 3 addressed under each issue. The evaluation criteria defined in the RFP for this project was used as a guide for determining potential impacts. The available data was used to predict reasonable foreseeable mining scenarios and is used in the analysis of the four wildlife issues.

3.2.1 Wildlife Issue 1

Any changes in water flow and quality in perennial drainages and reservoirs or to riparian vegetation/wetlands could affect habitat for terrestrial and aquatic species.

3.2.1.1 Alternative 1: No-Action

No leasing or mining would occur under this alternative. No changes in water flow or quality of perennial streams and drainages or to riparian and wetland ecosystems would be expected. Therefore, habitat for terrestrial and aquatic species would not be affected.

3.2.1.2 Alternative 2: Standard Lease Terms and Conditions Only

Under this alternative, the tract would be leased and mined with BLM standard lease terms and conditions. This alternative would allow longwall mining (full extraction) throughout the tract, which could result in subsidence of perennial drainages, escarpments, and surface facilities. The duration of mining for complete coal recovery would be approximately 20 years. Localized impacts associated with mining, such as subsidence and subsidence-induced tension cracks as discussed below, are estimated to occur over one to two years, with the majority of the subsidence occurring in the first three weeks after coal extraction.

Mining activities would result in subsidence-induced ground movements and other changes in geology and topography. These changes include variations in stream gradient, tension cracks, and rock failures. Subsurface disturbances could cause temporary cracks to open up in streambeds, which could divert flow underground. Temporary disruptions of ground and surface water flows could reduce water availability for fish and aquatic invertebrates. In addition, subsidence could disturb escarpments in localized areas (MTI 2004), which could lead to major disruptions of the natural sediment delivery process to streams (Nelson et al. 2003). These changes could influence the abundance and community structure of aquatic species.

As stated in the Surface and Ground Water Technical Report prepared for the Muddy Creek Tract, cracks resulting from subsidence could enhance the rate of vertical flow from ground waters, and thus lead to reduced flow at springs originating above the mined area. The likelihood of springs drying up completely is low, and due to differences in the overburden thickness, the risk of permanent impacts from vertical fractures is expected to be low, with the exception of the Box Canyon springs. If flow is permanently affected at these springs, the water diverted underground would be expected to discharge at a different location further down slope.

Perennial streams that would be undermined under Alternative 2 and may be affected by subsidence include Muddy Creek and tributaries of Cowboy Creek. As discussed in the Surface and Ground Water Technical Report, subsidence of streams could intercept flowing water and divert it into underground workings or enhance subsurface flow in the shallow bedrock underlying the stream valley. Stream segments occurring within the Castlegate Sandstone outcrop along Greens Hollow and Cowboy Creek and the segments with low overburden cover along Muddy Creek, Horse Creek, and Greens Canyon present the highest risk of subsidence. The risk of water diversion into underground workings is greatest for Muddy Creek, and would likely result in a loss of stream flow and alluvial groundwater. Subsidence fracturing would also pose a significant risk of enhanced water losses from Greens Canyon, and would likely reduce the length of perennial flow of the stream segment. Subsidence could lead to the temporary reduction in intermittent flows of Greens Hollow and Cowboy Creek and an increase in subsurface flow in the fractured bedrock.

Mining could also impact ponds (natural basins and stock ponds) and wetlands. Although these habitats make up a small portion of the analysis area and impact zone, they are important for a large number of wildlife species. Subsidence-induced tension cracks could divert surface water to underwater networks on a temporary basis (less than 2 years), thus reducing the availability of water for aquatic and terrestrial wildlife. These cracks could potentially cause ponds to dry up. Over time, as the tension is released, the cracks would close, and organic debris would fill the remaining gaps. Stress on riparian vegetation, loss of wetlands, and/or changes in species composition could result from the temporary changes in water availability, thus indirectly impacting wildlife. No reservoirs are inside of the zone of potential impact. Therefore, no impacts would result to this form of aquatic habitat.

Mining could impact escarpments in areas located near Box, Greens, and Muddy canyon. Potential effects in these areas include the formation of cracks and spalling of escarpments (MTI 2004). In addition, localized areas could also be affected by water withdrawals for exploration drilling.

Impacts to water quality from subsidence, as discussed in the Surface and Ground Water Technical Report, are expected to be minor and imperceptible. Therefore, the remainder of this section focuses only on potential impacts to wildlife from changes in water flow or to riparian vegetation and wetlands. Potential impacts to specific species or groups are described below.

3.2.1.2.1 Fisheries

The cutthroat trout, thought to belong to the native Colorado River subspecies, is the only fish species of concern within the analysis area. This native species occurs in Muddy Creek and thus could be impacted by flow reductions caused by diversion of water to underground workings. The risk of flow diversions is greatest in the area of low overburden cover along Muddy Creek. The magnitude of potential impact to cutthroat trout depends on the volume of surface water that could be lost to subsurface flows. Fish migrating upstream to spawn require suitable water velocities and depths to succeed. Thompson (*in* Bjorn and Reiser 1991) quantified the minimum water depth that would allow trout migration. According to his estimates, migration would succeed in depths of 0.12 to 0.18 meters. Based on recent surveys, the average stream depth in Muddy Creek is 0.57 meters. A reduction in water depth of 68 to 78 percent in this stream could influence the spawning success of the native cutthroat trout species. In addition, flow regulates the amount of spawning area available by regulating the area covered by water and the water velocity

over gravel beds (Bjorn and Reiser 1991). Hunter (*in* Bjorn and Reiser 1991) determined that cutthroat trout prefers water depths greater than 6 centimeters and velocities between 13 and 72 centimeters per second. Reductions in depth and velocity below these levels could also impact the spawning success of cutthroat trout. Further, fish are not uniformly distributed at all depths in a stream. For example, Pratt (*in* Bjorn And Reiser 1991) determined that cutthroat trout less than 100 mm used lower depths than fish larger than 100 mm (32 and 62 cm respectively). Consequently, potential reductions in flow and water depth could lead to more negative impacts on small cutthroat trout than on larger fish. These effects are expected to be temporary, as seasonal flows are likely to transport substrates downstream and thus fill in cracks within a short time period. According to the Geology Technical Report prepared for the Muddy Creek Tract, the natural recovery of tension cracks in a streambed could range from a few weeks to one or two years. Mitigation is recommended in section 3.3 to minimize potential impacts to fisheries habitat.

Activities that deplete water from the Colorado River have been identified by the FWS as having adverse cumulative effects to the endangered bonytail, Colorado pikeminnow, humpback chub, and razorback sucker. However, transbasin water diversions that could affect these species are not expected. Changes to water flow in the analysis area would not result in water depletions from the Colorado River Basin. A reduction in flow along Muddy Creek is expected, but because of the existence of water rights along this stream, the coal-mining permit would require mitigation by either sealing the subsidence fractures or providing alternative water supplies. With implementation of these and other recommended mitigation measures, formal consultation with FWS for the four endangered fish species would not be warranted.

3.2.1.2.2 Macroinvertebrates

Aquatic macroinvertebrates depend on the flow of seasonal and perennial waters. Higher abundance and diversity of macroinvertebrates are usually associated with cool water temperatures, substantial perennial flows, and diverse streambed substrate. Lower abundance and diversity are associated with intermittent streams with high water temperature, low flow, and streambeds with large amounts of fine sediment. Potential flow reductions in localized areas in Muddy Creek could modify the species composition and abundance at a stream reach scale. As discussed in the Surface and Ground Water Technical Report prepared for the Muddy Creek Tract, the risk of subsidence fractures intercepting stream water and diverting it into underground mine workings is greatest in areas of low overburden cover along Muddy Creek.

Potential damage from tensile strains that could cause surface cracks and spall of escarpments is also expected in areas located near Greens, Box, and Muddy canyons (MTI 2004). However, as discussed in section 3.2.1.2.1, tension cracks in the streambed would recover naturally, in as little as a few weeks, or up to two years. The Biotic Condition Index (BCI >80) at Greens Canyon indicated that this stream is in good condition. Under this alternative, potential flow losses or reductions in this stream could affect invertebrate habitat, abundance, and diversity. A reduction of 20 percent or more in the BCI would require further evaluation and possibly a change in management direction as required by the Standards and Guidelines defined in the Forest Management Plan for the MLNF (Forest Service 1986).

Minimal impacts on aquatic invertebrates within the tributaries of Cowboy Creek are expected, as the effects from subsidence on stream flow are expected to be minimal (MTI 2004). However, temporary reductions to intermittent flows and perennial reach lengths could occur in Greens Hollow and Cowboy Creek as a result of surface tensile fractures in the nearby Castlegate Sandstone, and could cause associated reductions to invertebrate habitat, abundance, and diversity. Impacts to Greens Hollow and to the intermittent portions of Cowboy Creek could potentially pose a greater risk to associated invertebrate communities than to those in Greens Canyon, given the low Biotic Condition Index (BCI <65; poor quality) of these stream reaches and the presence of grazing in these areas. However, as discussed above, tension cracks would recover naturally, thus the impacts to stream flow and associated invertebrate communities would be temporary.

3.2.1.2.3 Birds

Potential stress on the riparian vegetation from diversion of surface water could reduce the function and value of riparian habitat to many bird species. However, since the majority of impacts to surface water would be short-term (less than 2 years), associated impacts on vegetation and wildlife are expected to be temporary. An exception could be to Muddy Creek, where impacts to water flow could be longer term, potentially leading to a loss of riparian habitat. Riparian habitats provide important brood-rearing habitat for sage-grouse, as the young rely on insects and

succulent forbs. A reduction in riparian habitat or surface water availability could therefore impact brood-rearing habitat for sage-grouse. Riparian habitat represents a critical component of several non-game bird species, including many warblers. Stress on riparian vegetation could reduce the availability or quality of nesting and/or foraging habitat for these species. Several bird species also rely on pooled or flowing water as a water source. Changes in the availability of free water may result in modification in behavior of birds as they search for alternative water sources.

3.2.1.2.4 Mammals

Loss of surface water could impact deer and elk, but this impact would be limited to seasons when snow and succulent forage were not available. The impact would be minor, causing temporary modification in behavior and daily movements as they search for alternative water sources. The potential stress on riparian vegetation and temporary reductions of surface waters could impact the quality of habitat for several small mammals, particularly those that rely on riparian habitats for foraging, such as some shrew and bat species. However, these impacts would likely be temporary.

3.2.1.2.5 Amphibians

Changes in water flow could reduce the amount of habitat available to amphibians in the analysis area. Because amphibians are dependent on water for at least part of their life cycle, a decline in number of individuals would be expected if a substantial loss of flow resulted from mining-induced subsidence. These impacts would be temporary (less than 2 years), lasting until the tension cracks sealed and surface waters were restored.

The majority of amphibians observed in the analysis area were in ponds. The majority of these ponds, including those where boreal toads were observed, was near the perimeter of the buffer boundary, and would therefore not be impacted. However, a few ponds are within or near the area that would be undermined. If cracks occurred in these ponds, surface flow would be temporarily reduced or eliminated, thus eliminating potential amphibian breeding habitat for the duration of the affect. However, as discussed in the Surface and Ground Water Technical Report prepared for the Muddy Creek Tract, all of the ponds within the study area are located at least 1,000 feet above the mine coal. Therefore, the risk of tensile cracks from mine subsidence causing enhanced water loss from ponds would be relatively low. Furthermore, since the ponds are located within formations that contain abundant shales and clays, any surface tensile cracks that may occur as a result of mine subsidence would likely seal.

Wetlands provide an abundant source of insects for amphibians to feed on. Therefore, potential impacts to springs and associated wetlands from mining-induced subsidence, as discussed in the Vegetation Technical Report prepared for the Muddy Creek Tract, would affect amphibian habitat. With the potential exception of wetlands associated with springs SP_08, SP_09, and SP_39, impacts would be temporary for the reasons discussed above for ponds (overburden depth and clay substrate).

3.2.1.2.6 Reptiles

Reptiles would be minimally impacted, if at all, by changes in water flow and quality, as very few reptile species rely on riparian habitats. Species that commonly use riparian areas, such as the western terrestrial garter snake, could potentially be impacted. However, this species is not restricted to riparian habitats, and would likely travel over terrestrial habitat until an alternative water source was encountered.

3.2.1.3 Alternative 3: Standard Lease Terms and Conditions and Special Stipulations

Under this alternative the tract would be leased with special coal lease stipulations for the MLNF in addition to the standard terms and conditions. These stipulations would eliminate or minimize subsidence and its potential effects on perennial drainages. Stipulations associated with aquatic resources include Forest Service Stipulations 3, 7, 9, and 17. Stipulation 3 requires that the lessee obtain baseline data to quantify the existing surface resources. Stipulation 7 requires that baseline data be used for future monitoring and evaluation of effects. Stipulation 9 requires that mining operations be conducted in a manner that would prevent surface subsidence, which could lead to escarpment failures and landslides as well as to damage or alterations of flow in perennial streams. Stipulation 17 requires that any ground or surface waters identified for protection that may be impacted by mining would have to be restored by the lessee in order to maintain riparian and fishery habitat, wildlife, and other uses.

3.2.1.3.1 Aquatic and Terrestrial Wildlife

Impacts to aquatic and terrestrial wildlife would be similar to those previously described for Alternative 2. However, the special stipulations for the protection of wildlife and perennial drainages described above would minimize the potential impacts to perennial streams, riparian vegetation, and wetland habitat under this alternative. The perennial streams in Muddy and Box canyons would be protected from mining under this alternative by shortening the length of and/or eliminating some of the longwall panels. Therefore, associated subsidence impacts

to aquatic and terrestrial wildlife would not result. Since full longwall extraction of coal could still occur beneath isolated perennial segments of Cowboy Creek and Greens Hollow, impacts to wildlife discussed under Alternative 2 could still result. Impacts to riparian vegetation would be less than under Alternative 2, since Muddy Creek would not be undermined. However, impacts to riparian vegetation along Greens Hollow and Cowboy Creek could still occur. Impacts to ponds and wetlands associated with subsidence-induced tension cracks could still occur. However, risks of impacts to wetlands in Box Canyon and the small riparian zones associated with Box Canyon and Greens Canyon streams would be reduced because mining would not be allowed under these resources. Water depletion from the Colorado River system would not be expected since waters lost would be replaced. Therefore, formal consultation with FWS for the four endangered fish species would not be warranted.

3.2.2 Wildlife Issue 2

Subsidence of perennial streams could cause changes in stream morphology and aquatic habitat.

3.2.2.1 Alternative 1: No-Action

No leasing or mining would occur under this alternative. Subsidence of perennial streams would not occur under this alternative. Therefore, stream morphology and aquatic habitat would not be altered.

3.2.2.2 Alternative 2: Standard Lease Terms and Conditions Only

Mining activities under Alternative 2 would result in subsidence-induced ground movements and other changes in geology and topography in aquatic and terrestrial environments. These changes include variations in stream gradient, tension cracks, and rock failures. Subsurface disturbances could cause temporary cracks to open up in streambeds, which could divert flow underground. Temporary disruptions of ground and surface water flows could reduce water availability for aquatic species. Potential effects associated with changes in water flow are discussed above in section 3.2.1.2. In addition to these effects, subsidence could disturb escarpments in localized areas (MTI 2004), which could lead to major disruptions of the natural sediment delivery process to streams (Nelson et al. 2003). This disturbance could also cause streambank erosion and instability in localized areas.

Perennial streams that would be undermined under Alternative 2 and thus may be affected by subsidence include Muddy Creek, perennial sections of Cowboy Creek, Greens Hollow, and Greens Canyon. The maximum expected change in stream gradient in Muddy Tract would be 3 percent, and the maximum expected subsidence of the streambed would be 7 feet at localized areas of Muddy Creek. In addition, cracks could be formed and escarpments could be disturbed in areas located near Box and Greens canyons (MTI 2004).

As discussed in the Surface and Ground Water Technical Report prepared for the Muddy Creek Tract, changes in surface elevation caused by subsidence would be expected to occur in areas of low overburden cover along Muddy Creek and Horse Creek. Localized changes in surface elevation would be likely to create ponding in areas where slope reductions occurred. Due to the nature of these streams, channel incision may occur in areas of increased slope, while sediment deposition and ponding would be expected downstream at the end of the subsidence zone. These changes in stream morphology could alter habitat for aquatic species. Changes in surface elevation could also occur along Greens Hollow and Cowboy Creek. However, since natural pools, steep segments, and large boulders occur along these channels, these changes may not be apparent and functional changes in channel morphology are not expected.

Potential impacts to specific species or groups are described below.

3.2.2.2.1 Fisheries

The cutthroat trout, thought to belong to the native Colorado River subspecies, is the only fish species of concern within the analysis area. This native species occurs in Muddy Creek and thus could be impacted by changes in stream morphology and aquatic habitat. The severity of these impacts depends on the magnitude of the disturbance of escarpments near streambeds, as well as to the potential changes in stream flow. A subsidence of 7 feet, the maximum expected at Muddy Creek, has the potential to affect fish movements above this stream reach, thus it could limit the access to spawning habitat in the upper sections of the stream, as well as in the South and North Forks of Muddy Creek. Further, if the degree of subsidence is such that flow would be interrupted, this obstruction could lead to the isolation and decline of the cutthroat trout populations in these areas. It would be unlikely that a gradient change of 3 percent could change the composition and ratios of habitat types (Schmidt 2004). Mitigation is recommended in section 3.3 to minimize potential impacts to fisheries habitat.

The natural input of sediment to streams is a normal component of salmonid habitat. However, increased sediment delivery to streams can cause major disruptions to the aquatic habitat. These disruptions can lead to the movement and redistribution of spawning gravels, additions of new sediments, changes in accessibility to fish of spawning habitats, changes in availability of food organisms, and changes in seasonal and diurnal water temperatures (Swanston 1991).

Additional inputs of sediment to streams, led by subsidence and the potential disturbance of escarpments near streambeds, could cause short-term and long-term changes to aquatic organisms and their habitat. Short-term impacts (days to months) could result in increases in availability, transport, and deposition of sediment. The accumulation of fine sediment on spawning gravels could reduce the availability of spawning habitat and reduce spawning/hatching success. Increasing the amounts of suspended and bedload sediments could reduce light penetration and thus photosynthesis and primary production, as well as reduce survival by delaying fish movements (migration), disrupting fish feeding and thus growth, interfering with respiration, and increasing gill irritation and the potential for infection. Conversely, long-term impacts (years to decades) include changes that may actually improve habitat quality and productivity by increasing the total area available for spawning and rearing habitat. The addition of boulders, rubble, and gravel to the stream could lead to increases in habitat diversity and thus to the available habitat for fish. Obstructions caused by boulders and bedrock outcrops could modify channel velocity and direction, thus leading to the creation of pools, gravel bars, and side-channel rearing areas (Swanston 1991).

3.2.2.2 Macroinvertebrates

Aquatic macroinvertebrates depend on the flow of seasonal and perennial waters. As discussed above, higher abundance and diversity of macroinvertebrates are usually associated with cool water temperatures, substantial perennial flows, and diverse streambed substrate. Lower abundance and diversity are associated with ephemeral streams with high water temperature, low flow, and streambeds with large amounts of fine sediment.

Under this alternative, potential subsidence-induced changes in sediment inputs to Muddy Creek and Horse Creek and alterations in channel morphology could modify the species composition and abundance at a stream-reach scale. Potential increases to sediment loading in Greens Canyon could also affect macroinvertebrate communities. The estimated Biotic Condition Index indicated that this stream is in good condition (BCI >80). A reduction of 20 percent or more in the BCI would require further evaluation and possibly a change in management direction as required by the Standards and Guidelines defined in the Forest Management Plan for the MLNF (Forest Service 1986). The effects from subsidence on stream morphology within Greens Hollow and Cowboy Creek are expected to be minimal (MTI 2004). However, localized impacts to aquatic invertebrates could occur in these drainages. Changes in sediment input as well as changes in the number or distribution of pools in localized areas could lead to shifts in the composition and distribution of aquatic invertebrate communities at a small scale (e.g. habitat units and reaches). However, changes to invertebrate communities at a larger scale (e.g. drainages) are not expected. Potential effects on invertebrate communities from changes in water flow are discussed in section 3.2.1.2.2.

Potential damage from spill of escarpments also exists in areas located near Greens, Box, and Muddy canyons (MTI 2004). In these areas, increased bedload sediment could eliminate habitat for aquatic invertebrates, reduce abundance of invertebrates, and ultimately lead to reductions in fish production (Bjorn and Reiser 1991). Similar to the potential effects on fisheries discussed above, any damages to stream habitat could pose short-term and long-term effects. While short-term impacts may include the reduction in abundance and biodiversity of macroinvertebrates, the addition of boulders and rubble to the stream could result in a more complex habitat and thus increase species diversity in the long-term.

3.2.2.3 Alternative 3: Standard Lease Terms and Conditions and Special Stipulations

Under this alternative, the tract would be leased with special coal lease stipulations in addition to the standard terms and conditions, as described above in section 3.2.1.3. These stipulations would eliminate or minimize subsidence and its potential effects on perennial drainages and associated aquatic habitat.

3.2.2.3.1 Fish and Macroinvertebrates

Impacts to fish and macroinvertebrates would be similar as those previously described for Alternative 2 with the following exceptions. Special Stipulation 9 would prevent subsidence of perennial streams or escarpments thus eliminating associated impacts. Special Stipulation 17 would require the replacement of any waters lost due to the mining operation. Therefore, water depletion from the Colorado River system would not be expected, and formal consultation with FWS for the four endangered fish species would not be warranted.

3.2.3 Wildlife Issue 3

Exploration drilling and construction of mine vent holes could temporarily disrupt use of summer habitat by terrestrial species.

3.2.3.1 Alternative 1: No-Action

No leasing or mining would occur under this alternative. Exploration drilling and construction of mine vent holes would not occur. Therefore, use of summer habitat by terrestrial species would not be disrupted.

3.2.3.2 Alternative 2: Standard Lease Terms and Conditions Only

Mining activities under Alternative 2 would include coal exploration drilling and the construction of four mine vent holes and associated structures. Exploration drilling would occur at approximately 26 locations and would include the construction of associated drill pads, a staging area, and several short access roads. The majority of the drilling would be road supported, but helicopter supported drilling is proposed at three sites in canyon bottoms. Exploration activities are estimated to occur over 5 years and take place over a 2 month per year time period during the late summer and fall.

Disturbance associated with exploration drilling and construction of mine vent holes includes noise from equipment use and road travel. In addition, vegetation would be removed from small, localized areas. These areas would be reclaimed, but would constitute a temporary loss in wildlife habitat, and likely a change in vegetation type. The conceptual location of drill pads and roads, as depicted in the Geology Technical Report prepared for the Muddy Creek Tract (Plate 2), and the conceptual location of mine vent holes, as depicted in the Conceptual Mine Plan for the Muddy Tract (MTI 2002), were used as the basis for this analysis.

Total temporary disturbance due to exploration drilling would be approximately 17 acres. The approximate acres of disturbance associated with the construction of new roads, drill pads, and staging areas by vegetation type are depicted in Table 11.

Vegetation Type¹	Acres
Aspen	4.8
Sagebrush	4.3
Mahogany/Mountain Brush	4.1
Mixed Conifer	0.4
Limber Pine	0.3
Ponderosa Pine	0.2
Grassland	2.3
Pinyon/Juniper	0.1
Willow Riparian	0.1
Unidentified for staging areas	0.5
Total	16.6

¹Vegetation types are defined in section 3.1.1.

Potential impacts associated with exploration drilling and construction of mine vent holes to wildlife are discussed below.

3.2.3.2.1 TEPS Fishes

Exploration drilling would use water supplied by a relay system of pumps, water lines, and tanks. The streams proposed for water use have not yet been determined, but they would likely occur in the analysis area in the Muddy and/or Quitcupah drainages. Since these streams eventually flow into the Colorado River via the San Rafael and Green rivers, use of water for drilling, if not replaced or otherwise mitigated, would result in minor depletions to the Colorado River system, and thus impacts to the four endangered fish species could result. Therefore, formal consultation with the FWS could be required under this alternative.

Two drill pads are proposed near Muddy Creek and the South Fork of Muddy Creek, and helicopter-supported drilling is proposed in the bottom of Muddy Canyon. If construction activities took place too close to the streams,

potential impacts to habitat for cutthroat trout and other fishes could result. Mitigation is recommended in section 3.3 to minimize potential impacts to fisheries habitat.

3.2.3.2.2 TEPS Birds

Roads and drill pads associated with exploratory drilling would be in the vicinity of at least two goshawk territories and numerous flammulated owl territories. Noise associated with construction of roads and pads and with drilling could disrupt roosting and foraging behavior of these birds at the end of the nesting period. The magnitude of behavior modification would vary depending on the distance of the disturbance from the birds and nest sites, and the intensity and duration of the disturbance. Responses could vary from temporary startle responses (flush) and short avoidance flights, to longer-term avoidance of territories, and potential abandonment for a given year. As the majority of the exploratory drilling activities are not proposed to occur until late summer, the majority of the breeding season for these birds would be complete, and the young would be near, at, or past the fledgling stage. Mitigation is recommended in section 3.3 to minimize potential impacts to these species.

Drilling activities would not directly impact any other TEPS birds species. However, approximately 17 acres of habitat would be removed. Less than 5 acres occurs in each habitat type, thus the impact to wildlife foraging and nesting habitat would be negligible. Furthermore, these areas would be reclaimed, although the species composition would likely change. There would be no impacts to any TEPS bird species associated with construction of mine vent holes.

3.2.3.2.3 TEPS Mammals

There would be no direct impacts to TEPS mammal species from exploration drilling or construction of mine vent holes. There would be a minor reduction in habitat for moths, the primary prey species for spotted and Townsend's big-eared bats, but this impact would be negligible. Less than 6 acres of Canada lynx habitat would be removed. This impact would also be negligible. If a lynx were to use the tract as a travel corridor, it would be temporarily disturbed by noise associated with drilling, construction, and road use, but this species is not expected to occur in the analysis area.

3.2.3.2.4 MIS

One of the drill pad locations would be less than 0.25 miles from a golden eagle nest. The associated eagle pair could potentially be disturbed from the noise and human presence near this site, especially if the drill pad and road were visible from the nest site. Of the four proposed mine vent holes, three are in the vicinity of golden eagle nests. Noise associated with the construction of vent structures could temporarily disturb these pairs. Disturbance to eagles would likely be minor, if construction, road use, and drilling occurred late in or outside of the eagle breeding season. Mitigation is recommended in section 3.3 to minimize potential impacts to golden eagles.

The areas proposed for exploration drilling are associated with summer and winter range for mule deer and elk, primarily winter range. Deer and elk using these areas during the period of drilling activity could be temporarily disturbed. It is likely that they would avoid these areas at this time. Increased use of roads associated with exploratory drilling would also result during later summer and early fall, potentially resulting in vehicle-related mortality or habitat avoidance. Impacts of habitat removal and to available forage would be negligible since less than 17 acres of total habitat would be removed (and areas eventually reclaimed). Removal of habitat suitable for deer fawning and elk calving could occur. However, because so little of this habitat would be removed (less than 10 acres), these impacts are expected to be minor. Noise associated with the construction of vent structures could potentially disturb deer and elk in the analysis area. However, disturbance would be temporary. Noise from the operating vents would be continuous and audible. However, it is not expected to disturb these mammals, as they likely would become readily accustomed to it, as they are from the noise from the SUFCO Mine vents.

No impacts to macroinvertebrates are anticipated from exploration drilling or construction of vents. The only exception would be if the drill pads in the canyon bottoms or near perennial streams were too close to streams and impacted water quality. Mitigation is recommended in section 3.3 to minimize potential impacts to aquatic habitat.

3.2.3.2.5 Species of High Federal Interest

No impacts to migratory birds of high federal interest, other than those previously discussed for the golden eagle and flammulated owl, are anticipated from exploration drilling or construction of vents. Minor amounts of habitat would be removed for the construction of roads and drill pads (less than 17 acres total). However, the associated impact to bird habitat would be negligible.

3.2.3.2.6 Other Species

The locations of some of the drill pads and roads occur in potential habitat and near known locations of blue grouse. Therefore, drilling activities could potentially disturb grouse using the area. Disturbance would likely be short-term and include temporary displacement. No notable impacts to grouse habitat would occur.

The locations of some of the drill pads and roads are in the vicinity of a small number of ponds and springs in the analysis area. If drilling occurred in the ponds or springs, or associated hydric vegetation, amphibian habitat would be impacted. No drilling is proposed in the vicinity of the known boreal toad populations. Mitigation is recommended in section 3.3 to minimize potential impacts to amphibian habitat.

No impacts to reptiles, small mammals, or non-game birds are anticipated from exploration drilling or construction of vents. Minor amounts of habitat would be removed for the construction of roads and drill pads (< 17 acres total). However, the associated impact to wildlife habitat would be negligible.

3.2.3.3 Alternative 3: Standard Lease Terms and Conditions and Special Stipulations

Under Alternative 3, the tract would be leased with special coal lease stipulations in addition to the standard terms and conditions. Mining activities would include coal exploration drilling and the construction of four mine vent holes and associated structures. Exploration drilling would include the construction of associated drill pads, a staging area, and several short access roads. The number and location of drill pads and required time for exploration activities would likely be the same as under Alternative 2, since there would still be a need for geologic information throughout the tract.

3.2.3.3.1 Aquatic and Terrestrial Wildlife

Potential impacts to TEPS species, MIS, species of high federal interest, and other categories of wildlife would therefore be the same or similar to those outlined under Alternative 2, with two exceptions. The same mitigation measures suggested under Alternative 2 apply under this alternative.

Potential impacts to big-game species could be reduced under Special Stipulation 14. Measures could be put in place, as deemed necessary, that would curtail specific surface uses outside the mine development area during specified periods of the year in order to protect big-game wintering areas, elk calving and deer fawning areas, and other key wildlife habitat and/or activities. However, given that the impacts to deer and elk discussed under Alternative 2 were considered minor, implementation of such measures would unlikely be necessary.

Water depletion from the Colorado River system would not be expected under this alternative since under Special Stipulation 17, ground or surface waters identified for protection that may be impacted would have to be restored by the lessee in order to maintain riparian and fishery habitat, wildlife and other uses. Therefore, formal consultation with FWS for the four endangered fish species would not be warranted.

3.2.4 Wildlife Issue 4

Construction and operation of mine facilities and haul roads and coal traffic could remove habitat and associated noise/activity could displace dispersed wildlife (avoidance) including threatened, endangered, proposed, and sensitive species.

3.2.4.1 Alternative 1: No-Action

No leasing or mining would occur under this alternative. Construction and operation of mine facilities and haul roads and coal traffic would not occur. Therefore, habitat would not be removed and wildlife would not be displaced or dispersed from associated noise and mining activities.

3.2.4.2 Alternative 2: Standard Lease Terms and Conditions Only

Under Alternative 2, no mining facilities (storage units, offices, warehouses, truck loadouts, portals, conveyors, power lines, etc.) or roads would be constructed. Existing mining facilities and haul roads associated with the SUFCO mine would be used. These facilities are located outside of but adjacent to the analysis area. No aboveground mining activities would occur within the analysis area. Therefore, there would be no impacts to TEPS species, MIS, species of high federal interest, and other categories of wildlife associated with construction of mine facilities and hauling coal on haul roads. There would, however, be impacts to wildlife from subsidence of escarpments and spalling resulting from mine operations.

Escarpment failure could occur in lower Box Canyon, Greens Canyon, and Muddy Canyon, with the potential for failure considered very low, medium to high, and high, for the three canyons, respectively (MTI 2004). Potential impacts of escarpment failure and cliff face spalling are discussed below.

3.2.4.2.1 TEPS Fishes

Potential impacts to fish habitat from escarpment failure and spalling are discussed above in section 3.2.2.2.

3.2.4.2.2 TEPS Birds

Of the seven TEPS bird species discussed in this analysis, only the Mexican spotted owl and peregrine falcon use cliffs for nesting. Hypothetically, if escarpment failure were associated with a cliff nest site, the nest would be destroyed and breeding success for the raptor would be reduced until a new nest were built. Only peregrine falcons are known to nest in the analysis area, and the nest site is outside of the tract and the potential area of subsidence. Therefore, no impacts to TEPS birds would occur.

3.2.4.2.3 TEPS Mammals

Spotted bats and Townsend's big-eared bats often roost in rock crevices on steep cliff faces. Spotted bats have been observed in Box Canyon, Greens Canyon, and Muddy Canyon, thus escarpment failure and cliff face spalling could reduce suitable habitat for this species. Roost habitat for big-eared bats would also be affected, although it is unlikely that this species occurs in the analysis area.

3.2.4.2.4 MIS

As discussed in section 3.4.2.2, escarpment failure could impact nest sites and breeding success of cliff-nesting raptors. Golden eagles nest on cliff faces in Box and Muddy canyons. Therefore, escarpment failure would impact this species.

Impacts to mule deer and elk from escarpment failure and spalling would be minor or non-existent. Potential impacts would be limited to isolated incidents of rocks or boulders falling on individuals below cliff faces.

Potential impacts to macroinvertebrate habitat from escarpment failure and spalling are discussed above in section 3.2.2.2.

3.2.4.2.5 Species of High Federal Interest

Migratory birds of high federal interest that use cliffs for nesting include golden eagles, peregrine falcons, prairie falcons, and Mexican spotted owls. Black swifts also may use cliff habitats. The potential impacts of Alternative 2 on golden eagles, peregrine falcons, and spotted owls were discussed earlier in this section. No black swifts are present in the analysis area, and their presence is unlikely. However, potential habitat could be impacted by escarpment failure. A prairie falcon nest occurs about 0.5 miles from the tract boundary in Muddy Canyon. This nest could potentially be impacted by escarpment failure. However, the nest is on the northern side of the canyon, reducing the potential for impact.

3.2.4.2.6 Other Species

Potential impacts from escarpment failure for other species would be minor to non-existent. There is the potential for small mammal or reptile burrows to be crushed from large boulders. However, these impacts would occur in localized areas and would not impact populations. No impacts to amphibian habitat are expected. Failure of escarpments and spalling of cliff faces could potentially impact other species of birds that use cliff faces and rocky habitats, such as ravens, canyon wrens, and rock wrens.

3.2.4.3 Alternative 3: Standard Lease Terms and Conditions and Special Stipulations

Under Alternative 3, the tract would be leased with special coal lease stipulations in addition to the standard terms and conditions. Forest Service Special Stipulations associated with wildlife resources include stipulations 2 and 9. Since no surface uses would occur in the analysis area, special stipulations 4 and 14 for wildlife would not apply. Stipulation 2 requires that the lessee conduct an intensive field inventory for threatened and endangered species and migratory bird species of high federal interest. These surveys were conducted and survey results were used in the development of the Conceptual Mine Plan for the Muddy Tract for this alternative (MTI 2002), so that impacts to cliff-nesting raptors were avoided. Stipulation 9 requires that mining operations be conducted in a manner that would prevent surface subsidence, which could lead to escarpment failures and landslides as well as to damage or alterations of flow in perennial streams.

3.2.4.3.1 Aquatic and Terrestrial Wildlife

As under Alternative 2, no mining facilities or roads would be constructed. Existing mining facilities and haul roads associated with the SUFCO mine would be used. No aboveground mining activities would occur in the analysis area. Therefore, there would be no associated impacts to TEPS species, MIS, species of high federal interest, and

other categories of wildlife. In addition, special stipulation 9 defined above for Alternative 3 would eliminate the risk of escarpment failure. Therefore, impacts to wildlife associated with escarpment failure and cliff face spalling would not occur.

3.3 Mitigation and Monitoring Recommendations

- Include special stipulations to shorten longwall panels in order to prevent significant losses of surface and ground water flow in Muddy Creek.
- Conduct removal of debris, construction of fishways, and/or installation of culverts to enhance fish and aquatic habitat in areas that lose flow permanently, or where connectivity is interrupted as a result from subsidence (see Reeves et al. 1991).
- Conduct exploration drilling outside of streambeds and associated riparian areas (or riparian conservation areas or buffers, if defined) to reduce or eliminate potential impacts to aquatic habitat for fishes and macroinvertebrates.
- Conduct clearances for special status bird species (federally listed or proposed species, Forest Service sensitive species, MIS, and other raptors of federal interest) prior to mining activities. If species are observed, identify and map the location of nest sites.
- Implement seasonal and spatial buffers as described in Romin and Muck (1999) around any occupied goshawk, flammulated owl, golden eagle, or other known or identified raptor nest sites that may be impacted by mining-related activities.
- Conduct exploration drilling outside of/away from ponds, springs, and wetland habitats to reduce or eliminate potential impacts to amphibian habitat.

3.4 Cumulative Effects

Several land management activities have recently occurred, are currently occurring, or could occur in the near future in the Muddy analysis area. The activities that have the greatest potential to add cumulatively to the impacts of proposed mining on wildlife include cattle grazing, mining in the Pines Coal Tract, and recreation.

In general, livestock grazing poses a potential threat to aquatic habitat. Improper grazing practices can degrade streams, riparian habitats, and fish populations. It can also reduce the quality of habitat for terrestrial species associated with riparian systems. Degradation occurs when soils are compacted and the vegetation composition is changed. This can lead to increased runoff and erosion, reduced streambank vegetation and stability, changes to aquatic habitat, and adverse impacts to fish and other aquatic species (Platts 1991). Impacts from cattle grazing could add cumulatively to the impacts to aquatic habitat from mining-induced subsidence and escarpment failure.

Present and future mining activities in the Pines Coal Tract could affect fish and aquatic macroinvertebrate habitat in the Muddy Creek Tract, as small flow reductions and additional sediment inputs into Muddy Creek are anticipated (Forest Service 1999). Potential escarpment failure and cliff-face spalling, and mining-induced tension cracks associated with this mining lease could also add cumulatively to the impacts to other aquatic and terrestrial wildlife. No disturbance to terrestrial wildlife associated with mining activities would occur since the above ground activities for the Pines Tract occur outside of the Muddy analysis area.

Recreation in the analysis area is associated primarily with hunting. Increased visitation and vehicle use during the hunting season could add cumulatively to disturbances associated with coal exploration activities.

4.0 Literature Cited and Contacts

Andrews, R., and R. Righter. 1992. Colorado Birds. Denver Museum of Natural History, Denver, CO.

Beck, D.L. 1980. Wintering bald eagles in the Wells Resource Area, Elko District, Nevada, 1970-1980. U.S. Department of the Interior, Bureau of Land Management.

- Berg, L. 2002a. FAX addressed to Eric Duffin, Cirrus Ecological Solutions from Lois Berg, Aquatic Program Manager, Southeastern Region UDWR, regarding fish species presence in essential waters in Emery, Sanpete, and Sevier counties. May 10.
- Berg, L. 2002b FAX addressed to Rebecca Thompson, Cirrus Ecological Solutions from Lois Berg, Aquatic Program Manager, Southeastern Region UDWR, regarding fish surveys in the Muddy, Quitcupah, Straight Canyon, and Cottonwood drainages. September 18.
- Berg, L. 2002c Email addressed to Rebecca Thompson, Cirrus Ecological Solutions from Lois Berg, Aquatic Program Manager, Southeastern Region UDWR, regarding fish surveys in the Muddy, Quitcupah, Straight Canyon, and Cottonwood drainages. November 12.
- Bjornn, T.C., and D.W. Reiser. 1991. Habitat requirements of salmonids in Streams. American Fisheries Society Special Publication 19:83-138.
- BLM (USDI-Bureau of Land Management). 2003. Draft BLM Sage-Grouse Habitat Conservation Strategy. BLM Sage-Grouse Team, Washington, D.C. Released August 21, 2003.
- Bond, F. M., G. R. Craig, J. H. Enderson, A. W. Heggen, J. V. Kussman, D. L. Wills, A. Jenkins, and J. P. Hubbard. 1984. American Peregrine Falcon: Rocky Mountain and Southwest Population Recovery Plan. U.S. Fish and Wildlife Service, Denver, Colorado.
- Colorado Division of Wildlife (CDOW). 2001. Colorado Gap Analysis Project, Colorado Division of Wildlife, Habitat Section. Web access: <http://ndis.nrel.colostate.edu/cogap/gapframe.html>.
- Dalton, L.B., J.S. Price, and L.A. Romin. 1990. Fauna of Southeastern Utah and life requisites regarding their ecosystems. Publication No. 90-11. Utah Department of Natural Resources, Division of Wildlife Resources.
- Ehrlich, P. R., D. S. Dobkin, and D. Wheye. 1988. The Birder's Handbook: A Field Guide to the Natural History of North American Birds. Simon and Schuster/Fireside Books, New York.
- Forest Service (USDA Forest Service). 1986. Land and Resource Management Plan for the Manti-La Sal National Forest. 1986 version with 2003 revisions.
- Forest Service. 1993. Draft northern goshawk survey protocol and data collection forms. USDA-FS, Intermountain Region, Ogden, Utah. June, 1993.
- Forest Service. 1999. Final Environmental Impact Statement for the Pines Tract Project. Manti-La Sal National Forest. January.
- Forest Service. 2002a. Vegetation for the Manti-La Sal National Forest, Mani/Sanpitch Divisions. Manti-La Sal National Forest. April 22. <<http://www.fs.fed.us/r4/mantilasal/downloads/mantiveg.e00>>
- Forest Service. 2002b. Manti-La Sal National Forest News Release. Evidence of Canada Lynx Found on Manti-La Sal National Forest. 3 September.
- Forest Service 2003a. Manti-La Sal National Forest coal lease special stipulations. United States Department of Agriculture Forest Service. Manti-La Sal National Forest, Price, Utah.
- Forest Service. 2003b. Intermountain Region Proposed, Endangered, Threatened, and Sensitive Species. Known/Suspected Distribution by forest. December 2003.

- FWS (USDI Fish and Wildlife Service). 2002. Federally Listed and Proposed (P), Endangered (E), Threatened (T), and Candidate (C) Species and Habitat in Utah By County, dated August 2002. Salt Lake City, UT.
- Hart, J.M. and L.N. Berg. 2003. Colorado River cutthroat trout conservation and management in the Southeastern Region during 2002. Project F-44-R. Utah Department of Natural Resources, Division of Wildlife Resources, Salt Lake City. June.
- Hawkins, C. J. Ostermiller, and M. Vinson. 1998. Stream Invertebrate Sampling and Environmental Sampling Associated with Biological Water Quality Assessments: Field Protocols. Department of Fisheries of Wildlife and BLM National Aquatic Monitoring Center, Utah State University, Logan, UT. June.
- Hodson, R. 2004. Personal communication between Rebecca Thompson, Cirrus Ecological Solutions and Ron Hodson, Wildlife Biologist, Northern Region Utah Division of Wildlife Resources, regarding potential elk calving habitat in the Muddy analysis area. February 19.
- Jewkes, J. 2004a. Personal communication between Rebecca Thompson, Cirrus Ecological Solutions and Jeff Jewkes, Wildlife Biologist, Ferron Ranger District, regarding goshawk territory occupancy on the Manti-La Sal National Forest.
- Jewkes, J. 2004b. Digital coverage of modeled elk calving grounds emailed to Rebecca Thompson, Cirrus Ecological Solutions from Jeff Jewkes, Wildlife Biologist, Ferron Ranger District. February 23.
- Johnson, A., and D. Janetski. 2003. Field trip report: Quitchupah Creek and Water Hollow tributary. Report to the Utah Division of Wildlife Resources. Salt Lake City, Utah.
- Koehler, G. M. 1990. Population and habitat characteristics of lynx and snowshoe hares in north central Washington. *Canadian Journal of Zoology* 68:845-851.
- Lentsh, L., Y. Converse, and J. Perkins. 1997. Conservation agreement and strategy for Bonneville cutthroat trout (*Oncorhynchus clarki Utah*) in the state of Utah. Utah Division of Wildlife Resources, Publication 97-19, Salt Lake City, Utah.
- Licht, L. 1986. Food and feeding behavior of sympatric red-legged frogs, *Rana aurora*, and spotted frogs, *Rana pretiosa*, in southwestern British Columbia. *Canadian Field Naturalist* 100: 22-31.
- Mowat, G. K., G. Poole, and M. O'Donoghue. 2000. Ecology of lynx in northern Canada and Alaska. Chapter 9 in Ruggiero, L. F., K. B. Aubry, S. W. Buskirk [and others]. *Ecology and conservation of lynx in the United States*. University Press of Colorado, Boulder, CO. 480 p.
- MTI. 2002. Conceptual Mine Plan for the Muddy Creek Tract. Prepared by Maleki Technologies, Inc. Spokane WA. Prepared for Cirrus Ecological Solutions, LC.
- MTI. 2004. Assessment of the effects of surface impacts resulting from longwall mining in the Muddy Tract, Utah. Report to Cirrus Ecological Solutions, Logan, Utah, 2004. Maleki Technologies, Inc, Spokane, Washington.
- NatureServe Explorer: An online encyclopedia of life [web application]. 2002. Version 1.6 . Arlington, Virginia, USA: NatureServe. Available: <http://www.natureserve.org/explorer>. (Accessed: January 2003).

- Nelson, R.L., McHenry, M.L., and W.S.Platts. 1991. Mining. American Fisheries Society Special Publication 19:425-457.
- Oliver, G. 2001. Zoologist, Utah Division of Wildlife Resources. Personal communication with R. Thompson, Cirrus Ecological Solutions, regarding audible detections of bats. October.
- Oliver, G. V. 2000. The bats of Utah: A literature review. Utah Division of Wildlife Resources, Salt Lake City. Publication 00-14. 141 pages.
- Osmundson, D.B., and L.R. Kaeding. 1991. Flow recommendations for maintenance and enhancement of rare fish habitat in the "15 mile" reach during October-June. Final Report. U.S. Fish and Wildlife Service, Grand Junction, CO.
- Perkins, J.M. and J.R. Peterson. 1997. Bat survey for the SUFCO mine, Emery County, Utah. September, 1997. Salt Lake City, Utah.
- Peterson, R. T. 1990. A Field Guide to Western Birds. The Peterson Field Guide Series. Houghton Mifflin Company, Boston.
- Platts, W.S. 1991. Livestock grazing. American Fisheries Society Special Publication 19: 389-423.
- Reeves, G.H., Hall, J.D., Roelofs, T.D., Hickman, T.L., and C.O.Baker. 1991. American Fisheries Society Special Publication 19:519-556.
- Reynolds, R. T., and B. D. Linkhart. 1987. The nesting biology of flammulated owls in Colorado. Pages 239-309 in *Biology and Conservation of Northern Forest Owls*, Symposium Proceedings. USDA-Forest Service, Fort Collins, CO. Gen. Tech. Rep. RM-142.
- Reynolds, R.T., R.T. Graham, M.H. Reiser, R.L. Basselt, P.L. Kennedy, D.A. Boyce Jr., G. Goodwin, R. Smith, and E.L. Fisher. 1992. Management recommendations for the northern goshawk in the southwestern United States. Rocky Mountain Forest and Range Experiment Station, South-west Region Forest Service. USDA Forest Service, General Technical Report GTR-RM-217. 90 pp.
- Rodriguez 1998. Life History and Analysis of Endangered, Threatened, Candidate, Sensitive, and Management Indicator Species of Dixie National Forest. Dixie National Forest, Cedar City, Utah. March.
- Romin, L.A. and J.A. Muck. 1999. Utah Field Office Guidelines for Raptor Protection from Human and Land Use Disturbances. US Fish and Wildlife Service, Salt Lake City, Utah.
- Ruediger, B., J. Claar, S. Gniadek, B. Holt, L. Lewis, S. Mighton, B. Naney, G. Patton, T. Rinaldi, J. Trick, A. Vandehey, F. Wahl, N. Warren, D. Wenger, and A. Williamson. 2000. Canada Lynx Conservation Assessment and Strategy. USDA Forest Service, USDI Fish and Wildlife Service, USDI Bureau of Land Management, and USDI National Park Service. Missoula, MT.
- Ruggiero, L. F., K. B. Aubry, S. W. Buskirk, L. J. Lyon, W. J. Zielinski. 1994. The Scientific Basis for Conserving Forest Carnivores. American Martin, Fisher, Lynx, and Wolverine in the Western United States. USDA Forest Service, Rocky Mountain Forest and Range Experiment Station, Fort Collins, Colorado. General Technical Report RM-254.
- Sauer, J.R., J.E. Hines, and J. Fallon. 2003. The North American Breeding Bird Survey, Results and Analysis 1966 - 2002. Version 2003.1, USGS Patuxent Wildlife Research Center, Laurel, MD.

- Schmidt, J. 2004. Associate professor. Department of Aquatic, Watershed, & Earth Resources Utah State University, Logan, Utah. Personal communication.
- Sigler, W.F., and J.W. Sigler. 1996. Fishes of Utah: a natural history. University of Utah Press. Salt Lake City, Utah. 375pp.
- Spahr, R., L. Armstrong, D. Atwood, and M. Rath. 1991. Threatened, Endangered, and Sensitive Species of the Intermountain Region, USDA-Forest Service, Intermountain Region, Ogden, UT.
- Swanston, D.N. 1991. Natural Processes. American Fisheries Society Special Publication 19:139-179.
- Thomas, J.W., H. Black, R.J. Scherzinger, and R.J. Pedersen. 1979. Deer and Elk. Chapter 8 in Wildlife Habitats in Managed Forests: The Blue Mountains of Oregon and Washington. USDA Forest Service, Agricultural Handbook 553.
- Toone, R. 1993. General inventory for spotted bats (*Euderma maculatum*) on the Wasatch Plateau, Manti-La Sal National Forest, and the Old Woman Plateau and Thousand Lakes Mountain, Fishlake National Forest. Utah Natural Heritage Program, Utah Department of Natural Resources, Salt Lake City. January.
- Tyus, H.M. 1987. Distribution, reproduction, and habitat use of the razorback sucker in the Green River, Utah, 1979-1986. Transactions of the American Fisheries Society 116:111-116.
- UDWR (Utah Division of Wildlife Resources). 1992. Forest Owl Inventory Protocol. Utah Department of Natural Resources, Division of Wildlife Resources. Received from the Manti-La Sal National Forest, 2001.
- UDWR. 1997. Inventory of Sensitive Species and Ecosystems in Utah. Inventory of Sensitive Vertebrate and Invertebrate Species: A Progress Report. Department of Wildlife Resources. September 30.
- UDWR. 2001a. Elk aerial census data from January 2001 in the South Muddy, North Muddy/Ferron, North Horn, and South Horn. Received from Ron Hodson, Wildlife Manager for the Northern Region. Feb 2004.
- UDWR. 2001b. Utah big game annual report - 2001. Utah Department of Natural Resources, Utah Division of Wildlife Resources Publication Number 01-30. December 14.
- UDWR and UNHP (Utah Natural Heritage Program). 1992. Northern three-toed woodpecker (*Picoides tridactylus dorsalis*) inventory protocol. Received from the Manti-La Sal National Forest, 2001.
- Utah Conservation Data Center (UCDC). 2003. Utah Division of Wildlife Resources, UCDC Web home page, species information and Utah distribution maps. Available: <http://www.utahcdc.usu.edu/ucdc>.
- Utah Gap Analysis: An Environmental Information System. 1997. USDI National Biological Service and Utah State University. Contains species-specific maps depicting predicted critical, high value, substantial value, and limited value habitat in Utah. (<http://dwrcdc.nr.utah.gov/ucdc/>).
- Utah Gap Analysis: An Environmental Information System: DWR Neotrops Revision. 1999. USDI National Biological Service and Utah State University. Contains species-specific maps depicting breeding and wintering habitat in Utah. (<http://dwrcdc.nr.utah.gov/ucdc/>).
- Vinson, M. 2002a. Aquatic Macroinvertebrate Monitoring Report, Manti-La Sal National Forest, June 2001 - August 2001. National Aquatic Monitoring Center, Department of Aquatic, Watershed, and Earth Resources, Utah State University, Logan Utah. February.

- Vinson, M. 2002b. Aquatic Macroinvertebrate Monitoring Report, Manti-La Sal National Forest, May 2002 - September 2002. National Aquatic Monitoring Center, Department of Aquatic, Watershed, and Earth Resources, Utah State University, Logan Utah. December.
- Vinson, M. 2004. Aquatic Macroinvertebrate Monitoring Report for Greens, Cowboy, and White Mountain Cabin Creeks. June 2001 - September 2003. National Aquatic Monitoring Center, Department of Aquatic, Watershed, and Earth Resources, Utah State University, Logan Utah. January.
- Wai-Ping, V. and M.B. Fenton. 1989. Ecology of spotted bat (*Euderma maculatum*): roosting and foraging behavior. *Journal of Mammalogy*. 70(3):617-622.
- Wiley, D.W. 1992. Distribution and habitat ecology of Mexican spotted owls on the Colorado Plateau: Ann. Rep. 1991-1992. Utah Div. Wildl. Res. Salt Lake City, UT.

5.0 List of Preparers with Qualifications of Preparers

Rebecca Thompson (MS), Lead Wildlife Biologist, Cirrus Ecological Solutions, LC. Ms. Thompson has over nine years of experience in wildlife and forest ecology. She has worked in a wide array of natural systems, from coastal Oregon and Washington to the Intermountain West. Her graduate work at Oregon State University focused on habitat associations of red-backed voles and other small mammal species. Her work with the Forest Service focused on collection and analysis of aquatic vertebrate assemblages. Her work with the BLM included NEPA analysis of recreation and road improvement projects; managing wildlife habitat development projects; and coordinating and conducting timber sale clearance surveys for threatened, sensitive, and special status species. As a consultant she has functioned as the lead wildlife biologist and as a resource specialist on a variety of natural resource management projects (ski area developments/expansions, recreation permits, coal mining, grazing permits, and housing developments), conducting NEPA analyses and wildlife field surveys. She has conducted wildlife and botanical field surveys in range, forest, and riparian environments. Her field experience includes: wildlife field surveys for terrestrial and aquatic amphibians, small and arboreal mammals, mollusks, neo-tropical migrant birds, and federally listed threatened species and Forest Service sensitive species, and botanical surveys for vegetation community classifications (habitat typing) in sagebrush and subalpine habitats.

Anne Brown (MS), Wildlife Biologist, Cirrus Ecological Solutions, LC. Ms. Brown has 10 years of professional experience with wildlife and other natural resources in a wide variety of ecosystems ranging from forests to deserts to wetlands to agricultural systems, throughout North America. Her graduate work at Utah State University focused on sage grouse chick foraging ecology in sagebrush grasslands. Her extensive field experience includes point-count and broadcast avian surveys, nest searches and monitoring, behavioral observations, mist-netting, banding, color-marking, radio-tagging, and radio-tracking of various bird species, including passerines, game birds, hawks, and owls. She was also involved in small mammal, big game, and predator research. More recently, she assisted with water quality sampling of springs and the collection and processing of hydrologic data. She has demonstrated expertise with data analysis, technical writing, and editing.

Ernesto A. de la Hoz (MS), Aquatic Ecologist, Cirrus Ecological Solutions, LC. Mr. de la Hoz has over 7 years of experience in fisheries biology and aquatic ecology. He has worked in a wide variety of natural systems from Caribbean coral reefs to glacially formed lakes in the Intermountain West. His technical expertise includes fisheries biology, aquatic ecology, water quality, ichthyology, aquaculture, and modeling. He has worked on projects with sensitive/endangered species components and prepared environmental documents for government agencies in the U.S. and South America. In addition, Mr. de la Hoz has participated in stream, lake, coastal, and marine monitoring programs, water quality assessments, natural resource inventories, and numerous other fishery and aquatic ecology research projects. He holds a B.S. degree in Marine Biology from the University of Bogota- Jorge Tadeo Lozano (Colombia), a 2nd B.S. degree in Fisheries and Wildlife Management, and a M.S. degree in Ecology from Utah State University.

In addition to the list of preparers above, the following people (former Cirrus employees) assisted with wildlife field data collection and data analysis for this project:

- Don McIvor, Lead Wildlife Biologist and Crew Leader in 2001.
- Creed Clayton, Wildlife Technician and Crew Leader in 2001.
- Sunny McBride, Wildlife Technician in 2001.

6.0. APPENDICES: SUPPORTING DATA

Appendix A. Wildlife survey figures.

Appendix B. Macro-invertebrate taxa collected in the Muddy analysis area, 2001-2003.

Appendix C. Wildlife species observed in the Muddy analysis area, 2001-2003.

Appendix D. Small mammals.

Appendix E. Non-game birds.

Appendix A. Wildlife Survey Figures

- Figure A-1. Fisheries survey sites in the Muddy analysis area, 2001-2002.
- Figure A-2. Northern goshawk surveys in the Muddy analysis area, 2001-2003.
- Figure A-3. Flammulated owl surveys in the Muddy analysis area, 2001-2003.
- Figure A-4. Three-toed woodpecker surveys in the Muddy analysis area, 2001-2002.
- Figure A-5. Spotted bat surveys in the Muddy analysis area, 2001-2002.
- Figure A-6. Golden eagle surveys in the Muddy analysis area, 1998-2003.
- Figure A-7. Mule deer winter and summer range in the Muddy analysis area.
- Figure A-8. Elk winter and summer range in the Muddy analysis area.
- Figure A-9. Blue grouse sightings in the Muddy analysis area, 2001-2003.
- Figure A-10. Macroinvertebrate sampling sites in the Muddy analysis area, 2001-2003.
- Figure A-11. Species of high federal interest in the Muddy analysis area, 2001-2003.
- Figure A-12. Sage-grouse surveys in the Muddy analysis area, 2001-2003.
- Figure A-13. Pond locations and amphibian observations in the Muddy analysis area, 2001-2003.

Appendix B. Macro-
invertebrate taxa collected in the muddy analysis area, 2001-2003.

Order	Family	Subfamily/Genus/species
Phylum: Annelida		
Class: Oligochaeta		
	Lumbriculida	Lumbriculidae
		Lumbriculus
Phylum: Arthropoda		
Class: Arachnida		
Trombidiformes		
Class: Entognatha		
Collembola		
Class: Insecta		
Coleoptera		
Coleoptera	Dryopidae	Helichus
Coleoptera	Dytiscidae	
Coleoptera	Dytiscidae	Agabus
Coleoptera	Dytiscidae	Oreodytes
Coleoptera	Dytiscidae	Stictotarsus
Coleoptera	Elmidae	
Coleoptera	Elmidae	Optioservus
Coleoptera	Elmidae	Optioservus divergens
Coleoptera	Hydrophilidae	
Coleoptera	Hydrophilidae	Ametor
Coleoptera	Hydrophilidae	Berosus
Coleoptera	Hydrophilidae	Paracymus
Diptera		
Diptera	Ceratopogonidae	
Diptera	Ceratopogonidae	Bezzia
Diptera	Ceratopogonidae	Culicoides
Diptera	Ceratopogonidae	Probezzia
Diptera	Chironomidae	
Diptera	Chironomidae	Chironominae
Diptera	Chironomidae	Orthocladiinae
Diptera	Chironomidae	Tanypodinae
Diptera	Culicidae	
Diptera	Culicidae	Culiseta
Diptera	Dixidae	
Diptera	Dixidae	Dixa
Diptera	Dixidae	Dixella
Diptera	Dixidae	Meringodixa
Diptera	Dolichopodidae	
Diptera	Empididae	
Diptera	Empididae	Chelifera
Diptera	Empididae	Clinocera
Diptera	Empididae	Hemerodromia
Diptera	Empididae	Oreogeton
Diptera	Empididae	Wiedemannia
Diptera	Ephydriidae	
Diptera	Muscidae	
Diptera	Psychodidae	Pericoma
Diptera	Ptychopteridae	Ptychoptera
Diptera	Simuliidae	
Diptera	Simuliidae	Metacnephia
Diptera	Simuliidae	Prosimulium
Diptera	Simuliidae	Simulium
Diptera	Stratiomyidae	
Diptera	Stratiomyidae	Caloparyphus
Diptera	Stratiomyidae	Euparyphus
Diptera	Tabanidae	
Diptera	Tabanidae	Chrysops
Diptera	Tabanidae	Tabanus
Diptera	Tipulidae	
Diptera	Tipulidae	Dicranota
Order		
Family		
Subfamily/Genus/species		
Diptera	Tipulidae	Hexatoma
Diptera	Tipulidae	Ormosia

Diptera	Tipulidae	Pedicia
Diptera	Tipulidae	Tipula
Ephemeroptera		
Ephemeroptera	Ameletidae	Ameletus
Ephemeroptera	Baetidae	
Ephemeroptera	Baetidae	Acentrella
Ephemeroptera	Baetidae	Baetis
Ephemeroptera	Baetidae	Callibaetis
Ephemeroptera	Baetidae	Dipheter hageni
Ephemeroptera	Baetidae	Fallceon quilleri
Ephemeroptera	Ephemerellidae	
Ephemeroptera	Ephemerellidae	Drunella
Ephemeroptera	Ephemerellidae	Drunella coloradensis
Ephemeroptera	Ephemerellidae	Drunella doddsi
Ephemeroptera	Heptageniidae	
Ephemeroptera	Heptageniidae	Cinygmula
Ephemeroptera	Heptageniidae	Epeorus
Ephemeroptera	Heptageniidae	Rhithrogena
Ephemeroptera	Leptohyphidae	
Ephemeroptera	Leptohyphidae	Tricorythodes
Ephemeroptera	Leptophlebiidae	
Ephemeroptera	Leptophlebiidae	Paraleptophlebia
Ephemeroptera	Siphonuridae	
Ephemeroptera	Siphonuridae	Siphonurus
Hemiptera	Gerridae	
Hemiptera	Gerridae	Aquarius
Plecoptera		
Plecoptera	Capniidae	
Plecoptera	Chloroperlidae	
Plecoptera	Chloroperlidae	Suwallia
Plecoptera	Nemouridae	
Plecoptera	Nemouridae	Malenka
Plecoptera	Nemouridae	Zapada
Plecoptera	Nemouridae	Zapada cinctipes
Plecoptera	Nemouridae	Zapada columbiana
Plecoptera	Perlodidae	
Plecoptera	Perlodidae	Isoperla
Plecoptera	Taeniopterygidae	
Trichoptera		
Trichoptera	Brachycentridae	
Trichoptera	Brachycentridae	Brachycentrus
Trichoptera	Brachycentridae	Brachycentrus americanus
Trichoptera	Brachycentridae	Micrasema
Trichoptera	Hydropsychidae	
Trichoptera	Hydropsychidae	Hydropsyche
Trichoptera	Hydropsychidae	Parapsyche
Trichoptera	Hydroptilidae	
Trichoptera	Hydroptilidae	Hydroptila
Trichoptera	Hydroptilidae	Leucotrichia
Trichoptera	Limnephilidae	
Trichoptera	Limnephilidae	Hesperophylax
Trichoptera	Limnephilidae	Limnephilus
Trichoptera	Limnephilidae	Onocosmoecus
Trichoptera	Rhyacophilidae	Rhyacophila
Trichoptera	Rhyacophilidae	Rhyacophila alberta
Trichoptera	Rhyacophilidae	Rhyacophila brunnea
Trichoptera	Uenoidae	Neophylax
Trichoptera	Uenoidae	Neothremma
Trichoptera	Uenoidae	Oligophlebodes
Class: Maxillipoda		
Cyclopoida		
Harpacticoida		
Class: Maxillipoda, subclass copepoda		
Class: Ostracoda		
Podocopida		

Order	Family	Subfamily/Genus/species
Phylum: Mollusca		
Class: Bivalvia		
Veneroida	Pisidiidae	
Veneroida	Pisidiidae	Pisidium
Class: Gastropoda		
Basommatophora	Lymnaeidae	
Phylum: Nemata		
Phylum: Platyhelminthes		
Class: Turbellaria		

A total of 126 taxa were collected in 49 samples.

Appendix C. Wildlife species observed in the muddy analysis area, 2001 - 2003.

Birds

American crow	Mallard
American dipper	Mountain bluebird
American goldfinch	Mountain chickadee
American kestrel	Mourning dove
American robin	Northern flicker
Ash-throated flycatcher	Northern goshawk
Bald eagle	Northern pintail
Black-billed magpie	Northern pygmy-owl
Black-capped chickadee	Northern saw-whet owl
Black-chinned hummingbird	Olive-sided flycatcher
Black-headed grosbeak	Orange-crowned warbler
Blue grouse	Peregrine falcon
Brewer's blackbird	Pine grosbeak
Brewer's sparrow	Pine siskin
Broad-tailed hummingbird	Pinyon jay
Brown creeper	Prairie falcon (nest)
Brown-headed cowbird	Red-breasted nuthatch
Canyon wren	Red-naped sapsucker
Chipping sparrow	Red-tailed hawk
Clark's Nutcracker	Rock wren
Cliff swallow	Ruby-crowned kinglet
Common nighthawk	Ruffed grouse
Common poorwill	Sage-grouse
Common raven	Sage thrasher
Cooper's hawk	Sharp-shinned hawk
Dark-eyed junco	Short-eared owl
Downy woodpecker	Song sparrow
Dusky/Hammond's flycatcher	Sora
Evening grosbeak	Spotted sandpiper
Flammulated owl	Spotted towhee
Flycatcher sp.	Steller's jay
Golden eagle	Three-toed woodpecker
Gray flycatcher	Townsend's solitaire
Gray jay	Tree swallow
Gray vireo	Turkey vulture
Great horned owl	Vesper sparrow
Green-tailed towhee	Violet-green swallow
Hairy woodpecker	Vireo sp.
Hermit thrush	Warbling vireo
House wren	Western kingbird
Killdeer	Western meadowlark
Lincoln's sparrow	Western scrub jay
Long-eared owl	Western tanager
MacGillivray's warbler	Western wood-peewee

Bird, continued

White-crowned sparrow
White-throated swift
Williamson's sapsucker
Yellow warbler
Yellow-rumped warbler

Mammals

Badger
Beaver
Black bear
Black-tailed jackrabbit
Bushy-tailed woodrat
Cliff chipmunk
Cougar
Coyote
Elk
Golden-mantled ground squirrel
Moose
Mountain cottontail
Mule deer
Northern grasshopper mouse
Porcupine
Red squirrel
Snowshoe hare
Spotted bat
Uintah chipmunk
Uintah ground squirrel
Unidentified chipmunk species
Unidentified ground squirrel species
Unidentified pocket gopher species
White-tailed jackrabbit
Yellow-bellied marmot

Amphibians

Great basin spadefoot toad
Striped chorus frog
Tiger salamander

Reptiles

Eastern fence lizard
Sagebrush lizard
Short-horned lizard
Tree lizard
Unidentified garter snakes
Western terrestrial garter snake

Appendix D. Small mammals.

Small Mammal Species Potentially Occurring in the Muddy Analysis Area.							
Species	Ecological Association ¹	Elevation Range (CO) ²	Habitat ³	Relative Abundance ⁴	County Record ⁵	Predicted Habitat ⁶	Occur Expe
Masked shrew	all elevations	5,000-11,000	Prefers moist (riparian) habitats in mountainous areas	c	Unknown	Yes	Yes
Merriam's shrew	submontane /montane	4,500-9,600	Dry habitats, especially sagebrush; also grasslands, mixed woodlands	u	No	Yes	Possibl
Montane shrew	all elevations	5,300-11,500	Boreal forests and alpine habitats in mountainous areas	c	Yes	Yes	Yes
Dwarf shrew	submontane /montane	5,300-10,000	Rocky habitats in alpine tundra or subalpine conifer forests, talus slopes	k	No	Yes	No
Northern water shrew	submontane /montane	3,000-12,500	Near mountain streams, lakes, and marshes	c	Yes	Yes	Yes
Vagrant shrew	submontane /montane	Not mentioned	Near water	c	Unknown	Yes	Yes
Pallid bat	desert /submontane	3,000-7,000	Arid desert and grassland habitats, near water and rocky cliffs; also buildings	c	Unknown	Yes	Yes
Big brown bat	submontane /montane	3,000-10,000	Forests and urban areas; caves, mines, rock crevices, trees, buildings	c	Yes	Borderline	Yes*
Spotted bat	desert /submontane	Not mentioned	Desert, shrub steppe, moutain grassland or woodland; near cliffs	k	No	No	Observ
Silver-haired bat	montane	4,500-9,500	Forests/woodlands near water	c	Yes	Yes	Yes*
Red bat	submontane	3,000-5,000	Wooded areas near water, caves	r	No	Yes	No
Hoary bat	submontane /montane	3,000-10,000	Woodland habitats, roosts in trees	u	Unknown	Yes	Yes
California myotis	all elevations	4,500-7,500	Rock crevices, caves, buildings; forages near trees or over water	c	Unknown	No	Yes*
Long-eared myotis	submontane /montane	4,000-9,000	Prefers forested areas with rocky outcrops; also caves, mines, buildings	c	Unknown	Yes	Yes*
Small-footed myotis	all elevations	4,000-8,500	Wide variety of habitats, mostly forested; trees, crevices, caves, mines	u	Yes	Yes	Yes*
Little brown bat	all elevations	5,000-11,000	Buildings, caves, trees, mines; forages near trees and water	c	Unknown	Yes	Yes*
Fringed myotis	desert /submontane	3,000-7,500	Desert to woodland habitats; caves, mines, rock crevices, buildings	u	No	No	Possibl
Long-legged myotis	all elevations	4,000-12,500	Pine forests, deserts, riparian habitats; buildings, crevices, trees, mines	c	Unknown	Yes	Yes
Yuma myotis	desert /submontane	3,000-6,000	Various habitats near open water; caves, bridges, old buildings, mines	u	Unknown	Borderline	Yes*
Western pipistrelle	desert /submontane	3,000-6,000	Desert and rocky habitats near water, canyons; crevices, mines, caves, buildings	c	Yes	Borderline	Yes*
Townsend's big-eared bat	all elevations	3,000-9,500	Many habitat types, usually near forested areas; needs caves or mines	c	Yes	Yes	Possibl

Small Mammal Species Potentially Occurring in the Muddy Analysis Area.							
Species	Ecological Association¹	Elevation Range (CO)²	Habitat³	Relative Abundance⁴	County Record⁵	Predicted Habitat⁶	Expected⁷
Brazilian free-tailed bat	desert /submontane	3,000-9,500	Mostly warm, low, open habitats, including urban areas; caves, buildings	r	Yes	Yes	Possible
Ringtail	desert /submontane	3,000-9,500	Rocky deserts and woodlands, with cliffs and rocky outcrops, usually near water	c	Yes	Yes	Possible
Raccoon	desert /submontane	3,000-10,000	Wooded areas near water	c	Unknown	Yes	Yes
Marten	montane	9,000-12,000	Late-successional coniferous forests in remote mountainous areas	k (extirpated)	No	Yes	No
Ermine	submontane /montane	3,000-10,000	Prefers heavily wooded areas along streams	u	Unknown	Yes	Yes
Long-tailed weasel	all elevations	3,000-14,500	Habitat generalist: occurs in numerous habitat types; tolerant of human presence	c	Yes	Yes	Yes
Black-footed ferret	desert /submontane	3,000-10,000	Associated with prairie dog towns, for prey and den sites	k	No	Borderline	No
Mink	submontane /montane	3,000-14,500	Wetlands, marshes, and riparian areas, particularly near forested areas	l	Unknown	Yes	Possible
Badger	all elevations	4,500-14,500	Open areas such as grasslands and deserts, with sufficient soil for burrowing	c	Yes	Yes	Observed
Striped skunk	all elevations	3,000-10,000	Forest edges and open areas with sufficient soil for burrowing; also urban areas	c	Yes	Yes	Yes
Spotted skunk	all elevations	4,000-8,000	Rocky, brushy areas	c	Unknown	Yes	Yes
River otter	all elevations	4,000-12,500	Riparian habitats, from montane forests to desert canyons	r	No	No	No
Yellow-bellied marmot	submontane /montane	5,400-14,500	Rocky areas and meadows near forested areas	c	Yes	Yes	Observed
White-tailed prairie dog	all elevations	3,000-10,000	Open areas with well-drained soil for burrowing	l	Yes	Borderline	Possible
Utah prairie dog	Not mentioned	Not mentioned	Open areas below 9,000 ft with non-alkaline soils and succulent vegetation	Not mentioned	Yes	No	No
White-tailed antelope squirrel	desert /submontane	4,500-7,000	Desert and shrubland areas with sparse vegetation, rocky or gravelly soil	c	Yes	Yes	Yes
Uintah ground squirrel	submontane /montane	Not mentioned	Open, well-drained meadows, grasslands, and cultivated fields near water	c	Yes	Yes	Observed
Golden-mantled ground squirrel	submontane /montane	5,200-12,500	Rocky outcrops and talus slopes, open forests at high elevation and alpine tundra	c	Yes	Yes	Observed
Rock squirrel	desert /submontane	3,000-8,300	Rocky habitats	c	Yes	Yes	Yes
Cliff chipmunk	all elevations	5,500-7,000	Cliff dwellers in many types of habitats ranging from saltbrush to pine forests	u	Yes	Yes	Observed

Small Mammal Species Potentially Occurring in the Muddy Analysis Area.							
Species	Ecological Association¹	Elevation Range (CO)²	Habitat³	Relative Abundance⁴	County Record⁵	Predicted Habitat⁶	Occur Expe
Least chipmunk	all elevations	5,500-12,000	Many types of habitats, ranging from deserts to mountain forests	c	Yes	Yes	Yes
Uintah chipmunk	submontane /montane	6,500-12,000	Talus slopes and openings in coniferous forests, or forest edges	c	Unknown	Yes	Observ
Hopi chipmunk	Not mentioned	4,500-8,000	Prefers rocky habitats, especially with pinyon-juniper woodlands	Not mentioned	Unknown	Yes	Possibl
Red squirrel	montane	6,000-12,000	Dense stands of montane coniferous forests	c	Unknown	Yes	Observ
Northern flying squirrel	montane	Not mentioned	Mountainous areas, primarily in mature coniferous forests and riparian areas	c	Yes	Yes	Yes
Botta's pocket gopher	all elevations	4,000-8,500	Occurs in many types of habitats and soils	c	Yes	Yes	Yes
Northern pocket gopher	submontane /montane	5,000-14,500	Prefers deep, sandy soils, and high elevation prairies, meadows, and open forests	c	Yes	Yes	Yes
Ord's kangaroo rat	desert /submontane	3,000-8,000	Grasslands, shrublands, and woodlands with sandy soils and sparse vegetation	c	Yes	Yes	Yes
Plains pocket mouse	desert /submontane	3,000-7,500	Open grassland or desert habitats with sandy soils	c	Yes	No	Possibl
Great basin pocket mouse	desert /submontane	5,000-8,000	Arid grassland, sagebrush, and pinyon-juniper areas with sandy soils	c	Yes	No	Possibl
Bushy-tailed woodrat	desert /submontane	4,500-14,000	Rocky habitats (rocky outcrops), particularly at high elevations	c	Yes	Yes	Observ
Desert woodrat	desert /submontane	4,500-7,000	Rocky slopes and desert areas with sparse vegetation	c	Yes	Borderline	Possibl
Northern grasshopper mouse	desert /submontane	4,500-8,000	Grassland, desert, sagebrush, or pasture, with sandy soils and sparse vegetation	k	Unknown	Yes	Observ
Brush mouse	desert /submontane	4,000-8,500	Rocky sites with heavy brush	c	Unknown	Yes	Yes
Canyon mouse	submontane	4,500-8,000	Arid rocky habitats, such as deserts	c	Yes	Yes	Yes
Deer mouse	all elevations	3,000-14,000	Dryland habitats ranging from deserts to grasslands to coniferous forests	c	Yes	Yes	Yes
Pinyon mouse	submontane	4,000-8,500	Rocky terrain in pinyon-juniper, desert scrub, and woodland habitats	c	Yes	Yes	Yes
Western harvest mouse	desert /submontane	3,000-7,500	Dense vegetation near water; meadows, fields, weedy areas, grasslands	c	Yes	Yes	Yes
Western jumping mouse	desert /submontane	6,500-11,000	Mountain meadows near streams or marshes	c	Yes	Yes	Yes
Long-tailed vole	submontane /montane	3,500-14,000	Forests, mountain meadows, sagebrush, and riparian habitats	c	Yes	Yes	Yes
Montane vole	submontane /montane	6,000-14,500	Meadows and fields in mountain valleys	c	Yes	Yes	Yes

Small Mammal Species Potentially Occurring in the Muddy Analysis Area.

Species	Ecological Association ¹	Elevation Range (CO) ²	Habitat ³	Relative Abundance ⁴	County Record ⁵	Predicted Habitat ⁶	Expected ⁷
Meadow vole	all elevations	3,000-9,500	Variety of habitats ranging from dry open areas to marshes	c	Unknown	No	Possible
Water vole	submontane /montane	Not mentioned	Prefer alpine and subalpine meadows near fast-moving, clear streams	c	Yes	Yes	Yes
House mouse	all elevations	Not mentioned	Buildings and cultivated fields (weedy fields); usually associated with man	c	Unknown	Yes	Yes
Norway rat	all elevations	Not mentioned	Cities; near buildings, farms, dumps	c	Yes	Borderline	Possible
Black rat	all elevations	Not mentioned	Associated with man; buildings or fields near buildings; seaports	c	Unknown	Borderline	Possible
Muskrat	desert /submontane	3,000-14,500	Marshes, ponds; shallow, relatively still water surrounded with dense vegetation	c	Unknown	Yes	Yes
Beaver	all elevations	3,000-14,000	Depend upon permanent water sources within 1/4 mile of woodlands	c	Unknown	Yes	Observed
Porcupine	all elevations	3,000-14,500	Prefers coniferous or mixed forests; also riparian zones, deserts, shrublands	c	Yes	Yes	Observed
Pika	montane	10,000-14,500	Rocky slopes above the treeline (talus slopes and rockslides)	l	Yes	Yes	Yes
Snowshoe hare	submontane /montane	8,000-11,500	Mountain coniferous forests interspersed with thickets of aspen, willow, or alder	c	Yes	Yes	Observed
Black-tailed jackrabbit	desert /submontane	3,000-7,000	Open areas or brushlands of foothills, lower valleys, and desert areas	c	Yes	Yes	Observed
White-tailed jackrabbit	desert /submontane	4,000-14,500	Mountains statewide, also foothills and valleys in N Utah; mostly open areas	c	Yes	Yes	Observed
Desert cottontail	desert /submontane	3,000-7,000	Often concentrate in brushy areas along streams or dry washes	c	Yes	Borderline	Possible
Mountain (Nuttall's) cottontail	submontane /montane	6,000-14,500	Thickets, loose rocks, and cliffs; brushy areas along streams or dry washes	c	Unknown	Yes	Observed

¹ From Dalton et al. 1990; desert = 3,700 to 5,800 ft., submontane = 5,500 to 8,500 ft., and montane = 6,500 to 12,700 ft. elevation.

² From Colorado GAP analysis website (CDOW 2001); elevation range in ft.

³ Mostly based on narrative from UDWR web site (UCDC 2003) and also UDWR 1997 (for sensitive species), and Dalton et al. 1990 (for a few species).

⁴ From Dalton et al. 1990; c = common, u = uncommon. l = limited, r = rare. k = not known to inhabit the Wasatch Plateau area, Not mentioned et al. 1990.

⁵ From UDWR 1997, for sensitive species, and Durrant 1952 (Mammals of Utah); "unknown" if species was not mentioned in either publication.

⁶ Based on predicted habitat maps from UDWR web site (Utah Gap Analysis 1997).

⁷ Based on the information presented in the other columns of the table or whether we observed that species in the field.

⁸ From UDWR web site (UCDC 2003) and UDWR 1997 (for sensitive species).

* These bat species were observed during a 1997 survey conducted as part of the SUFCO and Dugout Mine's permit requirements (Perkins and

Appendix E. non-game birds

Non-Game Bird Species Potentially Occurring in the Muddy Analysis Area.							
Species	Ecological Association ¹	Elevation Range (CO) ²	Habitat ³	Seasonal Status ⁴	Relative Abundance ⁵	Predicted Habitat ⁶	Occur Expe
Common loon	all elevations	3,000-9,000	Large bodies of open water	Tr	u	No	Possibl
Western grebe	all elevations	3,000-9,500	Lakes, marshes, coasts	Tr	o	Borderline	Possibl
Clark's grebe	not mentioned	3,000-9,000	Lakes, marshes, coasts	Not mentioned	Not mentioned	Borderline	Possibl
Horned grebe	all elevations	3,000-6,000	Marshes, lakes, ponds, coasts	Su	u	No	Possibl
Eared grebe	all elevations	3,000-9,000	Shallow lakes and ponds with large macroinvertebrate communities; islands	Su	u	No	Possibl
Pied-billed grebe	all elevations	3,000-9,000	Riparian areas, shorelines, marshy wetlands	Su	c	Borderline	Possibl
American white pelican	all elevations	3,000-9,000	Reservoirs, large bodies of water	Tr	r	No	Possibl
Double-crested cormorant	all elevations	3,000-9,000	Ocean coasts, bays, lakes, rivers, reservoirs	Tr	r	Borderline	Possibl
Great blue heron	Desert /submontane	3,000-9,000	Shorelines of lakes and rivers, marshes	Su	u	Borderline	Possibl
Snowy egret	Desert /submontane	3,000-9,000	Marshes, lakes, coastlines	N/A	k	Borderline	Possibl
Black-crowned night-heron	Desert /submontane	3,000-10,000	Wetland areas, marshes along lakes	Su	u	No	Possibl
White-faced ibis	Desert /submontane	3,000-9,000	Marshy freshwater areas, swamps, ponds, rivers	Tr	r	Borderline	Possibl
Turkey vulture	all elevations	3,000-9,000	Open habitats in both lowlands and mountains	Su	c	Yes	Observ
Osprey	all elevations	3,000-10,000	Rivers, lakes, and ocean coasts	Su	r	Yes	Possibl
Cooper's hawk	all elevations	3,000-10,000	Coniferous and deciduous forests, riparian woodlands	Su	c	Yes	Observ
Northern goshawk	Montane; all (winter)	3,000-11,500	Mature mountain forests (conifer/aspens), usually within 1/4 mile of water	Yl	u	Yes	Observ
Sharp-shinned hawk	Submontane /montane	3,000-11,500	Forests and woodlands; heavy brush areas	Yl	u	Yes	Observ

Non-Game Bird Species Potentially Occurring in the Muddy Analysis Area.

Species	Ecological Association ¹	Elevation Range (CO) ²	Habitat ³	Seasonal Status ⁴	Relative Abundance ⁵	Predicted Habitat ⁶	Expected
Red-tailed hawk	all elevations	3,000-13,500	Open country with scattered trees, edge of woodlands	Yl	c	Yes	Observed
Rough-legged hawk	desert /submontane	3,000-9,500	Grasslands, fields, marshes, sagebrush flats and other open habitats (in winter)	Wt	c	Yes	Yes
Ferruginous hawk	desert /submontane	3,000-9,500	Grasslands and shrub steppes, edge of pinyon-juniper woodlands	Su	r	Yes	Possible
Swainson's hawk	all elevations	3,000-10,000	Shrub and grassland habitats, deserts, agricultural areas with scattered trees	Su	r	Yes	Possible
Northern Harrier	all elevations	3,000-9,500	Open habitats such as marshes, fields, and grasslands	Su	c	Yes	Possible
Golden eagle	all elevations	3,000-14,000	Open country, especially in mountainous regions; nests on cliffs or in trees	Yl	c	Yes	Observed
Bald eagle	all elevations	3,000-8,000	Coasts, rivers, lakes, or reservoirs, in open areas with available perching sites	Yl	e*	Borderline	Observed
Prairie falcon	all elevations	3,000-14,000	Open habitats (prairie, desert, alpine tundra) adjacent to cliffs	Yl	c	Yes	Observed
Peregrine falcon	all elevations	3,000-10,000	Open habitats from seacoasts to high mountains, open forests, cliffs, tall buildings	Yl	e*	Borderline	Observed
American kestrel	all elevations	3,000-10,000	Open or partly open habitats with scattered trees, also cultivated and urban areas	Su	c	Yes	Observed
Merlin	desert	3,000-9,000	Nests in coniferous woodlands or wooded prairies, often near water; open habitats during non-breeding season	N/A	k	Yes	Possible
American coot	all elevations	3,000-9,500	Ponds, lakes, marshes, rivers	Su	l	Borderline	Yes
Sora	desert /submontane	3,000-11,000	Freshwater wetlands, wet/flooded fields	Su	u	No	Observed
Virginia rail	desert /submontane	3,000-9,000	Freshwater or occasionally brackish marshes; also saltwater marshes in winter	Su	c	No	Possible
Sandhill crane	all elevations	3,000-10,000	Shallow wetlands, freshwater margins, also forages in open grasslands, meadows	Tr	u	Yes	Possible
Snowy plover	all elevations	3,000-6,000	Beaches, mudflats, saltflats, shorelines of rivers, lakes, ponds	Tr	r	No	Possible
Mountain plover	all elevations	4,500-6,000	Disturbed semi-arid grasslands (typically shortgrass prairie), also shrubsteppe	Tr	r	No	No
Semi-palmated plover	all elevations	3,000-6,000	During migrations: mudflats, beaches, flooded fields, marshes; breeds on tundra	Tr	u	No	Possible
Killdeer	all elevations	3,000-10,000	Fields, meadows, pastures, mudflats, freshwater margins	Su	c	Yes	Observed

Non-Game Bird Species Potentially Occurring in the Muddy Analysis Area.

Species	Ecological Association ¹	Elevation Range (CO) ²	Habitat ³	Seasonal Status ⁴	Relative Abundance ⁵	Predicted Habitat ⁶	Occur Expe
American (lesser) golden-plover	all elevations	3,000-6,000	During migrations: lake shores, marshes; breeds on tundra	Tr	u	No	Possibl
Black-bellied plover	all elevations	3,000-6,000	Lake shores in Utah during migrations; breeds on tundra	Tr	u	No	Possibl
Black-necked stilt	all elevations	3,000-6,000	Along freshwater and alkaline lakes, marshes, mudflats, shallow ponds, wet fields	Su	c	Borderline	Possibl
American avocet	all elevations	3,000-9,500	Shallow alkaline wetlands, ponds, mudflats of lakes and impoundments, estuaries	Su	c	Borderline	Possibl
Spotted sandpiper	all elevations	3,000-11,500	Rocky shorelines and marshy habitats, from sea level to alpine areas	Su	c	Yes	Observ
Sanderling	all elevations	3,000-6,000	Beaches, mudflats; breeds on arctic tundra	Tr	u	Borderline	Possibl
Baird's sandpiper	all elevations	3,000-9,500	Mudflats, shallow water, beaches; breeds on tundra	Tr	u	No	Possibl
Western sandpiper	all elevations	3,000-8,000	Breeds on tundra (no info about habitat used in Utah during migrations)	Tr	c	Borderline	Possibl
Pectoral sandpiper	all elevations	3,000-6,000	Wetlands, shallow water in marshes and at pond edges; breeds on tundra	Tr	u	No	Possibl
Least sandpiper	all elevations	3,000-9,500	Lakeshores and pond edges; some stay in Utah over winter near warm springs	Tr	c	No	Possibl
Semi-palmated sandpiper	not mentioned	3,000-6,500	Forages at water edges and on floating vegetation; breeds on tundra	Not mentioned	Not mentioned	Borderline	Possibl
Solitary sandpiper	all elevations	3,000-8,000	Streams, woodland swamps and ponds; also drainage ditches, puddles of manure	Tr	u	No	Possibl
Willet	submontane /montane	3,000-8,200	Shorelines of marshes and lakes, mudflats, coastal beaches	Su	c	No	Possibl
Common snipe	all elevations	3,000-10,500	Wetlands; nests in wet grass habitats	Yl	c	Borderline	Possibl
Short-billed dowitcher	desert /submontane	3,000-6,000	Shallow (salt)water with mud; breeds on tundra, wet meadows, forest bogs	Tr	u	No map	Possibl
Long-billed dowitcher	all elevations	3,000-9,500	Shallow water with mud, freshwater ponds; breeds on tundra, wet meadows	Su	c	No	Possibl
Marbled godwit	all elevations	3,000-9,500	Mud and alkali flats, shallow water; breeds in prairie wetlands, pastures, marshes	Tr	c	Borderline	Possibl
Long-billed curlew	all elevations	3,000-5,000	Uncultivated rangelands and pastures, grassy meadows, prairies	Su	r	Yes	No

Non-Game Bird Species Potentially Occurring in the Muddy Analysis Area.							
Species	Ecological Association¹	Elevation Range (CO)²	Habitat³	Seasonal Status⁴	Relative Abundance⁵	Predicted Habitat⁶	Expe
Lesser yellowlegs	all elevations	3,000-9,000	Marshes, mudflats, edges of ponds; breeds in open forests, muskeg, tundra	Tr	c	Borderline	Possibl
Greater yellowlegs	all elevations	3,000-9,500	Marshes, mudflats, lakes, ponds, flooded fields; breeds in muskeg, tundra, bogs	Tr	u	Borderline	Possibl
Red-necked phalarope	all elevations	3,000-9,000	Lacustrine wetlands, open water of bays, lakes, ponds, ocean; breeds on tundra	Su	c	No	Possibl
Wilson's phalarope	all elevations	3,000-9,500	Freshwater marshes, sloughs, wet meadows, islands; occasionally saline habitat	Su	c	No	Possibl
Herring gull	all elevations	3,000-6,000	Along coasts and near lakes, rivers, and landfills; breeds on tundra, coasts, islands	Tr	u	No	Possibl
California gull	all elevations	3,000-10,000	Mudflats, marshes, irrigated fields, lakes; also dumps, cities, agricultural lands	Su	c	Borderline	Possibl
Ring-billed gull	all elevations	3,000-9,500	Beaches, estuaries, water bodies, fields, parking lots, garbage dumps	Wt	c	Borderline	Possibl
Bonaparte's gull	all elevations	3,000-6,000	Feeds in open water; breeds in open coniferous woodlands near ponds, lakes	Tr	u	No	Possibl
Franklin's gull	all elevations	3,000-9,500	Breeds in prairie freshwater marshes, sloughs, marshy lakes; feeds in fields	Su	c	Yes	Possibl
Black tern	all elevations	3,000-8,500	Freshwater marshes, sloughs, wet meadows; nests in dense emergent vegetation	Su	c	No	
Caspian tern	all elevations	3,000-5,500	Large lakes, marshes, islands (in lakes and rivers), beaches, bays, coastal waters	Tr	r	No	No
Forster's tern	all elevations	3,000-8,500	Cattail and bullrush marshes along ponds, lakes, and sloughs; bays, sea coasts	Su	c	No	Possibl
Common tern	all elevations	3,000-6,000	Lakes, bays, sea coasts; breeds on islands and coastal beaches	Tr	u	Borderline	Possibl
Yellow-billed cuckoo	desert /submontane	3,000-6,000	Riparian (cottonwood/willow) or open woodlands with dense undergrowth, parks	Su	r	No	No
Barn owl	desert /submontane	3,000-6,000	Open and semi-open habitats, especially grassland, farmland, often near towns	Yl	u	Yes	Possibl
Northern saw-whet owl	submontane /montane	5,500-10,000	Dense conifer and mixed forests, wooded swamps, bogs, brushy areas	Yl	u	Yes	Observ
Short-eared owl	desert	3,000-8,500	Open habitats: grasslands, shrublands, meadows, marshes, tundra	N/A	k	No	Observ
Long-eared owl	all elevations	3,000-9,000	Woodlands bordered with open habitats, often near water; also parks, orchards	Yl	c	Yes	Observ
Burrowing owl	desert /submontane	3,000-9,000	Open grasslands and prairies; also golf courses, airports, cemeteries; need burrows	Su	l	Yes	Possibl

Non-Game Bird Species Potentially Occurring in the Muddy Analysis Area.

Species	Ecological Association ¹	Elevation Range (CO) ²	Habitat ³	Seasonal Status ⁴	Relative Abundance ⁵	Predicted Habitat ⁶	Occur Expe
Great horned owl	all elevations	3,000-11,500	Conifer or deciduous forests, woodlands, orchards, parks, wetlands, semidesert	YI	c	Yes	Observ
Northern pygmy-owl	submontane /montane	5,000-10,000	Woodland habitats; mixed conifer-deciduous and pine-oak forests	YI	u	Yes	Observ
Flammulated owl	montane	6,000-10,000	Montane forests, especially ponderosa pine associations	Su	u	No	Observ
Western screech-owl	all elevations	3,000-9,000	Woodlands (especially oak and riparian), scrub, orchards, woodlots, urban areas	YI	u	Yes	Possibl
Common nighthawk	all elevations	3,000-10,000	Open and semi-open habitats, such as grasslands, fields, open forests, towns	Su	c	Yes	Observ
Common poorwill	submontane	3,000-9,000	Semi-arid and arid grasslands and shrublands, rocky canyons, open woodlands	Su	c	No	Observ
White-throated swift	desert /submontane	5,500-10,000	Rocky cliffs and canyons in mountainous areas, occasionally coastal sea cliffs	Su	c	Yes	Observ
Black swift	submontane /montane	7,500-14,000	Cliffs near waterfalls in mountainous areas, steep rocky canyons, also sea cliffs	Su	u	Borderline	Possibl
Black-chinned hummingbird	desert /submontane	3,000-7,000	Riparian and open woodlands, shrublands, parks and gardens, often in arid regions	Su	c	Yes	Observ
Broad-tailed hummingbird	all elevations	3,000-11,000	Riparian areas, open woodlands, mountain forests near openings, brushy slopes	Su	c	Yes	Observ
Rufous hummingbird	submontane /montane	5,500-12,000	Coniferous forests with adjacent meadows, thickets, brushy slopes	Su (Tr?)	c	Yes	Yes
Calliope hummingbird	submontane /montane	5,500-8,500	Mountainous areas; open forests, meadows, and canyon, often along streams	Su	u	No	Possibl
Belted kingfisher	all elevations	3,000-9,000	Streams, lakes; particularly forested wetland areas near clear fishing waters	YI	u	Yes	Possibl
Northern flicker	all elevations	3,000-11,500	Open forest areas; nearly ubiquitous where nest sites and open ground occur	YI	c	Yes	Observ
Red-headed woodpecker	all elevations	3,000-5,500	Deciduous woodlands (esp. beech or oak), open areas with scattered trees, parks	Su	u	Not mentioned	No
Downy woodpecker	all elevations	3,000-11,000	Deciduous, mixed, and riparian woodlands, esp. aspen forests; parks, orchards	YI	c	Yes	Observ
Three-toed woodpecker	submontane /montane	8,000-11,500	Coniferous or sometimes mixed forests, burnt tracts; in Utah: spruce-fir forests	YI	c	Yes	Observ
Hairy woodpecker	all elevations	3,000-11,500	Deciduous or coniferous forests, wooded swamps, orchards, towns, parks	YI	c	Yes	Observ

Non-Game Bird Species Potentially Occurring in the Muddy Analysis Area.							
Species	Ecological Association¹	Elevation Range (CO)²	Habitat³	Seasonal Status⁴	Relative Abundance⁵	Predicted Habitat⁶	Expe
Williamson's sapsucker	submontane /montane	5,500-11,000	Conifer (fir, lodgepole pine) and mixed aspen-conifer forests; also aspen groves	Su	r	Yes	Observ
Red-naped sapsucker	all elevations	3,000-11,500	Coniferous forests with aspen, montane riparian woodlands	Yl	c	Yes	Observ
Olive-sided flycatcher	all elevations	3,000-11,500	Open coniferous and mixed forests with standing dead trees	Su	u	Yes	Observ
Western wood-pewee	all elevations	3,000-10,000	Coniferous and mixed forests, forest edges, riparian woodlands	Su	c	Yes	Observ
Cordilleran (western) flycatcher	all elevations	3,000-11,500	Deciduous and coniferous woodlands and forests, riparian areas	Su	c	Yes	Yes
Hammond's flycatcher	montane	7,000-11,000	Mature coniferous and aspen forests (desert, scrublands, woodlands in winter)	Su	u	No	Possibl
Dusky flycatcher	submontane /montane	5,500-11,000	Open and semi-open areas with dense brush; open conifer forests, aspen, willows	Su	c	Yes	Yes
Willow flycatcher	all elevations	3,000-10,000	Low scrub, swamps, thickets, especially willows, groves of small trees near water	Su	c	Borderline	Possibl
Gray flycatcher	submontane	5,000-7,000	pinyon-juniper, sagebrush, desert shrublands, open pine-oak woodlands	Su	u	Yes	Observ
Say's phoebe	desert /submontane	3,000-9,500	Open woodlands, farmlands, savannas, usually near water	Yl	c	Borderline	Possibl
Ash-throated flycatcher	desert /submontane	3,000-9,000	Scrub, chaparral, open and riparian woodlands, especially oak and pinyon-juniper	Su	c	Yes	Observ
Eastern kingbird	desert /submontane	3,000-9,000	Open and riparian woodlands, forest edges, agricultural areas, urban parks	Su	c	No	Possibl
Western kingbird	desert /submontane	3,000-10,000	Open and semi-open habitats: deserts, grasslands, agricultural and riparian areas	Su	c	Yes	Observ
Cassin's kingbird	submontane /montane	4,500-7,000	Mixed deciduous-conifer woodlands, dry savanna, scrub; also riparian areas	Su	u	Borderline	Possibl
Horned lark	desert/mont	3,000-9,000	Open habitats: desert, grasslands, agricultural areas, tundra, alpine meadows	Yl	c	Yes	Yes
Cliff swallow	desert /submontane	3,000-10,000	Open areas near running water; nests on cliffs, bridges, buildings, or in culverts	Su	c	Yes	Observ
Barn swallow	desert /submontane	3,000-10,000	Open habitats, especially near water; nests on rafters, bridges, cliffs	Su	c	Borderline	Possibl
Bank swallow	all elevations	3,000-8,000	Open country, especially near running water; nests along cliffs and bluffs	Su	c	No	Possibl
Northern rough-winged swallow	desert /submontane	3,000-9,000	Open country, especially near water; nests in earthen banks along waterways	Su	c	No	Possibl

Non-Game Bird Species Potentially Occurring in the Muddy Analysis Area.

Species	Ecological Association ¹	Elevation Range (CO) ²	Habitat ³	Seasonal Status ⁴	Relative Abundance ⁵	Predicted Habitat ⁶	Occur Expe
Tree swallow	all elevations	3,000-10,500	Open woodlands near water; coniferous forests in Utah; nests in tree cavities	Su	c	Yes	Observ
Violet-green swallow	all elevations	3,000-13,000	Coniferous or deciduous open forests or woodlands, particularly aspen	Su	c	Yes	Observ
Purple martin	montane	6,500-10,000	Open country, urban areas; in Utah: aspen-conifer forests near mountain lakes	Su	r	Yes	Possibl
Western scrub-jay	desert /submontane	5,000-7,000	Scrub oak, pinyon-juniper, brush, chaparral, pine-oak woodlands	Yl	c	Borderline	Observ
American crow	all elevations	3,000-10,000	Open habitats: agricultural areas, sparse woodlands, towns, orchards, tidal flats	Tr	o	Yes	Observ
Common raven	all elevations	5,000-14,000	Wide variety of habitats, often in mountainous or hilly areas	Yl	c	Yes	Observ
Steller's jay	montane; submontane (winter)	5,000-12,000	Coniferous and mixed forests, pine-oak woodlands	Yl	c	Yes	Observ
Pinyon jay	all elevations	5,000-7,000	Pinyon-juniper woodlands, pine woodlands	Yl	c	Yes	Observ
Gray jay	montane; submontane (winter)	5,000-11,500	Boreal and subalpine coniferous and mixed forests, open woodlands, bogs	Yl	u	Yes	Observ
Clark's nutcracker	submontane /montane	5,500-12,000	Breeds in montane coniferous forests; also uses pinyon-juniper in winter	Yl	c	Yes	Observ
Black-billed magpie	all elevations	3,000-13,000	Open country with scattered trees, brush, riparian and open woodlands, farmlands	Yl	c	Yes	Observ
Black-capped chickadee	all elevations	3,000-9,000	Deciduous or mixed woodlands, riparian woodlands, thickets, parks, suburbs	Yl	c	Yes	Observ
Mountain chickadee	all elevations	5,000-11,500	Montane coniferous forests; lower elevations in winter, including riparian areas	Yl	c	Yes	Observ
Plain titmouse	submontane	5,000-7,000	Pinyon-juniper and oak woodlands	Yl	u	Yes	Possibl
Bushtit	desert /submontane	5,000-8,500	Pinyon-juniper, oak scrub, chaparral, and other brushy habitats	Yl	c	Yes	Yes
Red-breasted nuthatch	montane	3,000-11,500	Montane coniferous and mixed forests, aspen; mature stands with decaying trees	Yl	c	Yes	Observ
White-breasted nuthatch	all elevations	3,000-11,500	Deciduous, mixed, and coniferous forests, riparian woodlands, pinyon-juniper	Yl	c	Yes	Yes
Pygmy nuthatch	montane	5,500-10,000	Pine forests (ponderosa pine in Utah, also yellow and Jeffrey)	Yl	c	Yes	Yes
Brown creeper	all elevations	3,000-11,500	Forested areas in high mountains, pine forests; lower elevations in winter	Yl	c	Yes	Observ

Non-Game Bird Species Potentially Occurring in the Muddy Analysis Area.

Species	Ecological Association ¹	Elevation Range (CO) ²	Habitat ³	Seasonal Status ⁴	Relative Abundance ⁵	Predicted Habitat ⁶	Expected ⁷
Canyon wren	all elevations	5,000-8,500	Cliffs, steep rocky canyons, rock outcrops, buildings, in arid and semi-arid areas	YI	c	Yes	Observed
Marsh wren	desert /submontane	3,000-9,000	Fresh- and brackish-water marshes with abundant reeds	Su	l	Yes	Possible
Rock wren	all elevations	3,000-12,000	Arid and semi-arid canyons, rock outcrops, talus slopes, scrublands, dry washes	YI	c	Yes	Observed
Bewick's wren	desert /submontane	3,000-7,000	Open woodlands, shrublands, farms, suburbs; pinyon-juniper and deserts in Utah	YI	c	Borderline	Possible
House wren	all elevations	3,000-11,000	Open and semi-open brushy areas; open woodlands, shrublands, farmlands, suburbs	Su	c	Yes	Observed
Winter wren	not mentioned	3,000-5,500	Forests (usually coniferous) or open habitats with dense brush or other groundcover	Not mentioned	Not mentioned	Yes	Possible
American dipper	submontane /montane	5,000-11,500	Fast-flowing mountain streams	YI	c	No	Observed
Blue-gray gnatcatcher	desert /submontane	5,000-7,000	Pinyon-juniper; deciduous forests, woodlands, swamps, scrub, chaparral, deserts	Su	c	No	Possible
Ruby-crowned kinglet	all elevations	3,000-11,500	Coniferous and mixed forests; mountains in summer, lower elevations in winter	YI	c	Yes	Observed
Golden-crowned kinglet	montane: submontane (winter)	3,000-11,500	Mountain coniferous forests in summer; lower elevation forests in winter	YI	c	Yes	Observed
Veery	desert /submontane	3,000-8,500	Shaded moist woodlands (esp. poplar, aspen) with understory	Su	u	Borderline	Possible
Hermit thrush	submontane /montane	3,000-11,500	Conifer, mixed, and deciduous forests, forest edges, riparian areas	Su	c	Yes	Observed
Swainson's thrush	submontane /montane	3,000-11,000	Dense shrublands, woodlands, and riparian areas, coniferous forest edges, orchards	Su	c	Yes	Yes
Townsend's solitaire	all elevations	3,000-12,000	Open montane coniferous forests on steep rocky slopes; lower elevations in winter	YI	c	Yes	Observed
Mountain bluebird	all elevations	3,000-13,500	Subalpine meadows, open forests and forest edges, rangelands, other open country	YI	c	Yes	Observed
Western bluebird	all elevations	3,000-8,000	Open, riparian, burnt, or cutover woodlands, open country with scattered trees	YI	r	No	Possible
American robin	all elevations	3,000-11,500	Forests, woodlands, scrublands, wetlands, fields, parks, suburbs	YI	c	Yes	Observed
Gray catbird	desert /submontane	3,000-7,000	Dense brush, shrublands, wooded suburbs, forest edges	Su	u	Borderline	Possible

Non-Game Bird Species Potentially Occurring in the Muddy Analysis Area.

Species	Ecological Association ¹	Elevation Range (CO) ²	Habitat ³	Seasonal Status ⁴	Relative Abundance ⁵	Predicted Habitat ⁶	Occur Expe
Northern mockingbird	desert /submontane	3,000-7,000	Low open areas with scattered trees, farmlands, second growth areas, suburbs	Su	u	Borderline	Possibl
Sage thrasher	submontane	3,000-14,000	Sagebrush and greasewood communities in low-elevation deserts	Yl	c	Yes	Observ
American (water) pipit	all elevations	3,000-14,000	Alpine and arctic tundra; winters at lower elevations	Yl	c	Yes	Yes
Cedar waxwing	desert /submontane	3,000-8,500	Woodlands, forest edges, well-planted suburbs	Wt	c	No	Possibl
Bohemian waxwing	all elevations	3,000-10,500	Breeds in coniferous and mixed woodlands; often frequents suburbs in winter	Wt	u	Yes	Possibl
Northern shrike	desert /submontane	3,000-9,500	Open deciduous and coniferous woodlands, taiga, scrub, thickets	Wt	u	No	Possibl
Loggerhead shrike	desert /submontane	3,000-9,000	Grasslands, pastures, fields with scattered trees, desert scrub, open woodlands	Yl	c	Yes	Yes
European starling	desert /submontane	not mentioned	Urban areas, farmlands, woodlands	Yl	c	Yes	Yes
Warbling vireo	montane	3,000-10,500	Open deciduous and mixed woodlands, riparian woodlands, montane aspen	Su	c	Yes	Observ
Solitary vireo	all elevations	3,000-8,000	Montane coniferous and mixed forests (ponderosa, PJ, aspen), riparian woodlands	Su	c	Yes	Yes
Gray vireo	submontane	5,000-7,000	Pinyon-juniper on arid slopes in Utah; oak-juniper, arid thorn scrub, chaparral	N/A	k	Yes	Observ
Yellow-rumped warbler	all elevations	3,000-11,000	Breeds in montane coniferous and mixed forests; lower elevations during migrations	Su	c	Yes	Observ
Magnolia warbler	montane	3,000-5,500	Open montane coniferous forests (spruce-fir-hemlock)	Tr	r	Not mentioned	No
Black-throated gray warbler	submontane	3,000-7,500	Open, dry coniferous and mixed forests, pinyon-juniper, chaparral, scrub, oak	Yl (Su?)	c	Yes	Yes
Yellow warbler	all elevations	3,000-10,000	Woodlands, scrublands, agricultural areas, suburbs, riparian areas	Su	c	Yes	Observ
Townsend's warbler	montane	3,000-12,000	Coniferous and mixed forests	Tr	u	Yes	Possibl
MacGillivray's warbler	all elevations	3,000-11,000	Dense riparian thickets (willow, alder), edges of coniferous or mixed forests	Su	c	Yes	Observ
Orange-crowned warbler	all elevations	3,000-9,000	Deciduous and mixed woodlands, riparian thickets, chaparral	Su	c	Yes	Observ
Nashville warbler	all elevations	3,000-7,000	Open deciduous, mixed, coniferous, or riparian woodlands, thickets	Tr	u	No	Possibl

Non-Game Bird Species Potentially Occurring in the Muddy Analysis Area.

Species	Ecological Association ¹	Elevation Range (CO) ²	Habitat ³	Seasonal Status ⁴	Relative Abundance ⁵	Predicted Habitat ⁶	Expe
Virginia's warbler	desert /submontane	3,000-10,000	Brush on dry hillsides, pinyon-juniper, scrub oak, chaparral, mountain mahogany	Su	c	Yes	Yes
Wilson's warbler	all elevations	3,000-13,500	Riparian woodlands, thickets and brush near water (esp. willow and alder bogs)	Su	c	Yes	Yes
American redstart	desert /submontane	3,000-6,500	Open deciduous and mixed woodlands, forest edges, second growth, riparian areas	Tr	r	No	Possibl
Common yellowthroat	submontane /montane	3,000-9,000	Marshes, riparian areas, brushy pastures, old fields, hedgerows, woodland margins	Su	l	No	Possibl
Yellow-breasted chat	all elevations	3,000-8,000	Dense brush or scrub, especially along streams and at swamp margins	Su	r	Yes	Possibl
Western tanager	all elevations	3,000-10,500	Breeds in conifer and mixed forests in mountains; riparian areas during migrations	Su	c	Yes	Observ
Lazuli bunting	all elevations	3,000-9,500	Arid brushy canyons, riparian thickets, chaparral, open woodlands	Su	c	Borderline	Possibl
Indigo bunting	desert /submontane	3,000-5,500	Brushy and weedy habitats, deciduous forest edges and clearings, weedy fields	Su	u	No	No
Rose-breasted grosbeak	all elevations	3,000-6,000	Deciduous forests and woodlands, second growth	Su	o	No	
Black-headed grosbeak	all elevations	3,000-11,500	Riparian woodlands and thickets, aspen, shrublands, open woodlands, pond edges	Su	c	Borderline	Oobserv
Sage sparrow	desert /submontane	3,000-7,000	Shrublands (sagebrush, arid brushlands, chaparral), grasslands, deserts	Su	u	Borderline	Possibl
Black-throated sparrow	desert /submontane	3,000-6,000	Dry brushy habitat, desert scrub, rocky uplands	Su	u	Borderline	Possibl
Lark sparrow	desert /submontane	3,000-9,000	Open habitats: grasslands, prairies, savannas, forest edges, cultivated areas	N/A	k	Yes	Possibl
Lincoln's sparrow	desert /submontane	3,000-12,000	Wet meadows, bogs, riparian thickets, mostly in mountains or boreal regions	Su	u	Borderline	Observ
Song sparrow	all elevations	3,000-10,500	Streamside thickets, marshes, wet meadows, bogs, forest edges, clearings, suburbs	Yl	c	No	Observ
Fox sparrow	all elevations	3,000-11,000	Forest undergrowth and edges, riparian thickets, scrub, montane brushland	Su	u	Yes	Yes
Vesper sparrow	all elevations	3,000-13,000	Dry grasslands and sagebrush, prairie, savanna, old fields, arid scrub, clearings	Su	c	Yes	Observ

Non-Game Bird Species Potentially Occurring in the Muddy Analysis Area.

Species	Ecological Association ¹	Elevation Range (CO) ²	Habitat ³	Seasonal Status ⁴	Relative Abundance ⁵	Predicted Habitat ⁶	Occur Expe
American tree sparrow	desert /submontane	3,000-10,000	Open areas with scattered trees, brush, scrub; fields, marshes, suburbs in winter	Wt	u	Yes	Possibl
Brewer's sparrow	desert /submontane	3,000-10,000	Arid brushland, mostly shrub steppe (sagebrush), also high desert scrub	N/A	k	Yes	Observ
Chipping sparrow	all elevations	3,000-11,000	Open coniferous forests, forest edges, oak, pine-oak, streamside habitats, parks	Su	c	Yes	Observ
White-crowned sparrow	all elevations	3,000-13,000	Stunted woody vegetation, wet and alpine meadows, farmlands, parks, roadsides	Yl	c	Yes	Observ
Harris' sparrow	desert /submontane	3,000-6,000	Stunted trees in forest-tundra ecotone; in winter: thickets, woodlands, scrub	Wt	u	No map	Possibl
Lark bunting	desert /submontane	3,000-9,000	Shortgrass prairie, grasslands, meadows, sagebrush	Tr	o	No	Possibl
Dark-eyed junco	montane: all (winter)	3,000-10,000	Coniferous and deciduous forests and edges, open woodlands, brushy areas, bogs	Yl	c	Yes	Observ
Green-tailed towhee	submontane /montane	3,000-11,500	Shrublands with interspersed conifers, pinyon-juniper, forest edges, riparian scrub	Su	c	Yes	Observ
Spotted (rufous-sided) towhee	desert /submontane	3,000-8,000	Brush, riparian thickets, dense shrubby areas, forest edges, chaparral, woodlands	Yl	c	Yes	Observ
Red-winged blackbird	desert /submontane	3,000-11,000	Freshwater and brackish marshes, riparian habitats, brushy areas near water, fields	Yl	c	Borderline	Possibl
Rusty blackbird	submontane	3,000-5,500	Moist coniferous woodlands, bogs, riparian habitats	Tr	o	Not mentioned	No
Brewer's blackbird	desert /submontane	3,000-12,000	Shrubby, brushy areas, riparian woodlands, aspen, marshes, farmlands, suburbs	Yl	c	Yes	Observ
Yellow-headed blackbird	desert /submontane	3,000-8,500	Freshwater marshes, wetlands	Su	c	No	Possibl
Bullock's (northern) oriole	desert /submontane	3,000-8,500	Open woodlands (cottonwood, willow, sycamore, oak), near fields or grasslands	Su	c	Borderline	Possibl
Scott's oriole	submontane	3,000-5,500	Yucca, pinyon-juniper, oak scrub, riparian woodlands, palms, Joshua trees-cactus	N/A	k	Yes	Possibl
Brown-headed cowbird	all elevations	3,000-12,000	Grasslands, prairies, fields, pastures, orchards, suburbs, woodlands, forest edges	Yl	c	Borderline	Observ
Western meadowlark	Desert	3,000-12,000	Grasslands, savannas, pastures, cultivated fields, mountain meadows, tidal flats	N/A	k	Yes	Observ
Common redpoll	all elevations	3,000-9,500	Subarctic coniferous forests, dwarf hardwoods, shrubby areas, tundra	Yl (Wt?)	c (r?)	No	Possibl

Non-Game Bird Species Potentially Occurring in the Muddy Analysis Area.

Species	Ecological Association ¹	Elevation Range (CO) ²	Habitat ³	Seasonal Status ⁴	Relative Abundance ⁵	Predicted Habitat ⁶	Expe
Pine siskin	submontane /montane	3,000-11,500	Coniferous and mixed forests, woodlands, parks, suburbs	Yl	c	Yes	Observ
Lesser goldfinch	desert /submontane	5,000-8,000	Scrub oak, pinyon-juniper, open areas with scattered trees or brush, fields, suburbs	Yl	c	Yes	Yes
American goldfinch	desert /submontane	3,000-9,000	Weedy fields, open deciduous and riparian woodlands, suburbs	Yl	c	Yes	Observ
Cassin's finch	all elevations	5,500-11,000	Semi-arid open coniferous forests at higher elevations, ponderosa pine	Yl	c	Yes	Yes
House finch	desert /submontane	3,000-10,000	Wide variety of habitats, arid scrub, open woodlands, urban areas, cultivated lands	Yl	c	Yes	Yes
Evening grosbeak	all elevations	5,500-10,000	Coniferous and mixed forests, second growth, parks	Wt	c	Yes	Observ
Black rosy-finch	montane; all (winter)	5,500-11,500	Barren rocky or grassy areas on alpine tundra, maritime island tundra, rocky cliffs	Yl	u	Yes	Yes
Grey-crowned rosy-finch	montane; all (winter)	5,500-11,500	Snowfields and rocky summits, alpine and maritime island tundra, rocky cliffs	Yl	u	Yes	Yes
Red crossbill	montane	6,000-11,000	Coniferous and mixed forests	Su	u	Yes	Yes
Pine grosbeak	montane; all (winter)	5,000-11,500	Open coniferous forests and forest edges	Yl	u	Borderline	Observ
House sparrow	desert /submontane	3,000-10,000	Human-modified habitats: agricultural, suburban, and urban areas; woodland edges	Yl	c	Yes	Possibl

¹ From Dalton et al. 1990; desert = 3,700 to 5,800 ft., submontane = 5,500 to 8,500 ft., and montane = 6,500 to 12,700 ft. elevation.

² From Colorado GAP analysis website (CDOW 2001); elevation range in ft.

³ Mostly based on narrative from UDWR web site (UCDC 2003) and on Ehrlich et al. 1988 (Birder's Handbook), also UDWR 1997 (for sensitive species).

⁴ From Dalton et al. 1990; Tr = transient, Su = summer resident, Wt = winter resident, Yl = yearlong resident, N/A = not known to inhabit the area mentioned in Dalton et al. 1990; when this information was inconsistent with other sources, corrected information was added in parentheses within the text.

⁵ From Dalton et al. 1990; c = common, u = uncommon, l = limited, r = rare, t = threatened, e = endangered (* The status of the bald eagle and peregrine falcon was delisted in 1999, o = occasional, k = plateau area, Not mentioned = species not mentioned in Dalton et al. 1990; when this information was inconsistent with other sources, corrected information was added in parentheses with a question mark).

⁶ Based on predicted habitat maps from UDWR web site (Utah Gap Analysis 1997 and 1999).

⁷ Based on the information presented in the other columns of this table or whether we observed that species in the field.

⁸ From UDWR web site (UCDC 2003) and UDWR 1997 (for sensitive species).

APPENDIX C

**BIOLOGICAL EVALUATION
AND
BIOLOGICAL ASSESMENT
SUFCO 2004 COAL EXPLORATION
DRILLING PROJECT
AND
SUFCO 2014 EXPLORATION WILDLIFE MEMO**

**BIOLOGICAL EVALUATION
and
BIOLOGICAL ASSESSMENT**

For the

**SUFCO 2004 HELICOPTER-ASSISTED
COAL EXPLORATION DRILLING PROJECT**

Ferron/Price Ranger District

Manti-La Sal National Forest

San Pete and Sevier Counties, Utah

Prepared by: /s/ Terry Nelson
Terry Nelson, Wildlife Biologist

Date 7/26/2004

Reviewed by: /s/ Pamela Jewkes
Pamela Jewkes, Forest Fisheries Biologist

Date 7/27/2004

/s/ Robert M. Thompson
Robert M. Thompson, Botanist

Date 7/26/2004

I. INTRODUCTION

The purpose of this Biological Evaluation /Biological Assessment (BE/BA) is to review the SUFCO 2004 Helicopter-Assisted Coal Exploration Drilling Project to determine the proposed action's potential effects on threatened, endangered, proposed or sensitive (TEPS) plant and animal species. TEPS species that may occur in the management unit where the proposed project is located are identified in Tables 1 through 4; those TEPS species that will not be affected by the project will not be carried through analyses in this report. Section 7 of the Endangered Species Act of 1973 (PL 93-205, as amended) requires federal agencies to ensure that any activity they authorize, fund, or carry out, does not jeopardize the continued existence of any wildlife species federally listed as threatened, endangered or proposed. Consultation with the U.S. Fish and Wildlife Service (Service) is required if threatened or endangered (T&E) species, or their critical habitat may be affected by proposed actions. One purpose of this BE/BA is to determine whether consultation with the service is necessary. This BE/BA is prepared in accordance with legal requirements set forth under Section 7 of the Endangered Species Act (16 U.S.C. 1536 (c)), and follows standards established in the Forest Service Manual (FSM 2671.2 and 2672.4).

A. PROPOSED ACTION

1. Summary of the Proposed Action

Canyon Fuel Co. Company has submitted a plan to conduct coal exploration and reclamation activities. Six drill holes are proposed for coal exploration during summer 2004. Five of the holes are proposed on unleased federal portions of the proposed Muddy Coal Area (Forest Service Surface/Federal Coal). One hole is proposed on Utah School and Institutional Trust Lands Administration (SITLA) portions of the Muddy Coal tract (Forest Service Surface/SITLA Coal). The project would be completed during the summer and early fall season, 2004. Access to three of the proposed drill sites would be along existing FS roads. Helicopters would be used to fly drill equipment to the other 3 remote sites where there are no existing roads. Since, helicopter-drilling techniques are proposed, there would be minimum disturbance (<100 ft² per site).

The proponent's proposed action as defined in its 2003 coal exploration license proposal is to access National Forest system lands, construct temporary drilling pads, drill holes to acquire needed geologic data from six coal exploration holes and reclaim disturbed areas on Forest Service managed land, using helicopter-assisted drilling methods. The proposed helicopter-assisted drilling project is outlined below:

- The planned drilling method is wireline core drilling from the surface down through to the lowest coal horizon. Equipment will include two heli-portable skid-mounted core drilling rigs together with all necessary equipment such as drill


rod trays, fuel tanks, water tanks, etc. The necessary equipment and vehicles include an 18,000 gallon frac tank, helicopter, jet fuel tank (trailer mounted), 4000 gallon water truck, two or three fifth-wheel flatbed trucks trailers used to haul drill equipment, four pick-up trucks, a covered tool supply trailer, and a geophysical logging truck.

- Hauling exploration equipment and transporting personnel to the staging area (see map) would be via frFDR 50007, 50044, and 50132 which traverses both the Fishlake and Manti-La Sal National Forests. Road-use permits would be obtained from the Forest Service before operation start.
- Site preparation would include removal of some vegetation with hand tools as needed for placement of the drill rig and needed equipment. Surface disturbance would be minimal; less than 100 square feet per site.
- The finished size of the hole will be nominally 2 3/16 inch diameter. Three-inch surface casing will be inserted through the surface alluvium and certain other intervals depending on hole conditions. Upon completion, holes would be geophysically logged.
- Soils would be protected from potential contamination by placement of brattice or similar impermeable material placed beneath mechanical equipment
- Water for drilling operations and road maintenance would be obtained from Muddy Creek and/or Quitchumpah Creek. Necessary arrangements would be made with shareholders and the Utah Division of Water Rights through a temporary water exchange permit. Completed drill holes would be plugged with a cement or cement/bentonite slurry to their full depth in accordance with BLM and Forest Service standards.
- Reclamation would include removal of equipment and trash immediately after hole completion. Topsoil would be scarified with hand tools. The disturbed areas would be reseeded (same as 2003 seed mix) with seed mix approved by the FS. The total plan, including reclamation, should be completed in 8 to 10 weeks.
- One hole may be completed as water monitoring well. Nominal 1.0 to 1.5 inch well screen and steel casing would be installed to below the deepest mineable coal seam. The screen zone would be sand packed and sealed from overlying strata and the overlying hole annulus would be cemented to the surface. Well casing with a locking lid would be left at the surface extending above the surface approximately two feet. The wellhead would be properly identified with either a brass marker or a welded-on identification. Once the monitor well is no longer in use, it would be completely plugged with a cement or cement/bentonite slurry to the top. The wellhead would be removed at the surface.

2. Description of the Project Location

The general locations are in San Pete and Sevier Counties about 10 miles northwest of the town of Emery, Utah. The proposed project area and drill hole locations are shown on Map 1. The proposed drill holes, lease tract administrator, location, depth and proposed access routes are summarized in the following table:

Drill Site	Tract	Location: T20S, R5E	Access Route
A	SITLA	SW, SE, Sec. 32	By Air FR 50044
B	BLM	NW, NW, Sec. 33	By Air FR 50132
C	BLM	SE, SW, Sec. 29	FR 50132
D	BLM	NW, NW, Sec. 32	FR 50132
E	BLM	NE, SE, Sec. 29	FR 50132
F	BLM	SE, NW, Sec. 29	By Air FR 50132



B. SPECIES OF CONCERN

1. Threatened and Endangered Plant Species

Table 1 lists plant species designated as threatened or endangered by the U.S. Fish and Wildlife Service (Service) that could occur within San Pete and Sevier Counties, Utah. No proposed plants are identified in San Pete or Sevier County. The table also describes habitats occupied by the threatened and endangered plants, the general distribution of their habitats, and whether or not those habitats are found within the project area. Habitat descriptions and distributions were obtained from Welsh et al. (1987) and Atwood et al. (1991). Habitat presence in the project area was determined through field visits and existing data review of soils, elevations, microclimate, and plant community composition within the project area. Although no formal rare plant surveys were conducted for this project, field reviews (including informal rare plant surveys) of East Mountain have been conducted by Bob Thompson (MLNF Botanist) on numerous occasions over the past several years. No listed plants or their habitats were detected in the project area or surrounding areas during any of the field reviews, nor are they expected to occur in this area of the Forest.

Table 1. Federally listed plant species that could occur in Sevier County, Utah and site-specific occurrence of their habitat within the project area.

SPECIES	SPECIES STATUS	HABITAT DESCRIPTION and DISTRIBUTION in SAN PETE and SEVIER COUNTIES	HABITAT PRESENT in PROJECT AREA?
Heliotrope Milkvetch <i>Astragalus montii</i>	Threatened	<i>Astragalus montii</i> was first discovered by Monte Lewis and Robert Thompson in 1976, and was listed as threatened in 1987. Its habitat is high elevation (10,500 to 11,000 ft) limestone barrens derived from the Flagstaff Geological Formation. All suitable habitat sites on the MLNF have been surveyed for populations of this species, it is known to occur in three populations on the MLNF. R. Thompson did not find <i>A. montii</i> in the proposed project area.	No
Last Chance Townsendia <i>Townsendia aprica</i>	Threatened	<i>Townsendia aprica</i> occurs in salt desert shrub and pinyon-juniper communities on clay or clay silt of the Arapien and Mancos Shale Formations, 5100' – 8000', occurs in Southeastern Emery County (off MLNF managed land). <i>T. aprica</i> was not found in the proposed project area.	No
Wright Fishhook Cactus <i>Sclerocactus wrightiae</i>	Endangered	<i>Sclerocactus wrightiae</i> occurs in salt desert shrub and juniper communities on Mancos Shale, 4800' – 6200', occurs in Southeastern Emery County (off MLNF managed land). <i>S. wrightiae</i> was not found in the proposed project area.	No
Winkler Cactus <i>Pediocactus winkleri</i>	Threatened	<i>Pediocactus winkleri</i> The Winkler cactus is a diminutive species that usually occurs solitarily. The plant grows in salt desert shrub communities at 4800 to 5200 feet elevation, in fine textured and poor quality saline substrates (Welsh et al. 1987). <i>P. winkleri</i> was not found in the proposed project area.	No

2. Sensitive Plant Species

Table 2 lists sensitive plant species on the Intermountain Regional Forester's sensitive species list that could occur on the Manti division of the MLNF. The table also describes habitats occupied by these sensitive plants, the general distribution of their habitats, and whether or not those habitats are found within the project areas. Habitat descriptions were obtained from Welsh, et al. (1987) and Spahr et al. (1991). Habitat presence in the project area was determined through field visits and existing data review of soils, elevations, microclimate, and plant community composition. Although no formal rare plant surveys have been conducted in the project area, field reviews (including informal rare plant surveys) the Pines Tract area have been conducted by Robert Thompson, MLNF Botanist. No sensitive plants or their habitats were identified in the project area or surrounding areas during any of the field reviews, nor are they expected to occur.

Table 2. Sensitive plants that could occur on the Manti Division of the Manti-La Sal National Forest (MLNF), and site-specific occurrence of their habitat within the project areas.

SPECIES	HABITAT DESCRIPTION, SPECIES OCCURRENCE IN THE PROJECT AREA AND CONSIDERATION IN THIS BE/BA
Link Trail Columbine <i>Aquilegia flavescens rubicunda</i>	Not considered. <i>Aquilegia flavescens rubicunda</i> occurs near spring seeps and perennial wetland sites on the east side of the Wasatch Plateau. The proposed project is located in fairly dry pinyon/juniper, sagebrush, mohogany habitats. This species was not found in the project area.
Creutzfeldt-flower <i>Cryptantha creutzfeldtii</i>	Not Considered. <i>Cryptantha creutzfeldtii</i> occurs in shallow, rocky, heavy clay soils; open Mancos shale slopes. It is endemic to central Utah in Carbon and Emery Counties at 5,000 to 6,500 ft. elevation. The proposed project is above 8,000 ft. elevation, and this species was not found in the project area.
Carrington Daisy <i>Erigeron carringtoniae</i>	Not Considered. <i>Erigeron carringtoniae</i> occurs in limestone outcrops and escarpments in subalpine vegetation type on wind blown ridge tops and snowdrift sites at high elevations of the Wasatch Plateau (9,000 to 11,000 feet). The proposed project is located in fairly dry pinyon/juniper, sagebrush, mohogany habitats on a fairly flat plateau area between 8,500 and 9,000 ft. elevation. This species was not found in the project area.
Canyon Sweetvetch <i>Hedysarum occidentale var. canone</i>	Not Considered. <i>Hedysarum occidentale var. canone</i> is usually found on sites that have a high water table, near springs or stream beds; riparian sites within the Pinyon/Juniper vegetation type at 5,500 to 7,000 ft. elevation. River birch and squaw brush are the most commonly associated species. It is endemic to Duchesne, and Carbon Counties. The proposed project is located in fairly dry pinyon/juniper, sagebrush, mohogany habitats. This species was not found in the project area.
Arizona Willow <i>Salix arizonica</i>	Not Considered. <i>Salix arizonica</i> occurs in wet meadows along perennial streams; occurs only in the Muddy Creek drainage on the MLNF. The proposed project is located in fairly dry pinyon/juniper, sagebrush, mohogany habitats, and will not impact the Muddy Creek drainage.
Musinea groundsel <i>Senecio musiniensis</i>	Not Considered. <i>Senecio musiniensis</i> occurs in limestone barrens and talus slopes of the southern Wasatch Plateau. This species was not found in the proposed project area, and is not expected to occur their.

Maguire Champion
Silene petersonii

Not Considered. *Silene petersonii* occurs at high elevations (10,000 to 11,800 ft.) on open calcareous and igneous soils derived from Flagstaff Limestone. The proposed project is located at approximately 8,700 ft. elevation in sedimentary soils. This species was not found in the project area.

3. Threatened and Endangered (T&E) Wildlife and Fish Species

Endangered species are species that have been identified, and listed in the Federal Register, by the U.S. Fish and Wildlife Service (Service) as being in danger of extinction throughout all or a significant portion of its range. Threatened species are species that have been identified, and listed in the Federal Register, by the Service as likely to become an endangered species within the foreseeable future throughout all or a significant portion of its range.

Table 3 lists wildlife and fish species designated as threatened or endangered by the U.S. Fish and Wildlife Service that could occur in San Pete or Sevier County, Utah. T&E species that may occur in San Pete or Sevier County but are not likely to occur in, and do not have suitable habitat in or near the proposed project area are also identified in Table 3, but they will not be considered further in this BE/BA. There are no proposed wildlife or fish species identified for Emery County.

Table 3. Listed and candidate wildlife and fish species that could occur in Emery County, Utah, and their potential for occurrence in the proposed project area and consideration in this BE/BA.

SPECIES	SPECIES STATUS	SPECIES OCCURRENCE IN THE PROJECT AREAS AND CONSIDERATION IN THIS BE/BA
Bald Eagle <i>Haliaeetus leucocephalus</i>	Threatened San Pete and Sevier Counties	Considered. A bald eagle pair has been known to nest in Emery County approximately 20 miles from the proposed project area. Bald eagles may occur incidentally in the proposed project area.
Yellow-billed Cuckoo <i>Coccyzus americanus occidentalis</i>	Candidate San Pete and Sevier Counties	Not Considered. The western yellow-billed cuckoo breeds in western U.S. states including Utah, and migrates to South America during winter. Cuckoos are riparian obligates. Nesting habitat is classified as dense lowland cottonwood/willow riparian forest characterized by a dense sub-canopy or shrub layer. In Utah, nesting habitats are found at elevations between 2,500 to 6,000 feet. They appear to require large tracts (100 to 200 acres) of contiguous riparian nesting habitat (Parrish et al. 1999). The proposed project is located in fairly dry pinyon/juniper, sagebrush, mohogany habitats at between 8,500 and 9,000 ft. elevation, there is no suitable habitat for this species in or near the project area.
Canada Lynx <i>Lynx canadensis</i>	Threatened San Pete County	Not Considered. The proposed project is located in open fairly dry pinyon/juniper, sagebrush, mohogany habitats, which does not provide suitable habitat for the Canada lynx.
Utah Prairie Dog <i>Cynomys parvidens</i>	Threatened San Pete and Sevier Counties	Not Considered. Utah prairie dogs are found in areas where there are deep, well-drained soils, burrows extend straight down for about 10-15 ft. and then branch into horizontal tunnels. They feed on insects (particularly cicadas), where available. Their preferred vegetative food type is alfalfa, but they generally prefer grasses over forbs and shrubs. Moist palatable forage must be available throughout the summer. The proposed project is located in fairly dry pinyon/juniper, sagebrush, mohogany habitats with mostly shallow soils over Castle Gate sandstone. No evidence of Utah prairie dogs was found in or near the project area.

4. Sensitive Wildlife and Fish Species

Sensitive species are species that are recognized by the Regional Forester as needing special management attention in order to prevent them from becoming threatened or endangered.

Table 4 lists the Intermountain Regional Forester's list of sensitive wildlife species that could occur on the Manti Division of the Manti-La Sal National Forest (MLNF). Sensitive wildlife species that do not occur or have suitable habitat in or near the proposed project area, or species that would not be impacted by proposed activities within the project area, are identified in Table 4 and will not be considered further in this Biological Evaluation/Biological Assessment (BE/BA).

Table 4. Sensitive wildlife and fish species that could occur on the Manti Division of the MLNF, and their potential occurrence in the proposed project area and consideration in this BE/BA.

SPECIES	SPECIES OCCURRENCE IN THE PROJECT AREAS AND CONSIDERATION IN THIS BE/BA
Spotted Bat <i>Euderma maculatum</i>	Considered. In Utah, the spotted bat likely occurs throughout the state. It is known to use a variety of vegetation types from approximately 2,500 to 9,500 feet, including riparian, desert shrub, ponderosa pine, montane forests, open pastures and meadows. Spotted bats roost alone in rock crevices high up on steep cliff faces. There are potentially suitable roosting cliffs near the proposed project area. Spotted bats may occasionally forage in the sagebrush/shrub habitat in the vicinity of the proposed project, and in the nearby ponderosa pine habitat.
Townsend's Big-eared Bat <i>Plecotus townsendii pallescens</i>	Considered. In Utah, Townsend's big-eared bats roost and hibernate in caves and mines, they also roost (but not hibernate) in buildings (Oliver 2000). These bats use juniper/pine forests, shrub/steppe grasslands, deciduous and mixed conifer forests. There is potentially suitable roost sites and forage habitat in or near the proposed project area.
Greater Sage Grouse <i>Centrocercus urophasianus</i>	Considered. Sage grouse are generally found where there are large tracts of sage brush habitat with a diverse and substantial understory of native grasses and forbs or in areas where there is a mosaic of sagebrush, grasslands, aspen Wet meadows, springs, seeps, or other green areas within sagebrush shrublands are generally needed for the early brood-rearing period. There is suitable breeding habitat near the proposed project area.
Northern Goshawk <i>Accipiter gentilis</i>	Not Considered. Goshawks forage in fairly dense (generally greater than 40 percent canopy cover) conifer forests, and they nest in even denser stands (generally greater than 60 percent canopy cover), many nest and forage sights contain an aspen component. The proposed project is located in fairly dry pinyon/juniper, sagebrush, mohogany habitats. There is no suitable goshawk habitat in or near the project area.
Peregrine Falcon <i>Falco peregrinus</i>	Considered. Peregrine falcons may travel more than 18 miles from the nest site to hunt for food, however average foraging distance from the eyrie extends out to 10 miles, with 80 percent of peregrine falcon foraging occurring within a mile of the nest. The nearest known peregrine falcon eyrie is located approximately 3 1/2 miles from the project area. Nesting peregrine falcons may forage in the vicinity of the proposed project.
Flammulated Owl <i>Otus flammeollus</i>	Not Considered. Flammulated owls prefer mature ponderosa pine/Douglas fir forests with open canopies, but they can be found in second growth ponderosa pine, aspen and mixed conifer forests that contain a ponderosa pine component. The proposed project is located in fairly dry pinyon/juniper, sagebrush, mohogany habitats, and will not alter or disturb flammulated owl habitat.
Three-toed woodpecker <i>Picoides tridactylus</i>	Not Considered. Three-toed woodpeckers are found in northern coniferous and mixed forest types up to 9,000 feet elevation. Forests containing spruce, grand fir, ponderosa pine, tamarack, and lodgepole pine are used. Nests may be found in spruce, tamarack, pine, cedar, and aspen trees. The proposed project is located in fairly dry pinyon/juniper, sagebrush, mohogany habitats, and will not alter or disturb three-toed woodpecker habitat.
Spotted Frog <i>Rana pretiosa</i>	Not Considered. Spotted frogs are most commonly found in cold, still, permanent water in such habitats as marshy edges of ponds or lakes, in algae-grown overflow pools of streams, and near flat water springs with emergent vegetation. This frog has a broad distribution throughout the previously glaciated regions of British Columbia. They also occur in the Rocky Mountains of Alberta, and have patchy distribution in the United States, from Washington to Montana and south to Nevada and Utah. In Utah, the spotted frog occurs in isolated populations, and is considered to be a relict from the last ice age. The spotted frog has not been found on the Manti - La Sal National Forest or in the proposed project area.
Colorado Cutthroat Trout	Not Considered. Colorado cutthroat trout require cool, clear water in streams with well vegetated banks, which provides cover and bank stability. Deep pools and structures such as boulders and logs provide instream cover. This species is believed to have formerly been widespread in lakes, rivers, and streams in Utah, however now it is

*Oncorhynchus
clarki pleuriticus*

limited to isolated headwater streams and other rigorous environments where other species such as rainbow trout and Yellowstone cutthroat trout have not been introduced. Colorado cutthroat trout are not found in the proposed project area, and the project would not adversely impact drainages where it is found.

**Bonneville
Cutthroat Trout**
*Oncorhynchus
clarki utah*

Not Considered. Bonneville cutthroat trout require cool, clear, well-oxygenated water and the presence of clean, well-sorted gravels with minimal fine sediments for successful spawning. They are found at high, moderate and low elevations in small head water streams in the Bonneville basin (USDI 2001b). Bonneville cutthroat trout are not found in the proposed project area, and the project would not adversely impact drainages where it is found.

II. TES SPECIES POTENTIALLY AFFECTED BY THE PROJECT

A. THREATENED AND ENDANGERED SPECIES

Bald Eagle

Bald eagle nests are typically located in multi-storied (uneven aged) coniferous forest stands that contain elements of old growth structure, and are located near bodies of water that support prey species. Nest trees are generally one of the largest trees in the stand, which provides good visibility and a clear flight path to and from the nest (Stalmaster 1987). Bald Eagles typically construct large, conspicuous stick nests in sizeable trees.

Prey species commonly include fish, waterfowl, jackrabbits, and carrion; results of food-habit studies have indicated that bald eagle diets included: 56 percent fish, 28 percent birds, 14 percent mammals, and 2 percent miscellaneous sources (Stalmaster 1987).

Bald eagles spend over 90 percent of the daylight hours perching. Important perch sites generally have 3 fundamental elements: a direct view of potential food sources, located within 50 meters of water, and are located in areas isolated from human disturbance (Stalmaster 1987).

Unlike nesting and perch sites, roosting sites are not necessarily located close to water; during breeding season, nesting adults often roost in the nest or at the nest tree (Stalmaster 1987). Roost sites generally provide thermal cover, and are isolated from human disturbance. Bald eagles often roost communally during winter.

During the winter, Bald Eagles tend to concentrate wherever food is available; food availability is probably the single most important factor affecting winter eagle distribution and abundance, but availability of night roosts and diurnal perches are also fundamental elements of bald eagle winter range. Eagles are often attracted to wintering concentrations of waterfowl. In some regions, such as Utah, carrion can also be an important food source. At wintering areas, Bald Eagles often roost in large groups. These communal roosts are located in forested stands that provide protection from harsh weather.

There are only a few known nesting pairs of bald eagles in Utah. There is a bald eagle nest site located approximately 20 miles from the proposed project area, and located approximately 7 miles from Forest Service managed land. A nesting pair had been observed at this site during the nesting and fledgling period for several years prior to 1997. This nesting territory was not occupied in 2001 or 2002. The nest was blown out of the tree in the winter of 2003, and a pair built a new nest approximately ½ mile southeast of the old one, but did not nest successfully in 2003. The pair worked on the nest again in early 2004, but did not nest. A 1997 study by N. Boschen indicated that the pair did not forage on national forest system lands; nesting adults and fledglings were found to forage within a 5 mile radius of the nest tree (Boschen, 1997). No bald eagles are known to nest on Manti-La Sal NF managed lands. Most bald eagle sightings on the Forest have been at Joe's Valley Reservoir and Huntington Canyon during late fall and early winter prior to freeze over.

D. SENSITIVE SPECIES

Spotted Bat

The spotted bat ranges from Mexico through the western states to the southern border of British Columbia; it is probably widely distributed in low numbers throughout western North America (Toone 1994). And it probably occurs throughout Utah, but its distribution appears to be patchy. Hasenyager (1980) thought that "the range of the spotted bat in Utah could incorporate the southern third of the state and central portions of the west desert where suitable roosts exist, excluding the higher portions of the central mountain range." Habitat occupied by this bat ranges from low desert to montane coniferous forests normally below 8,000 feet in elevation (Watkins 1977). They have been found in a variety of habitat types including open ponderosa pine, desert shrub, pinyon/juniper, and open pasture and hay fields. In Utah, the spotted bat has been captured in several habitats: lowland riparian habitat (open meadows), desert shrub communities (sagebrush/rabbitbrush), ponderosa pine forest, montane grassland (grass/aspen), and montane forest and woodland (grass/spruce/aspen). This species has also been occasionally found in or on buildings in Utah towns and cities (Oliver 2000).

They typically roost singly in crevices in steep cliff faces. Cracks and crevices in limestone or sandstone cliffs provide important roosting sites (Spahr et al. 1991), especially where rocky cliffs occur in proximity to riparian areas. Day roosts and maternal roosts are typically within small (up to 6 cm) cracks and crevices in cliff faces (Toone 1994). The relative inaccessibility of cliff roosts may insulate spotted bats from human disturbance, but the species has been observed roosting (and foraging) near campgrounds (Toone 1994). Spotted bats are thought to feed mainly on moths high above the vegetation canopy. They forage alone after dark using echolocation, which is effective for fast flight feeding on tympanate moths (moths that can detect ultra-sonic sounds). As is common with many bats, spotted bats may forage a considerable distance (up to 6 miles) from roost sites (Toone 1994).

Roosting habitat in the Wasatch Plateau region is likely to occur in numerous cliffs along the edges of the plateau and on canyon walls that cut through the plateau. It is likely that spotted bats forage in a variety of habitats on the Plateau that are located within 6 miles of suitable roost cliffs and at elevations lower than 9,500 ft. Various surveys on the MLNF have detected spotted bats in several major canyons (and their tributaries) on the east side of the plateau, including Muddy, Ferron, Straight, Cottonwood, and Huntington Canyons (Perkins and Peterson 1997, and Sherwin et al. 1997). These surveys also detected spotted bats near Joes Valley Reservoir and Trail Mountain.

Observations made during the 1997 surveys on the MLNF indicated that spotted bats tolerate at least moderate human disturbance while foraging. Surveys were conducted at several sites near roads with light to moderate vehicular traffic (Crandall Canyon, Huntington Canyon, Straight Canyon), including tandem coal trucks. Spotted bats were observed foraging at low elevation sites, within 30 meters of the right-of-way. The fact that spotted bats were relatively common in active and previously mined areas may imply that subsidence caused cliff failures have not dramatically affected resident populations (Sherwin, et al. 1997).

Townsend's Big-eared Bat

Townsend's big-eared bats occur throughout North America, from British Columbia to southern Mexico; from California to South Dakota and western Texas and Oklahoma. They are widely distributed throughout the Intermountain Region, and they occur throughout Utah (Oliver 2000). They inhabit a wide variety of xeric and mesic habitats including: desert scrub, sagebrush, chaparral, deciduous and coniferous forests including, but not limited to pinyon/juniper, ponderosa pine, spruce/fir, redwood, mixed hardwood/conifer, and oak woodlands (Pierson et al. 1999), and their distribution is strongly correlated with the availability of caves or cave-like roosting habitat such as mines, buildings with cave-like attics, diversion tunnels or bridges (Pierson et al. 1999). They require relatively spacious, relatively cool cave-like roost sites; generally at least 30 meters in length, and at least 2 meters high with temperatures ranging from -2.0 to 13.0° C (Pierson et al. 1999).

These bats are relatively sedentary, and do not migrate long distances; generally seasonal movements are less than 32 km (Pierson et al. 1999). Detections in Utah have ranged from 3,300 feet to 9,520 feet (Oliver 2000). In Utah, night roosts are found in mines and caves; day roosts and maternity roosts are found in mines, caves and buildings (Oliver 2000).

Townsend's big-eared bats are insectivorous; a lepidopteran specialist eating mostly moths (Pierson et al. 1999). They forage after dark using echolocation on the wing (Sphar et al. 1991); a late flyer, emerging from the roost primarily after dark; well after sunset (Pierson et al. 1999).

Breeding occurs at winter sites between October and February, and parturition occurs in late spring and early summer. Each female usually gives birth to a single offspring.

Females and young roost in communal nurseries, which range in size from 12 to 200 individuals. The offspring fly at three weeks and are weaned in six to eight weeks. Nurseries break up by August.

During winter, these bats roost singly or in small clusters in hibernacula from October to February. They don't migrate, but will move to different roost locations within hibernacula and may even move to different hibernacula during a winter in response to temperature changes.

Most of the bat surveys conducted on the MLNF that employed the use of mist nets or bat detectors have not revealed Townsend's big-eared bats (Perkins and Peterson 1997, and Sherwin et al. 1997). This is not unusual, as these bats are most commonly located during direct surveys of roosts (Oliver 2000).

There is potentially suitable Townsend's big-eared bat foraging habitat in and around the proposed project area.

Greater Sage Grouse

Sage grouse are sagebrush ecosystem obligates; they occur in mosaics of sagebrush, grasslands, and aspen, and are associated with both tall and short species of sagebrush in foothills, sagebrush shrublands, and mountain slopes. They do not occur in pinyon-juniper woodlands or in shadscale shrublands (Paige and Ritter 1999). At one time sage grouse were found in virtually all areas where sage brush (especially *Artemisia tridentata*) occurred in Western North America. It is hypothesized that the sage grouse breeding population circa 1800 was 1.1 million birds. Today, the estimated breeding population is 0.2 million (Parrish et al. 2002).

In Utah, sage grouse inhabit sagebrush habitat of the Colorado Plateau and Great Basin geographic regions from 6,000 to 9,000 ft. elevation. During spring, they use sagebrush habitats for breeding, feeding, roosting, nesting and rearing young (Connelly et al. 2000). Large, relatively continuous sagebrush stands, often exceeding 50 sq. mi., are needed to provide all habitat characteristics used by sage grouse; summer home ranges may be as small as 1 to 2.5 square miles, and annual home ranges may be as large as 577 square miles (Page and Ritter 1999).

Sage grouse males appear to form breeding leks opportunistically at sites within or adjacent to potential nesting habitat. Leks are typically established in openings within large sagebrush stands; openings include old lakebeds, low sagebrush flats, ridge tops, burn areas, and other open areas within sagebrush stands (Connelly et al. 2000). Most nests are placed under sagebrush in stands that provide higher than average canopies and lateral cover (Connelly et al. 2000). Nest sites also generally contain taller and denser grass cover than average. As sage brush habitats dry out during summer sage grouse use a wider variety of habitats including meadow and riparian habitats. Hens with broods move to areas that support succulent vegetation including forbs (Parrish et al. 2002).

Sites used by broods have been reported to have twice as much forb cover as independent sites (Connelly et al. 2000).

There suitable sage grouse habitat near the proposed project area.

Peregrine Falcon

The peregrine falcon is cosmopolitan, ranging from coast to coast in North America. Pesticide accumulation in the mid 1900s drove the peregrine to the verge of extinction, and by 1965 fewer than 20 pairs were known west of the Great Plains. In 1990 there were 326 known pairs in the southwest region (Rodriguez 2002). The peregrine falcon was federally listed as an endangered species in 1970, and again in 1984. With the help of reintroductions and pesticide controls (primarily banning DDT, which caused eggshell thinning and drastically low reproduction), the peregrine falcon population increased sufficiently to be de-listed in 2000.

Peregrine falcon preferred nesting habitat is on cliff faces with recesses or protected shelves, although reintroduced birds regularly nest on man-made structures such as towers and high-rise buildings. A wide variety of habitats are used for foraging, including riparian woodlands, open country near rivers and marshes, coniferous and deciduous forest edges, shrublands, and prairies. They prey on a wide variety of birds including pigeons, shorebirds, waterfowl, grouse and other small to mediums sized terrestrial birds. Peregrine falcons may travel up to 18 miles from their nest site to forage for food, however a 10 mile radius around the nest is an average hunting area, and 80% of foraging occurs within a mile of the nest (Spahr et al. 1991). The nearest known peregrine falcon eyrie is located approximately 3 ½ miles from the project area. Nesting peregrine falcons may forage in the vicinity of the proposed project.

III. AFFECTED ENVIRONMENT

The proposed project is located on a relatively high elevation plateau on the Castle Gate sandstone formation. There are a variety of habitats on this plateau including: pinyon/juniper, mixed conifer dominated by ponderosa pine, mixed conifer and aspen, mountain brush, sage brush and perennial grassland habitats. There are 6 drill sites in the proposed project plan: 5 drill holes are located in sagebrush dominated habitat and one drill hole is located in mountain brush habitat that includes sagebrush/rabbit brush, service berry and mahogany.

IV. ANALYSIS OF EFFECTS

This analysis of effects is based on the existing conditions within the project planning area. The analysis reviews the potential “direct and indirect effects” of the proposed SUFCO Exploration Drilling project on threatened, endangered and sensitive (TES)

species, and the expected “cumulative effects” that could potentially accrue to TES species if project activities add cumulatively to other past, present or reasonably foreseeable future actions to impact the species of concern.

The past, present or reasonably foreseeable future actions that may add incrementally to impacts of the proposed CANYON FUEL CO. Company SUFCO (Federal Coal Leases U-76195 and U-63214) Coal Exploration Drilling Project (2003) include:

- Other exploration drilling activity
- Disbursed recreational activity
- Road construction and maintenance

A. Threatened and Endangered Species

This analysis of effects is based on the existing conditions within the project planning area. The analysis reviews the potential “direct and indirect effects” of the proposed SUFCO 2004 Helicopter-Assisted Coal Exploration Drilling Project on threatened, endangered and sensitive (TES) species, management indicator species (MIS), and priority migratory bird species. This report also states the expected “cumulative effects” that would potentially accrue to TES, MIS and priority migratory bird species if proposed project actions add cumulatively to other past, present or reasonably foreseeable future actions to impact the species of concern.

The past, present or reasonably foreseeable future actions that may add incrementally to impacts of the proposed SUFCO 2004 Helicopter-Assisted Coal Exploration Drilling Project include:

- Other exploration drilling activity
- Disbursed recreational activity
- Road construction and maintenance

A. Threatened and Endangered Species

Bald Eagle

Direct and Indirect Effects: There are no landscape characteristics in the vicinity of the proposed project that would attract bald eagles to the area; there are no water bodies that would provide suitable bald eagle forage habitat in or near the project area. The project area is not known or expected to be used by nesting, wintering or foraging bald eagles. However, bald eagles may occur incidentally while in transition during migration or dispersal during late fall or early winter months. These occurrences would only be incidental and of short duration, and the proposed project would not alter bald eagle habitat. Therefore, the proposed project is not likely to directly or indirectly affect the bald eagle.

Cumulative Effects: Since the proposed project is not likely to exert direct or indirect affects on the bald eagle, no cumulative affects will accrue to this species because of the SUFCO 2004 Helicopter-Assisted Coal Exploration Drilling Project.

B. Sensitive Species

Spotted Bat

Direct and Indirect Effects: There are numerous cliff faces that could provide suitable spotted bat roost habitat within 2 miles of the proposed project area. The nearest suitable roost habitat is located approximately ½ mile from the nearest drill site. Activity during project implementation would not likely disturb roosting bats, and the project would not directly or indirectly impact spotted bat roost habitat.

The project would be implemented over a short period of time (7 plus days at each drill site) over small segments of the landscape that potentially provides suitable spotted bat forage habitat. However, since project activity would occur during daylight hours, it would not impact the nighttime foraging spotted bat. The proposed project would not appreciably directly or indirectly impact spotted bat foraging habitat.

Cumulative effects: Since the proposed project would not likely exert appreciable direct or indirect impacts on the spotted bat, no appreciable cumulative affects would accrue to this species because of the SUFCO 2004 Helicopter-Assisted Coal Exploration Drilling Project.

Townsend's big-eared Bat

Direct and Indirect Effects: There are a number of alcoves and cave like structures located within 2 miles of the proposed project area. Activity during project implementation would not likely disturb roosting bats; the project would not directly or indirectly impact Townsend's big-eared bat roost habitat.

The project will be implemented for a short period of time (7 plus days at each drill site) over small segments of the landscape that potentially provides suitable Townsend's big-eared bat forage habitat. However, since project activity would occur during daylight hours, it would not impact this nighttime foraging species. The proposed project would not appreciably directly or indirectly impact Townsend's big-eared bat foraging habitat.

Cumulative effects: Since the proposed project would not likely exert appreciable direct or indirect affects on the Townsend's big-eared bat, no appreciable cumulative affects would accrue to this species because of the SUFCO 2004 Helicopter-Assisted Coal Exploration Drilling Project.

Greater Sage Grouse

Direct and Indirect Effects: The proposed project would occur outside the greater sage grouse lekking and breeding season, the project would not modify lekking or breeding habitat, and the project would not occur in brood rearing habitat. Therefore, the proposed project would not likely appreciably directly or indirectly impact the greater sage grouse.

Cumulative effects: Since the proposed project would not likely exert appreciable direct or indirect effects on the greater sage grouse, no appreciable cumulative effects would accrue to this species because of the SUFCO 2004 Helicopter-Assisted Coal Exploration Drilling Project.

Peregrine Falcon

Direct and Indirect Effects: The nearest known peregrine falcon eyrie is located approximately 3 ½ miles from the project area. Falcons may travel more than 18 miles from the nest site to hunt for food, however a 10 mile radius around the nest is an average hunting area, with 80% of foraging occurring within a mile of the nest. Nesting peregrine falcons may forage in the vicinity of the proposed project. Project implementation would not occur during the peregrine nesting period, and would only temporarily impact localized areas within potential forage habitat; therefore the proposed project would not likely appreciably directly or indirectly impact the peregrine falcon.

Cumulative effects: Since the proposed project would not likely exert appreciable direct or indirect effects on the peregrine falcon, no appreciable cumulative effects would accrue to this species because of the SUFCO 2004 Helicopter-Assisted Coal Exploration Drilling Project.

V. DETERMINATION OF EFFECTS

A. Threatened and Endangered Species

PLANT SPECIES: The U.S. Fish and Wildlife Service lists three plant species that could occur in or have suitable habitat in San Pete or Sevier County, Utah including: Heliotrope Milkvetch, Last Chance Townsendia, and Wright Fishhook Cactus. These plant species or their habitats do not occur within the proposed project area. Therefore, it is my determination that the proposed SUFCO 2004 Helicopter-Assisted Coal Exploration Drilling Project will not affect the Heliotrope Milkvetch, Last Chance Townsendia, or Wright Fishhook Cactus.

FISH SPECIES: The U.S. Fish and Wildlife Service (Service) does not list any fish species as threatened, endangered, proposed or candidate species for San Pete or Sevier County, Utah.

WILDLIFE SPECIES: The U.S. Fish and Wildlife Service lists four wildlife species that could occur in or have suitable habitat in San Pete or Sevier County, Utah including: bald eagle, Southwestern willow flycatcher, Western yellow-billed cuckoo and Utah prairie dog. Determinations for these species follow:

Bald Eagle

There are no landscape characteristics in the vicinity of the proposed project that would draw bald eagles to the area; the project area is not known or expected to be used by nesting, wintering or foraging bald eagles, and the proposed project will not impact bald eagle nest, forage or winter habitat. Therefore, it is my determination that the proposed SUFCO 2004 Helicopter-Assisted Coal Exploration Drilling Project will not affect the bald eagle.

Yellow-billed Cuckoo

Cuckoos are riparian obligates; nesting habitat is classified as dense lowland cottonwood/willow riparian forest characterized by a dense sub-canopy or shrub layer. In Utah, nesting habitats are found at elevations between 2,500 to 6,000 feet. They require large tracts (100 to 200 acres) of contiguous riparian nesting habitat (Parrish et al. 1999). The proposed project is located in relative dry upland habitat at approximately 8,700 ft. Therefore, it is my determination that the proposed SUFCO 2004 Helicopter-Assisted Coal Exploration Drilling Project will not affect the yellow-billed cuckoo.

Canada Lynx

The proposed project is located in open fairly dry pinyon/juniper, sagebrush, mohogany habitats, which does not provide suitable habitat for the Canada lynx. Therefore, it is my determination that the proposed SUFCO 2004 Helicopter-Assisted Coal Exploration Drilling Project will not affect the Canada Lynx.

Utah Prairie Dog

Utah prairie dogs are found in areas where there are deep, well-drained soils. The proposed project is located in an area where there are shallow soils on Castle Gate sandstone. Therefore, it is my determination that the proposed SUFCO 2004 Helicopter-Assisted Coal Exploration Drilling Project will not affect the Utah prairie dog.

B. Sensitive Species

PLANT SPECIES: The Intermountain Regional Forester lists seven sensitive plant species that could occur on the Manti Division of the Manti-La Sal National Forest including: Link Trail Columbine, Creutzfeldt-flower, Carrington Daisy, Canyon Sweetvetch, Arizona Willow, Musinea groundsel and Maguire Campion. None of these sensitive plants or their habitat occurs within or near the proposed project area. Therefore, it is my determination that the proposed SUFCO 2004 Helicopter-Assisted Coal Exploration Drilling Project will have no impact on the Link Trail Columbine, Creutzfeldt-flower, Carrington Daisy, Canyon Sweetvetch, Arizona Willow, Musinea groundsel or Maguire Campion.

FISH AND AMPHIBIAN SPECIES: The Intermountain Regional Forester lists three sensitive fish and amphibian species that could occur on the Manti Division of the Manti-La Sal National Forest including: spotted frog, Colorado cutthroat trout and the Bonneville cutthroat trout. Determinations for each of these species follow:

Spotted Frog

No suitable spotted frog habitat exists in or near the proposed project area, and no spotted frogs are known or thought to occur on the Manti-La Sal National Forest. Therefore, it is my determination that the proposed SUFCO 2004 Helicopter-Assisted Coal Exploration Drilling Project will have no impact on the spotted frog.

Colorado Cutthroat Trout

Colorado cutthroat trout are not found in the proposed project area, and the project would not adversely impact drainages where it is found. Therefore, it is my determination that the SUFCO 2004 Helicopter-Assisted Coal Exploration Drilling Project would have no impact on the Colorado cutthroat trout.

Bonneville Cutthroat Trout

Bonneville cutthroat trout are not found in the proposed project area, and the project would not adversely impact drainages where it is found. Therefore, it is my determination that the SUFCO 2004 Helicopter-Assisted Coal Exploration Drilling Project would have no impact on the Bonneville cutthroat trout.

WILDLIFE SPECIES: The Intermountain Regional Forester lists seven sensitive wildlife species that could occur on the Manti Division of the Manti-La Sal National Forest including: the spotted bat, townsend's big-eared bat, greater sage grouse, northern goshawk, peregrine falcon, flammulated owl and three-toed woodpecker. Determinations for each of these species follow:

Spotted Bat

The nearest suitable roost habitat is located approximately ½ mile from the project area. Activity during project implementation is not likely to disturb roosting or foraging bats; nor would it disturb roost or foraging habitat. Therefore, it is my determination that the proposed SUFCO 2004 Helicopter-Assisted Coal Exploration Drilling Project will have no impact on the spotted bat.

Townsend's big-eared bat

There is suitable Townsend's big-eared bat forage habitat in the proposed project area, and there is suitable roost habitat not far away. Activity during project implementation is not likely to disturb roosting or foraging bats; nor would it disturb roost or foraging habitat. Therefore, it is my determination that the proposed SUFCO 2004 Helicopter-Assisted Coal Exploration Drilling Project will have no impact on the Townsend's big-eared bat.

Greater Sage Grouse

The proposed project would occur outside the greater sage grouse lekking and breeding season, the project would not modify lekking or breeding habitat, and the project would not occur in brood rearing habitat. Therefore, it is my determination that the proposed SUFCO 2004 Helicopter-Assisted Coal Exploration Drilling Project will have no impact on the Greater Sage Grouse.

Northern Goshawk

Goshawks forage in fairly dense (generally greater than 40 percent canopy cover) conifer forests, and they nest in even denser stands (generally greater than 60 percent canopy cover); however the proposed project is located in fairly dry sagebrush and mohogany habitat types. Therefore, it is my determination that the proposed SUFCO 2004 Helicopter-Assisted Coal Exploration Drilling Project will have no impact on the northern goshawk.

Peregrine Falcon

The nearest known peregrine falcon eyrie is located approximately 3 ½ miles from the project area. Falcons may travel more than 18 miles from the nest site to hunt for food, however a 10 mile radius around the nest is an average hunting area, with 80% of foraging occurring within a mile of the nest. Nesting peregrine falcons may forage in the vicinity of the proposed project. Project implementation would not occur during the peregrine nesting period, and would only temporarily impact localized areas within potential forage habitat. Therefore, it is my determination that the proposed SUFCO 2004 Helicopter-Assisted Coal Exploration Drilling Project will have no impact on the peregrine falcon.

Flammulated Owl

Flammulated owls prefer mature ponderosa pine/Douglas fir forests with open canopies, but they can be found in second growth ponderosa pine, aspen and mixed conifer forests that contain a ponderosa pine component. The proposed project is located in fairly dry sagebrush and mohogany habitats. Therefore, it is my determination that the proposed SUFCO 2004 Helicopter-Assisted Coal Exploration Drilling Project will have no impact on the flammulated Owl.

Three-toed Woodpecker

Three-toed woodpeckers are found in northern coniferous and mixed forest types up to 9,000 feet elevation. Forests containing spruce, grand fir, ponderosa pine, tamarack, and lodgepole pine are used. Nests may be found in spruce, tamarack, pine, cedar, and aspen trees. The proposed project is located in fairly dry sagebrush and mohogany habitats. Therefore, it is my determination that the proposed SUFCO 2004 Helicopter-Assisted Coal Exploration Drilling Project will have no impact on the three-toed woodpecker.

VI. REFERENCES

- Atwood, D., J. Holland, R. Bolander, B. Franklin, D. E. House, L. Armstrong, K. Thorne, and L. England. 1991. Utah threatened, endangered, and sensitive plant field guide. USFS, NPS, BLM, UNHP, USFWS, EPA, Navajo Nation, and Skull Valley Goshute Tribe.
- Boschen, Nelson. 1997. Bald Eagles in Southeast Utah: 1997 Nesting Season. Moab Bureau of Land Management.
- Connelly, J.W., M.A. Schroeder, A.R. Sands, and C.E. Braun. 2000. Guidelines to manage sage grouse populations and their habitats. *Wildlife Society Bulletin*. 28(4): 967-985.
- Durrant, Stephen, D. 1952. *Mammals of Utah - Taxonomy and Distribution*. University of Kansas Publications, Museum of Natural History, Volume 6. pp. 549.
- Hasenyager, R. N. 1980. *Bats of Utah*. Utah Division of Wildlife Resources. Publication Number 80-15.
- Oliver, G.V. 2000. *The Bats of Utah: A Literature Review*. Utah Division of Wildlife Resources, Salt Lake City, Utah.
- Paige, C., and S.A. Ritter. 1999. *Birds in a Sagebrush Sea: Managing sagebrush habitats for Bird Communities*. Partners in Flight Western Working Group, Boise, Idaho.
- Parrish, J.R., F.P. Howe, R.E. Norvell. 2002. *Utah Partners in Flight Avian Conservation Strategy Version 2.0*. Utah Partners in Flight Program, Utah Division of Wildlife Resources, 1594 West North Temple, Salt Lake City, Utah 84116, UDWR Publication Number 02-27.
- Perkins, J.M. and J.R. Peterson. 1997. *Bat survey for the SUFCO Mine, Emery County, Utah*. 8pp.

- Pierson, E.D., M.C. Wackenhut, J.S. Altenbach, P. Bradley, P. Call, D.L. Genter, C.E.Harris, B. L. Keller, B. Lengus, L. Lewis, B. Luce, K.W. Navo, J.M. Perkins, S. Smith, and L. Welch. 1999. Species Conservation Assessment and Strategy for Townsend's Big-eared Bat (*Corynorhinus townsendii townsendii* and *Corynorhinus townsendii pallescens*). Idaho Conservation Effort, Idaho Department of Fish and Game, Boise, Idaho.
- Sherwin, R.E., D.S. Rogers, and C.A. Johansson. 1997. Assessment of spotted bat (*Euderma maculatum*) and Townsend's big-eared bat (*Corynorhinus townsendii*) in the proposed Cottonwood Canyon lease area. Manti La Sal National Forest, Emery County, Utah. Conducted for Energy West Mining Co. 18pp + append.
- Spahr, R.L., L. Armstrong, D. Atwood, and M. Rath. 1991. Threatened, Endangered, and Sensitive Species of the Intermountain Region. USDA Forest Service Fisheries and Wildlife Management Intermountain Region, Ogden, Utah.
- Stalmaster, M.V. 1987. The bald eagle. Universe Books, New York, 227 pp.
- Toone, R.A. 1994. General Inventory for Bats in the Abajo and La Sal Mountains, Manti-La Sal National Forest, with Emphasis on the Spotted Bat (*Euderma maculatum*) and the Townsend's Big-eared Bat (*Plecotus townsendii*). Heritage Program Utah Department of Natural Resources, Salt Lake City, Utah.
- USDA, U.S. Forest Service. 1986. Manti-La Sal National Forest Land and Resource Management Plan: Forest Plan Amendment, Appendix A. MLSNF, Price, Utah.
- USDI, U.S. Fish and Wildlife Service. 2002a. Bonytail (*Gila elegans*) Recovery Goals: Amendment and Supplement to the Bonytail Chub Recovery Plan. U.S. Fish and Wildlife Service, Mountain-Prairie Region (6), Denver, Colorado.
- USDI, U.S. Fish and Wildlife Service. 2002b. Humback Chub (*Gila cypha*) Recovery Goals: Amendment and Supplement to the Humback Chub Recovery Plan. U.S. Fish and Wildlife Service, Mountain-Prairie Region (6), Denver, Colorado.
- USDI, U.S. Fish and Wildlife Service. 2002c. Colorado Pikeminnow (*Ptychocheilus lucius*) Recovery Goals: Amendment and Supplement to the Colorado Squawfish Recovery Plan. U.S. Fish and Wildlife Service, Mountain-Prairie Region (6), Denver, Colorado.
- USDI, U.S. Fish and Wildlife Service. 2002d. Razorback sucker (*Xyrauchen texanus*) Recovery Goals: Amendment and Supplement to the Razorback Sucker Recovery Plan. U.S. Fish and Wildlife Service, Mountain-Prairie Region (6), Denver, Colorado.

- USDI, U.S. Fish and Wildlife Service. 2002e. Utah Field Office Guidelines for Raptor Protection from Human and Land Use Disturbances. Utah Field Office, Salt Lake City.
- USDI, Fish and Wildlife Service. 2001a. Final designation of critical habitat for the Mexican spotted owl. Federal Register 66(22): 8530-8553.
- USDI, U.S. Fish and Wildlife Service. 2001b. Status Review for Bonneville Cutthroat Trout (*Oncorhynchus clarki utah*). Regions 1 and 6 Portland, Oregon and Denver, Colorado.
- Watkins, L. C. 1977. Mammalian Species: *Euderma maculatum*. The American Society of Mammalogists. 77:1-4.
- Welsh, S.L., N.D. Atwood, S. Goodrich, L.C. Higgens. 1987. A Utah Flora. Brigham Young University, Provo, Utah.



136 East South Temple, Suite 910
Salt Lake City Utah, 84111
Tel 801-364-1064 Fax 801-364-2021
www.tetrattech.com

Technical Memorandum

To: Mark Bunnell
Company: Bowie Resource Partners LLC
SUFCO Mine
597 South SR 24
Salina, UT 84654

From: Tetra Tech:
Mike Egan, Project Manager
Justin DeCaro, Wildlife Biologist
James Hart, Wildlife Biologist

CC: Kreig Rasmussen, USDA Forest Service
Date: 06/10/2014

Re: Biological Surveys for the Proposed 2014 Exploration Drill Holes
Project: 114-520226

Mr. Bunnell:

On behalf of Bowie Resource Partners LLC (dba Canyon Fuel Company, LLC), Tetra Tech Inc. (Tetra Tech) conducted biological surveys in support of the 2014 SUFCO Mine proposed drilling exploration project located south of White Mountain in Sevier County. The project would occur within Skumpah Canyon on land administered by the U.S. Department of Agriculture, Forest Service (USFS), Fishlake National Forest (FLNF) and Manti-La Sal National Forest (MLNF) in Sections 13, 14, 15, 20, 21, 22, 27, 28, 29, 30, 31, 32 and 33, Township 21 South, Range 4 East. The project features consist of eighteen drill holes; temporary water tanks and pumps; temporary waterlines; helicopter flight paths; and two helicopter landing zone/staging areas (Figure 1). Two biological surveys were conducted, the first from May 19 to May 22, 2014 and the second from June 2 through June 5, 2014.

Prior to conducting surveys, Tetra Tech biologist Justin DeCaro consulted Mr. Kreig Rasmussen of the USFS for specific survey requirements for this project. In particular, Mr. Rasmussen requested that Tetra Tech biologists' thoroughly survey suitable greater sage-grouse (*Centrocercus urophasianus*) habitat near disturbance features. He stated that there is suitable habitat for the sage-grouse on the north end of the FLNF along the boundary with the Manti-La Sal National Forest, but that the USFS has not located any sage-grouse or identified signs that sage-grouse are using the area. He observed sage grouse this spring, approximately 1 mile east of Quitchupah Canyon towards Wildcat Knolls. The biological surveys conducted for this project include:

- Two broadcast vocalization surveys for northern goshawk (*Accipiter gentilis*)
- Raptor nest survey
- Visual encounter surveys for:
 1. Greater Sage Grouse
 2. Pygmy rabbit (*Brachylagus idahoensis*)
 3. Western boreal toad (*Bufo boreas*)

4. U.S. Fish and Wildlife Service (USFWS) federally listed plant, wildlife, and aquatic species for Sevier County
5. Other USFS Sensitive Species
6. USFS Management Indicator Species (MIS)
7. Utah Division of Natural Resources (UDWR) Species of Concern (SPC) that have potential to occur in Sevier County
8. Migratory Bird Species

SURVEY METHODS

Northern Goshawk Broadcast Vocalization Surveys

Northern goshawk broadcast vocalization surveys were conducted following the USFS protocol (Woodbridge and Hargis 2006). At each calling station, a northern goshawk alarm call was broadcast for 10 seconds, followed by 30 seconds of attentive listening and watching. Biologists would then repeat this sequence two more times, rotating the broadcaster 120 degrees between each call. Calls were broadcast on calm days no earlier than one-half hour before sunrise and no later than one-half hour after sunset.

The survey area was a 0.5-mile buffer around all project features associated with the exploration area. Survey calling points were located within suitable habitat along flight lines and at each project feature, at least 200 meters apart. The area was surveyed two times, with visits 11 days apart. Photos were taken at each drill site to document habitat conditions.

Raptor Nest Surveys

Raptor nest surveys were conducted following guidance from the 2010 State of Utah Department of Natural Resources, Division of Oil, Gas, and Mining *Raptor Survey Guidelines* (DOGM 2010). All suitable nesting habitat identified in the exploration area and the accompanying 0.5-mile buffer were surveyed (including stands of conifer, aspen (*Populus tremuloides*) or other trees; and cliff areas). Locations of any raptor nests encountered during the surveys were recorded using a Global Positioning System (GPS) unit. Photos were acquired if possible. Biologists also recorded any individual raptors that were observed while walking and driving throughout the project area. Spotting scopes and binoculars were used to identify raptor species, verify the presence of nests and determine nest condition and activity.

Visual Encounter Surveys

Surveys were conducted throughout the exploration area and the accompanying 0.5-mile buffer for the following:

- Suitable sagebrush habitat was surveyed for visual presence and signs of greater sage-grouse including scat and feathers.
- Suitable sagebrush habitat was also surveyed for visual presence and signs of pygmy rabbits including scat and burrows.
- All streams and wetlands occurring in the survey area were surveyed for western boreal toads by slowly walking around the perimeter of the feature.
- Visual encounter surveys for USFWS federally listed species, other USFS Sensitive, USFS MIS's, UDWR SPC, all migratory bird species, and any other incidental species were also conducted in conjunction with all of the above surveys. The locations of any of these species encountered during the survey were recorded.

RESULTS: HABITAT ASSESSMENT

Drill Sites

- Drill Site A-14: Located on top of a ridgeline. Vegetation is dominated by grass, snowberry (*Symphoricarpos albus*), aspen (*Populus tremuloides*), sagebrush (*Artemisia tridentata*), and Gambel oak (*Quercus gambelii*).
- Drill Site B-14: Located on top of small, open ridge. Vegetation consists of sparse sagebrush surrounded by serviceberry (*Amelanchier alnifolia*), Gambel oak, and limber pine (*Pinus flexilis*).
- Drill Site C-14: Located on top of a ridge. Vegetation consists of sparse sagebrush, mountain mahogany (*Cercocarpus montanus*), antelope bitterbrush (*Purshia tridentata*), and limber pine.
- Drill Site D-14: Located on top of a ridge. Vegetation consists of patchy Gambel oak, antelope bitterbrush and sparse Rocky Mountain juniper (*Juniperus scopulorum*).
- Drill Site E-14: Located on top of ridge with a northern aspect. Vegetation consists of sagebrush, scattered serviceberry, and scattered limber pine.
- Drill Site F-14: Located on bottom of slope near valley floor. Vegetation consists of sparse sagebrush intermixed with grass species. Drill site surrounded by small, scattered stands of aspen. Small, flowing creek approximately 100 meters west/northwest of drill site location.
- Drill Site G-14: Located in the bottom of drainage. Vegetation consists of sagebrush, grass species, and scattered Gambel oak.
- Drill Site H-14: Located on bottom of slope near the valley floor. Vegetation consists of sparse sagebrush intermixed with grass species, and mountain mahogany.
- Drill Site I-14: Located on bottom of slope near the valley floor. Vegetation consists sparse sagebrush intermixed with grass species.
- Drill Site J-14: Located on bottom of slope near the valley floor. Vegetation consists of sparse sagebrush intermixed with grass species, and sparse Rocky Mountain juniper.
- Drill Site K-14: Located on a south-west aspect, near the top of a ridge. Vegetation consists of sparse sagebrush, Gambel oak, antelope bitterbrush, and serviceberry. A small, isolated stand of aspen located approximately 100 meters to the west of drill site location.
- Drill Site L-14: Located on south-west aspect, on top of ridge. Vegetation consists of sparse sagebrush, serviceberry, and mountain mahogany.
- Drill Site M-14: Located in bottom of small, narrow drainage approximately 50 meters from flowing creek. Vegetation consists of sagebrush intermixed with grass species in the bottom of drainage. South facing slope composed of juniper species, mountain

mahogany and large, scattered boulders. North facing slope composed of aspen/mixed-conifer species.

- Drill Site N-14: Located near bottom of south facing slope. Vegetation consists of sagebrush, scattered Gambel oak, and serviceberry. There are a few small, isolated stands of aspen scattered in the drainage. Drainage has a small, intermittent stream channel with scattered willows (*Salix* spp.) present.
- Drill Site O-14: Located in bottom of drainage. Vegetation consists of sagebrush intermixed with grass species. Small stock pond with water located approximately 200 meters west of drill site location.
- Drill Site P-14: Located on a small saddle on top of a ridge. Vegetation consists of sparse sagebrush intermixed with sparse grass species.
- Drill Site Q-14: Located in small, open meadow on north-east aspect. Vegetation consists of sagebrush, snowberry, scattered aspen, scattered limber pine, and scattered Douglas-fir (*Pseudotsuga menziesii*).
- Drill Site R-14: Located in bottom of narrow drainage. Vegetation consists of sagebrush throughout drainage bottom. South facing slope composed of Gambel oak, and scattered pinyon-juniper. North facing slope composed of aspen/mixed-conifer species.

Flight Paths, Helicopter Staging Area, Water Tanks, and Waterlines

Flight paths between individual drill sites and the staging areas are located in Section 13, 14, 15, 20, 21, 22, 27, 28, 29, 30, 31, 32, 33, and T21S R4E on the FLNF and MLNF. Habitat within the flight paths consists of mixed conifer, aspen/mixed-conifer, aspen, Gambel oak woodland, pinyon-juniper woodland, montane shrubland, and sagebrush shrubland.

The helicopter staging areas are located in T21S R4E Section 13 and 33. Habitat around the staging area consists of sagebrush shrubland. USFS Road 007 is within the direct vicinity of the both staging areas.

RESULTS: SURVEY RESULTS

The weather on May 19th through May 22nd, 2014 was between 55-70 degrees Fahrenheit, with clear skies becoming partly cloudy in the afternoon, and variable winds of zero to five miles per hour (mph). Weather during the second visit on June 2nd through June 5th, 2014 was approximately between 60-75 degrees Fahrenheit, with clear skies, and variable winds of zero to fifteen mph.

Northern Goshawk Survey Results

The ability to detect northern goshawk or other raptor nests was excellent because a majority of the aspen trees were not leafed out during the first visit. No northern goshawks or northern goshawk nests were detected within the survey area. The majority of the habitat in the survey area is not suitable for goshawk nesting due to the lack of expansive, mature stands of timber. The most suitable northern goshawk habitat within the 2014 exploration area is the mixed conifer, aspen/mixed conifer stands in the area north and northwest of drill holes A-14, B-14, E-14, Q-14 and R-14.

Raptor Nest Survey Results

No new raptor nests were identified during the survey. A Cooper's hawk (*Accipiter cooperii*) and a pair of Golden Eagles were the only raptors observed within the survey area. The Golden Eagle pair was observed approximately 100 meters north of G-14. They appeared to be foraging. They were not exhibiting territorial behavior, and no nest was detected. The Cooper's hawk was observed approximately 200 meters west of R-14. It was not exhibiting territorial behavior, and no nest was detected.

Visual Encounter Survey Results

No greater sage-grouse, pygmy rabbit, or boreal toad or their sign were observed within the project area. Skutumpah creek was surveyed for boreal toad, but none were detected. Surveys for boreal toad were also conducted at Skutumpah spring, Lizonbee springs, and around Skutumpah reservoir. No detections were made at these locations.

No USFWS threatened, endangered, candidate, or proposed species; other USFS Sensitive; other USFS MIS's; or UDWR SPC species were documented within the project area. Table 1 lists migratory bird species observed within the project area during the 2014 surveys. Other wildlife species observed during the surveys include mule deer (*Odocoileus hemionus*) and elk (*Cervus canadensis*).

If you have any questions regarding this memo, please contact Mike Egan at 801-364-1064 extension 203.

Sincerely,
Tetra Tech, Inc.



Michael Egan
Sr. Project Manager

Attachments: Figure 1
Table 1


REFERENCES

Division of Oil, Gas and Mining (DOGM). 2010. *Raptor Survey Guidelines*. March, 2010. 23 pp.

Woodbridge, B. and C.D. Hargis. 2006. *Northern Goshawk Inventory and Monitoring Technical Guide*. Gen. Tech. Rep. WO-71. Washington, DC: U.S. Department of Agriculture, Forest Service. 80 pp.

Table 1: Migratory Bird Species Observed Within the SUFCO 2014 Exploration Drilling Project Area, May-June 2014

American Kestrel (<i>Falco sparverius</i>)
American Robin (<i>Turdus migratorius</i>)
Black-billed Magpie (<i>Pica hudsonia</i>)
Black-capped Chickadee (<i>Parus atricapillus</i>)
Black-headed Grosbeak (<i>Pheucticus melanocephalus</i>)
Brewer's Sparrow (<i>Spizella breweri</i>)
Cassin's Finch (<i>Carpodacus cassinii</i>)
Cassin's Vireo (<i>Vireo cassinii</i>)
Chipping Sparrow (<i>Spizella passerina</i>)
Clark's Nutcracker (<i>Nucifraga columbiana</i>)
Common Raven (<i>Corvus corax</i>)
Cooper's Hawk (<i>Accipiter cooperii</i>)
Dark-eyed Junco (<i>Junco hyemalis</i>)
Dusky Flycatcher (<i>Empidonax oberholseri</i>)
Gray Flycatcher (<i>Empidonax wrightii</i>)
Green-tailed Towhee (<i>Pipilo chlorurus</i>)
Hairy Woodpecker (<i>Picoides villosus</i>)
House Wren (<i>Troglodytes aedon</i>)
MacGillivray's Warbler (<i>Geothlypis tolmiei</i>)
Mountain Bluebird (<i>Sialia currucoides</i>)
Mountain Chickadee (<i>Parus gambeli</i>)
Northern Flicker (<i>Colaptes auratus</i>)
Orange-crowned Warbler (<i>Oreothlypis celata</i>)
Plumbeous Vireo (<i>Vireo plumbeus</i>)
Red-breasted Nuthatch (<i>Sitta canadensis</i>)
Red-naped Sapsucker (<i>Sphyrapicus nuchalis</i>)
Red-tailed Hawk (<i>Buteo jamaicensis</i>)
Ruby-crowned Kinglet (<i>Regulus calendula</i>)
Spotted Towhee (<i>Pipilo maculatus</i>)
Stellar's Jay (<i>Cyanocitta stelleri</i>)
Turkey Vulture (<i>Cathartes aura</i>)
Vesper Sparrow (<i>Pooecetes gramineus</i>)
Virginia's Warbler (<i>Oreothlypis virginiae</i>)
Warbling Vireo (<i>Vireo gilvus</i>)
Western Wood-pewee (<i>Contopus sordidulus</i>)
White-crowned Sparrow (<i>Zonotrichia leucophrys</i>)
Yellow-rumped Warbler (<i>Dendroica coronata</i>)
Yellow Warbler (<i>Setophaga petechia</i>)





136 East South Temple, Suite 910
 Salt Lake City, Utah, 84111
 Tel: 801-364-1064 Fax: 801-364-2021
 www.tetrattech.com

Technical Memorandum

To:	Mark Bunnell	From:	Tetra Tech:
Company:	Bowie Resource Partners LLC SUFCO Mine 597 South SR 24 Salina, UT 84854		Mike Egan, Project Manager Justin DeCaro, Wildlife Biologist
CC:	Kreig Rasmussen, USDA Forest Service	Date:	07/22/2014
Re:	Addendum - Biological Surveys for the Proposed 2014 Exploration Program	Project:	114-520226

Mr. Bunnell:

On behalf of Bowie Resource Partners LLC (dba Canyon Fuel Company, LLC), Tetra Tech Inc. (Tetra Tech) conducted biological surveys at an additional drill hole (S-14) in support of the 2014 SUFCO Mine proposed drilling exploration project. This report is an addendum to the original report submitted on June 20, 2014.

The additional drill hole S-14 is located on the Manti La Sal National Forest, in Section 18 Township 21 South, Range 5 East. This drill hole is near holes M-14, N-14, O-14, and P-14, and would be supported by the same staging area/landing zone located at the north fork of Quitchupah Creek (**Figure 1 (revised)**). The drill hole and a surrounding half-mile buffer was surveyed on July 20, 2014 for raptor nests. In addition, broadcast vocalization surveys were conducted for northern goshawk (*Accipiter gentilis*) and visual encounter surveys were conducted for greater sage-grouse (*Centrocercus urophasianus*), pygmy rabbit (*Brachylagus idahoensis*); federally threatened, endangered, candidate or proposed species; Forest Service Sensitive and Management Indicator Species (MIS); and state listed species. Migratory birds and other wildlife observed during the survey were also documented. Methods were the same used in the previous surveys (see memo from June 20, 2014).

RESULTS

Habitat Assessment

Drill Site S-14: Located near top of a ridgeline on south/southwest facing slope. Vegetation consists of Rocky Mountain juniper, pinyon pine, Gambel oak, limber pine, serviceberry, mountain mahogany, and sagebrush.

Wildlife Survey

No northern goshawk or other raptor nests were detected during the survey. No federally threatened, endangered, candidate or proposed species were documented. No Forest Service Sensitive or MIS or state-listed species were documented.

One adult red-tailed hawk was observed soaring approximately 200m to the west of proposed drill site.

The following migratory birds were recorded during the survey:

- Clark's Nutcracker
- Mountain Bluebird
- Black-billed Magpie
- Spotted Towhee
- Green-tailed Towhee
- Black-capped chickadee
- Chipping Sparrow
- Dark-eyed Junco

If you have any questions regarding the results of the survey or this report, please contact Mike Egan at 801-364-1064 extension 203.

Sincerely,
Tetra Tech, Inc.



Michael Egan
Sr. Project Manager

Attachments: **Figure 1 (revised)**

APPENDIX D

SUFCO RAPTOR SURVEY 2014

(SUBMITTED FOR PLACEMENT IN
CONFIDENTIAL FILE)

APPENDIX E

**CULTURAL RESOURCE INVENTORY OF 13
DRILL LOCATIONS, SUFCO MINE 2015 DRILLING,
BIG RIDGE AND NORTH FORK QUITCHUPAH AREAS,
FISHLAKE AND MANTI-LA SAL NATIONAL FOREST,
SEVIER AND SANPETE COUNTIES, UTAH
U-15-EZ-0056f**

**(SUBMITTED FOR PLACEMENT IN
CONFIDENTIAL FILE)**

